

Adjusted odds ratios — SHAP-informed confirmatory model, by age cohort
union of top-15 (50+) / top-15 (65+) SHAP-selected predictors
BOTH COHORTS (11)
Self-rated oral health
Current smoking rate indicator
Toothache in past year
Dong/Eup-Myeon classification
Diabetes (diagnosed or HbA1c ≥ 6.5%)
Education level
Daily brushing frequency (count)
Age
Income quartile
Frequency of fruit intake, past year
Daily total dietary fiber
50+ ONLY (4)
Region: Gyeongbuk
Housing type: Apartment
Uric acid
Estimated kidney function (eGFR, CKD-EPI)
65+ ONLY (4)
Brushed teeth yesterday
0.08 (0.04–0.15)
Region: Gyeongnam
Final systolic/diastolic BP
Daily total sugar
0.5
0.6
0.7
0.8
0.9
1
1.25
1.5
2
2.5
Adjusted odds ratio (95% CI), log scale
50+ (n=12,524, prevalence 25.9%)
65+ (n=6,471, prevalence 41.4%)
Filled = p < 0.05
Open = not significant
One marker per row = selected into only that cohort's SHAP-informed model | arrow markers = point estimate off the shared axis, exact aOR (95% CI) printed alongside