

Artificial Intelligence Foundation – JC3001

Lecture 34: Stochastic Planning - I

Prof. Aladdin Ayesh (aladdin.ayesh@abdn.ac.uk)

Dr. Binod Bhattarai (binod.bhattarai@abdn.ac.uk)

Dr. Gideon Ogunniye, (g.ogunniye@abdn.ac.uk)

October 2025

Material adapted from:

Russell and Norvig (AIMA Book): Chapter 17

Sutton and Barto (Reinforcement Learning: An Introduction 2nd ed.)

Sebastian Thrun (Stanford University / Udacity)

Course Progression



- Part 1: Introduction
 - ① Introduction to AI ✓
 - ② Agents ✓
- Part 2: Problem-solving
 - ① Search 1: Uninformed Search ✓
 - ② Search 2: Heuristic Search ✓
 - ③ Search 3: Local Search ✓
 - ④ Search 4: Adversarial Search ✓
- Part 3: Reasoning and Uncertainty
 - ① Reasoning 1: Constraint Satisfaction ✓
 - ② Reasoning 2: Logic and Inference ✓
 - ③ Probabilistic Reasoning 1: BNs ✓
 - ④ Probabilistic Reasoning 2: HMMs ✓
- Part 4: Planning
 - ① Planning 1: Intro and Formalism ✓
 - ② Planning 2: Algorithms & Heuristics ✓
 - ③ Planning 3: Hierarchical Planning ✓
 - ④ **Planning 4: Stochastic Planning**
- Part 5: Learning
 - ① Learning 1: Intro to ML
 - ② Learning 2: Regression
 - ③ Learning 3: Neural Networks
 - ④ Learning 4: Reinforcement Learning
- Part 6: Conclusion
 - ① Ethical Issues in AI
 - ② Conclusions and Discussion

Objectives

- Planning with Utilities
- Stochastic Planning Formalisms
- Dynamic Programming Algorithms for Stochastic Planning



Outline

1 Recap

► Recap

► Utilities and Expectation


Recap

1 Recap

- We know how to get from a current state to a goal state
 - Via domain specific search
 - Via domain independent planning
- We also know how to calculate the probability of future events via probability theory
- But, how can we get the agent to decide between different actions, particularly in the face of uncertainty?

Planning Under Uncertainty

1 Recap



Recap

1 Recap

	Deterministic	Stochastic
Fully Observable		
Partially Observable		

Recap

1 Recap

	Deterministic	Stochastic
Fully Observable	A*, DFS BFS, GP	
Partially Observable		

Recap

1 Recap

	Deterministic	Stochastic
Fully Observable	A*, DFS BFS, GP	MDP
Partially Observable		

Recap

1 Recap

	Deterministic	Stochastic
Fully Observable	A*, DFS BFS, GP	MDP
Partially Observable		POMDP



Outline

2 Utilities and Expectation

► Recap

► Utilities and Expectation

How Good is an action

2 Utilities and Expectation

- Consider an agent capable of planning and executing different actions to achieve different goals
- Example you can either:
 - Stay home and study
 - Go out and have fun
- Which should you do, and why?

Rationality

2 Utilities and Expectation

- A rational agent should pursue the goal that is, in some sense, “best for it”
- How can we identify whether the achievement of one goal is “better” for the agent than another?

Rationality

2 Utilities and Expectation

- A rational agent should pursue the goal that is, in some sense, “best for it”
- How can we identify whether the achievement of one goal is “better” for the agent than another?
- Utility theory
 - We capture preferences between world states via a utility function $U(S)$

$$U : States \rightarrow \mathbb{R}$$

- In other words, we assign a single number to a state to express the **desirability** of the state

Rationality

2 Utilities and Expectation

- A rational agent should pursue the goal that is, in some sense, “best for it”
- How can we identify whether the achievement of one goal is “better” for the agent than another?
- Utility theory
 - We capture preferences between world states via a utility function $U(S)$

$$U : States \rightarrow \mathbb{R}$$

- In other words, we assign a single number to a state to express the **desirability** of the state
- Winning the lottery $U(w) = 1000$

Rationality

2 Utilities and Expectation

- A rational agent should pursue the goal that is, in some sense, “best for it”
- How can we identify whether the achievement of one goal is “better” for the agent than another?
- Utility theory
 - We capture preferences between world states via a utility function $U(S)$

$$U : States \rightarrow \mathbb{R}$$

- In other words, we assign a single number to a state to express the **desirability** of the state
- Winning the lottery $U(w) = 1000$ Failing AI: $U(f) = -10$

Rationality

2 Utilities and Expectation

- A rational agent should pursue the goal that is, in some sense, “best for it”
- How can we identify whether the achievement of one goal is “better” for the agent than another?
- Utility theory
 - We capture preferences between world states via a utility function $U(S)$

$$U : States \rightarrow \mathbb{R}$$

- In other words, we assign a single number to a state to express the **desirability** of the state
Winning the lottery $U(w) = 1000$ Failing AI: $U(f) = -10$
- But what if actions have uncertain effects?
Buying a lottery ticket does not give us a utility of 1000

Expected Utility

2 Utilities and Expectation

- A nondeterministic action A could lead to several possible outcomes, e.g. $Result_1(A), Result_2(A), \dots$
 $Result(BuyTicket) \in \{win, \neg win\}$
- We can associate a probability with the different outcomes

$$P(Result_i(A) \mid Do(A), E)$$

- E captures the evidence we have about the world
 - $Do(A)$ is the proposition that action A is executed in the current state
- The expected utility of executing A given E is then

$$\mathbb{E}[U(A) \mid E] = \sum_i P(Result_i(A) \mid Do(A), E)U(Result_i(A))$$

- Principle of maximum expected utility (MEU):
A rational agent should choose the action that maximises its expected utility

The Expectation Operator

2 Utilities and Expectation

More generally

- We can use the expectation operator ($\mathbb{E}[X]$) for **any random variable** X

$$\mathbb{E}[X] \doteq \sum_{x \in X} P(x)x$$

- Example:
Expected value of a six sided die:

The Expectation Operator

2 Utilities and Expectation

More generally

- We can use the expectation operator ($\mathbb{E}[X]$) for **any random variable** X

$$\mathbb{E}[X] \doteq \sum_{x \in X} P(x)x$$

- Example:

Expected value of a six sided die:

x	$P(X=x)$	$x * P(X=x)$
1	1/6	1/6
2	1/6	2/6
3	1/6	3/6
4	1/6	4/6
5	1/6	5/6
6	1/6	6/6
Total		21/6 = 3.5

Maximum Expected Utility

2 Utilities and Expectation

- An agent following the principle of MEU is rational, acting optimally in any situation

Maximum Expected Utility

2 Utilities and Expectation

- An agent following the principle of MEU is rational, acting optimally in any situation
- But choosing the best sequence of actions requires enumerating all action sequences; infeasible for long sequences

Maximum Expected Utility

2 Utilities and Expectation

- An agent following the principle of MEU is rational, acting optimally in any situation
- But choosing the best sequence of actions requires enumerating all action sequences; infeasible for long sequences
- Computations are often too expensive:
 - $P(Result_i(A) \mid Do(A), E)$ requires a complete causal model of the world. Even if all information is available, this is computationally intractable
 - Computing $U(Result_i(A))$ requires searching or planning as we do not know how good a state is until we know what goals we can reach from it (and how expensive they are)

Maximum Expected Utility

2 Utilities and Expectation

- MEU is no silver bullet for the problems of AI
 - But does give us a nice framework for building an agent,

Maximum Expected Utility

2 Utilities and Expectation

- MEU is no silver bullet for the problems of AI
 - But does give us a nice framework for building an agent,
 - An MEU maximising agent will achieve the highest possible performance averaged over all environments in which the agent can be placed

Maximum Expected Utility

2 Utilities and Expectation

- MEU is no silver bullet for the problems of AI
 - But does give us a nice framework for building an agent,
 - An MEU maximising agent will achieve the highest possible performance averaged over all environments in which the agent can be placed
 - This moves from a performance measure in the external environment, to an internal criteria of utility maximisation

Lotteries

2 Utilities and Expectation

- A lottery is a probability distribution over a set of actual outcomes

Lotteries

2 Utilities and Expectation

- A lottery is a probability distribution over a set of actual outcomes
- A lottery L with possible outcomes C_1, \dots, C_n and associated probabilities p_1, \dots, p_n is written as

$$L = [p_1, C_1; \dots p_n, C_n]$$

Lotteries

2 Utilities and Expectation

- A lottery is a probability distribution over a set of actual outcomes
- A lottery L with possible outcomes C_1, \dots, C_n and associated probabilities p_1, \dots, p_n is written as

$$L = [p_1, C_1; \dots p_n, C_n]$$

- Each outcome C_i is an atomic state, or another lottery

Lotteries

2 Utilities and Expectation

- A lottery is a probability distribution over a set of actual outcomes
- A lottery L with possible outcomes C_1, \dots, C_n and associated probabilities p_1, \dots, p_n is written as

$$L = [p_1, C_1; \dots p_n, C_n]$$

- Each outcome C_i is an atomic state, or another lottery
- Example: There is a 30% likelihood I will go out tonight.
If I do so, there is a 60% likelihood I will do badly in my test.
If I do not go out, I will do well in my test with a 90% likelihood

$$L = [0.3, [0.6, f; 0.4, p]; 0.7, [0.9, p; 0.1, f]]$$

Lotteries

2 Utilities and Expectation

- A lottery is a probability distribution over a set of actual outcomes
- A lottery L with possible outcomes C_1, \dots, C_n and associated probabilities p_1, \dots, p_n is written as

$$L = [p_1, C_1; \dots p_n, C_n]$$

- Each outcome C_i is an atomic state, or another lottery
- Example: There is a 30% likelihood I will go out tonight.
If I do so, there is a 60% likelihood I will do badly in my test.
If I do not go out, I will do well in my test with a 90% likelihood

$$L = [0.3, [0.6, f; 0.4, p]; 0.7, [0.9, p; 0.1, f]]$$

- Our goal is to understand the relationships between preferences between lotteries and preferences between states

Properties of Preferences

2 Utilities and Expectation

- Preferences should be transitive

$$A \succ B, B \succ C \rightarrow A \succ C$$

- Proof:
 - Suppose $A \succ B \succ C \succ A$
 - And assume A, B, C are exchangeable goods
 - Then if the agent has A , we could offer it C for A and a bit of money
 - Since the agent prefers C , they would agree to this deal
 - Repeat, trading C for B , B for A and we are back at the start, with the agent having less money
 - Clearly the agent has not acted rationally

Utility

2 Utilities and Expectation

- Utility Principle: If an agent's preferences obey the axioms of utility, then there is a real-valued function U that operates on states such that $U(A) > U(B)$ if and only if A is preferred to B , and $U(A) = U(B)$ if and only if the agent is indifferent between A and B

$$U(A) > U(B) \leftrightarrow A \succ B \quad U(A) = U(B) \leftrightarrow A \sim B$$

- Maximum Expected Utility Principle: The utility of a lottery is the sum of the probability of each outcome times the utility of that outcome

$$U([p_1, S_1; \dots, p_n, S_n]) = \sum_i p_i U(S_i)$$

Utility

2 Utilities and Expectation

- Utility Principle: If an agent's preferences obey the axioms of utility, then there is a real-valued function U that operates on states such that $U(A) > U(B)$ if and only if A is preferred to B , and $U(A) = U(B)$ if and only if the agent is indifferent between A and B

$$U(A) > U(B) \leftrightarrow A \succ B \quad U(A) = U(B) \leftrightarrow A \sim B$$

- Maximum Expected Utility Principle: The utility of a lottery is the sum of the probability of each outcome times the utility of that outcome

$$U([p_1, S_1; \dots, p_n, S_n]) = \sum_i p_i U(S_i)$$

- Given probabilities and utilities of outcomes, we can determine the utility of a compound lottery

- Utility Principle: If an agent's preferences obey the axioms of utility, then there is a real-valued function U that operates on states such that $U(A) > U(B)$ if and only if A is preferred to B , and $U(A) = U(B)$ if and only if the agent is indifferent between A and B

$$U(A) > U(B) \leftrightarrow A \succ B \quad U(A) = U(B) \leftrightarrow A \sim B$$

- Maximum Expected Utility Principle: The utility of a lottery is the sum of the probability of each outcome times the utility of that outcome

$$U([p_1, S_1; \dots, p_n, S_n]) = \sum_i p_i U(S_i)$$

- Given probabilities and utilities of outcomes, we can determine the utility of a compound lottery
- Since outcomes of nondeterministic actions are lotteries, we obtain the MEU decision rule

Utility

2 Utilities and Expectation

- Utility Principle: If an agent's preferences obey the axioms of utility, then there is a real-valued function U that operates on states such that $U(A) > U(B)$ if and only if A is preferred to B , and $U(A) = U(B)$ if and only if the agent is indifferent between A and B

$$U(A) > U(B) \leftrightarrow A \succ B \quad U(A) = U(B) \leftrightarrow A \sim B$$

- Maximum Expected Utility Principle: The utility of a lottery is the sum of the probability of each outcome times the utility of that outcome

$$U([p_1, S_1; \dots, p_n, S_n]) = \sum_i p_i U(S_i)$$

- Given probabilities and utilities of outcomes, we can determine the utility of a compound lottery
- Since outcomes of nondeterministic actions are lotteries, we obtain the MEU decision rule

- We now know what a utility function is
- But without structure, there is little we can do with it

Utility Functions

2 Utilities and Expectation

- A typical use of utility functions (stemming from economic theory) maps the amount of money an agent has to its utility
- Agents prefer having more money to less: such a preference is monotonic
- But what about lotteries about money?

Utility Functions

2 Utilities and Expectation

- A typical use of utility functions (stemming from economic theory) maps the amount of money an agent has to its utility
- Agents prefer having more money to less: such a preference is monotonic
- But what about lotteries about money?
- You are a participant in a game show. You are in the final round, and can take \$1,000,000, or gamble it on a coin flip.
If the coin lands on heads, you will win \$3,000,000, tails, you will take home nothing.
What do you do?

Utility Functions

2 Utilities and Expectation

- A typical use of utility functions (stemming from economic theory) maps the amount of money an agent has to its utility
- Agents prefer having more money to less: such a preference is monotonic
- But what about lotteries about money?
- You are a participant in a game show. You are in the final round, and can take \$1,000,000, or gamble it on a coin flip.
If the coin lands on heads, you will win \$3,000,000, tails, you will take home nothing.
What do you do?
- The expected monetary value (EMV) is \$1,500,000;
this is more than the prize
- If $\$1=1$ utility, then a rational agent would take the gamble

Utility Functions

2 Utilities and Expectation

- Instead, let us denote the state of having \$ n as S_n , and the your initial states as S_k .
Then

$$EU(flip) = 0.5U(S_k) + 0.5U(S_{k+3000000}) \quad EU(noflip) = U(S_{k+1000000})$$

- We now need to assign utilities to the different outcome states
 - Utility is typically not proportional to money:
having a 3,000,000 is **not** worth 3 times as much as having 1,000,000.

Utility Functions

2 Utilities and Expectation

- Instead, let us denote the state of having \$ n as S_n , and the your initial states as S_k .
Then

$$EU(flip) = 0.5U(S_k) + 0.5U(S_{k+3000000}) \quad EU(noflip) = U(S_{k+1000000})$$

- We now need to assign utilities to the different outcome states
 - Utility is typically not proportional to money:
having a 3,000,000 is **not** worth 3 times as much as having 1,000,000.
 - So perhaps $S_k = 5, S_{k+3000000} = 10, S_{k+1000000} = 8$

Utility Functions

2 Utilities and Expectation

- Instead, let us denote the state of having \$ n as S_n , and the your initial states as S_k .
Then

$$EU(flip) = 0.5U(S_k) + 0.5U(S_{k+3000000}) \quad EU(noflip) = U(S_{k+1000000})$$

- We now need to assign utilities to the different outcome states
 - Utility is typically not proportional to money:
having a 3,000,000 is **not** worth 3 times as much as having 1,000,000.
 - So perhaps $S_k = 5, S_{k+3000000} = 10, S_{k+1000000} = 8$
 - What are the expected utilities? $EU(flip) = ?$ $EU(noflip) = ?$

Utility Functions

2 Utilities and Expectation

- Instead, let us denote the state of having \$ n as S_n , and the your initial states as S_k .
Then

$$EU(flip) = 0.5U(S_k) + 0.5U(S_{k+3000000}) \quad EU(noflip) = U(S_{k+1000000})$$

- We now need to assign utilities to the different outcome states
 - Utility is typically not proportional to money:
having a 3,000,000 is **not** worth 3 times as much as having 1,000,000.
 - So perhaps $S_k = 5, S_{k+3000000} = 10, S_{k+1000000} = 8$
 - Then $EU(flip) = 7.5, EU(noflip) = 8$, and you would not flip the coin

Utility Functions

2 Utilities and Expectation

- Instead, let us denote the state of having \$ n as S_n , and the your initial states as S_k .
Then

$$EU(flip) = 0.5U(S_k) + 0.5U(S_{k+3000000}) \quad EU(noflip) = U(S_{k+1000000})$$

- We now need to assign utilities to the different outcome states
 - Utility is typically not proportional to money:
having a 3,000,000 is **not** worth 3 times as much as having 1,000,000.
 - So perhaps $S_k = 5, S_{k+3000000} = 10, S_{k+1000000} = 8$
 - Then $EU(flip) = 7.5, EU(noflip) = 8$, and you would not flip the coin
- Typically, for some lottery L , $U(L) < U(S_{EMV(L)})$. In other words, being given the cash without gambling is better than facing a lottery
 - Agents with this type of utility function are risk averse
 - If $U(S_{EMV(L)}) < U(L)$ the agent is risk seeking

Utility Functions

2 Utilities and Expectation

- The amount an agent will accept instead of a lottery is called the certainty equivalent of the lottery
- The difference between the expected monetary value of a lottery and its certainty equivalent is called the insurance premium
- The insurance premium is only positive if an agent is risk averse
- If a utility curve is linear (i.e. $U(S_{EMV(L)}) = U(L)$), the agent is

Utility Functions

2 Utilities and Expectation

- The amount an agent will accept instead of a lottery is called the certainty equivalent of the lottery
- The difference between the expected monetary value of a lottery and its certainty equivalent is called the insurance premium
- The insurance premium is only positive if an agent is risk averse
- If a utility curve is linear (i.e. $U(S_{EMV(L)}) = U(L)$), the agent is risk neutral
- Any affine transformation of a utility function is a utility function

To continue in the next session.