

Introduction to Markov genealogy processes

Aaron A. King

March 4, 2024


Contents

1	Context	1
1.1	Example: emerging variants	1
1.2	Phylodynamics	2
1.3	Problems of phylodynamics	3
2	Population process	4
2.1	Examples	4
2.2	Formalization	4
3	Genealogy process	6
3.1	Genealogies	6
3.2	Induced genealogy process	6
4	Pruned and obscured genealogies	8


1 Context

1.1 Example: emerging variants


Example: surveillance for emerging SARS-CoV-2 variants




nextstrain.org (Hadfield *et al.*, 2018)



nextstrain.org (Hadfield *et al.*, 2018)



$$\lambda_1 = \beta_1 \frac{I_1}{N} \quad \lambda_2 = \beta_2 \frac{I_2}{N}$$





(Mathieu *et al.*, 2020)

1.2 Phylodynamics

What is phylodynamics?



Broadly: Phylodynamics is the project of inferring *determinants of epidemic spread* using *genomic data from pathogen samples*.


In this talk: Phylodynamics means using *genomic data* to infer *stochastic dynamic transmission models*.



1.3 Problems of phylodynamics

Core problems of phylodynamics






Overview

- We show how a given population process induces a unique genealogy process.
- *Pruning* and *obscuration* project a genealogy onto observable data.

2 Population process

2.1 Examples

Population process



2.2 Formalization

Population process

- Non-explosive Markov jump process, $\mathbf{X}_t \in \mathbb{X}$, $t \in \mathbb{R}_+$: the *population process*.
- Initial-state distribution, p_0 :

$$\text{Prob}[\mathbf{X}_0 \in \mathcal{E}] = \int_{\mathcal{E}} p_0(x) dx$$

- Jump rates: $\alpha(t, x, x') = \text{rate of jump } x \rightarrow x'$

$$\alpha(t, x, x') \geq 0, \quad \int_{\mathbb{X}} \alpha(t, x, x') dx' < \infty$$

- Multiple events at each jump are allowed.

Kolmogorov forward equation (KFE): If

$$\frac{\partial w}{\partial t}(t, x) = \int w(t, x') \alpha(t, x', x) dx' - \int w(t, x) \alpha(t, x, x') dx'$$


and

$$w(0, x) = p_0(x)$$


then

$$\int_{\mathcal{E}} w(t, x) dx = \text{Prob}[\mathbf{X}_t \in \mathcal{E}].$$

KFE is sometimes called the *master equation* for \mathbf{X}_t .



$$\frac{\partial w}{\partial t}(t, x) = \int w(t, x') \alpha(t, x', x) dx' - \int w(t, x) \alpha(t, x, x') dx'$$



$$\mathbb{U} = \{\text{Trans, Prog, Recov, Wane, Sample}\}$$




$$\frac{\partial w}{\partial t}(t, x) = \sum_{u \in \mathbb{U}} \left\{ \int w(t, x') \alpha_u(t, x', x) dx' - \int w(t, x) \alpha_u(t, x, x') dx' \right\}$$

$$\begin{aligned} \frac{\partial w}{\partial t}(t, S, E, I, R) = & \frac{\beta(t)(S+1)I}{N} w(t, S+1, E-1, I, R) - \frac{\beta(t)SI}{N} w(t, S, E, I, R) \\ & + \sigma(E+1) w(t, S, E+1, I-1, R) - \sigma E w(t, S, E, I, R) \\ & + \gamma(I+1) w(t, S, E, I+1, R-1) - \gamma I w(t, S, E, I, R) \\ & + \omega(R+1) w(t, S-1, E, I, R+1) - \omega R w(t, S, E, I, R) \end{aligned}$$

3 Genealogy process


3.1 Genealogies

What is a genealogy?



3.2 Induced genealogy process

Event types




If we write

$$\alpha(t, x, x') = \sum_{u \in \mathbb{U}} \alpha_u(t, x, x'),$$

the KFE becomes

$$\frac{\partial w}{\partial t}(t, x) = \sum_u \int w(t, x') \alpha_u(t, x', x) dx' - \sum_u \int w(t, x) \alpha_u(t, x, x') dx'$$




A population process induces a genealogy process

- \mathbf{G}_t is a stochastic process on the space of genealogies.
- The map $\mathbf{X} \mapsto \mathbf{G}$ is random.
- **Key assumption:** Lineages within a deme are *exchangeable*. There is no more structure than is implied by the population process.


- Simulation code on github.com/kingaa/phylopomp
- Animations at <https://kingaa.github.io/manuals/phylopomp/vignettes/>

4 Pruned and obscured genealogies


Full genealogy



Pruned genealogy



Obscured genealogy



An obscured genealogy is specified by (T, Z) .

Summary and outstanding challenges


Summary

- A discretely structured Markov population process uniquely induces a genealogy-valued Markov process.
- The likelihood of an observed genealogy satisfies a nonlinear filtering equation, which can be efficiently computed via Feynman-Kač (sequential Monte Carlo) algorithms.
- In principle, these results liberate us to entertain models that more closely match our biological questions, without less hindrance from inference methodology.

References

- Hadfield J, Megill C, Bell SM, Huddleston J, Potter B, Callender C, Sagulenko P, Bedford T, Neher RA (2018). “Nextstrain: real-time tracking of pathogen evolution.” *Bioinformatics*, **34**(23), 4121–4123. doi: [10.1093/bioinformatics/bty407](https://doi.org/10.1093/bioinformatics/bty407).
- King AA, Nguyen D, Ionides EL (2016). “Statistical inference for partially observed Markov processes via the R package pomp.” *J Stat Softw*, **69**(12), 1–43. doi: [10.18637/jss.v069.i12](https://doi.org/10.18637/jss.v069.i12).
- Mathieu E, Ritchie H, Rodés-Guirao L, Appel C, Giattino C, Hasell J, Macdonald B, Dattani S, Beltekian D, Ortiz-Ospina E, Roser M (2020). “Coronavirus pandemic (COVID-19).” *Our World in Data* [Online resource]. URL <https://ourworldindata.org/coronavirus>.
- Vaughan TG, Leventhal GE, Rasmussen DA, Drummond AJ, Welch D, Stadler T (2019). “Estimating epidemic incidence and prevalence from genomic data.” *Mol. Biol. Evol.*, **36**, 1804–1816. doi: [10.1093/molbev/msz106](https://doi.org/10.1093/molbev/msz106).

License, acknowledgments, and links

- The materials build on [previous versions of this course and related courses](#).
- Licensed under the [Creative Commons Attribution-NonCommercial license](#). Please share and remix non-commercially, mentioning its origin. 
- Produced with R version 4.3.2 and **pomp** version 5.6.

- Compiled on March 4, 2024.

[Back to Lesson](#)

[R codes for this lesson](#)