he principles of probability help bridge the worlds of descriptive statistics and he principles of probability help or age the foundation for the probability ferential statistics. Probability principles are the foundation for the probability of ferential statistics. ferential statistics. Probability principles are tribution, the concept of mathematical expectation, and the binomial and Political Concept of mathematical expectation, and the binomial and Political Concept of mathematical expectation, and the binomial and Political Concept of mathematical expectation, and the binomial and Political Concept of mathematical expectation, and the binomial and Political Concept of mathematical expectation, and the binomial and Political Concept of mathematical expectation, and the binomial and Political Concept of mathematical expectation. distributions, topics that are discussed in Chapter 5. In this chapter, you will learn about productions to the concept of mathematical expectations and Policy distributions, topics that are discussed in Chapter 5. In this chapter, you will learn about productions to the concept of mathematical expectations are discussed in Chapter 5. ability to answer questions such as the following: What is the probability that a household is planning to purchase a large-screen HDT

- What is the probability that a household will actually purchase a large-screen HDTo
- What is the probability that a household is planning to purchase a large-screen HD
- Given that the household is planning to purchase a large-screen HDTV, what is the property of the propert
- Does knowledge of whether a household plans to purchase the television change likelihood of predicting whether the household will purchase the television?
- What is the probability that a household that purchases a large-screen HDTV will a chase a television with a faster refresh rate?
- What is the probability that a household that purchases a large-screen HDTV with faster refresh rate will also purchase a streaming media box?
- What is the probability that a household that purchases a large-screen HDTV will satisfied with the purchase?

With answers to questions such as these, you can begin to form a marketing strategy. can consider whether to target households that have indicated an intent to purchase or to cus on selling televisions that have faster refresh rates or both. You can also explore whether households that purchase large-screen HDTVs with faster refresh rates can be easily persuad to also purchase streaming media boxes.

4.1 Basic Probability Concepts

Student Tip

Remember, a probability cannot be negative or greater than 1.

What is meant by the word probability? A probability is the numerical value represent the chance, likelihood, or possibility that a particular event will occur, such as the price a stock increasing, a rainy day, a defective product, or the outcome five dots in a single II of a die. In all these instances, the probability involved is a proportion or fraction who value ranges between 0 and 1, inclusive. An event that has no chance of occurring impossible event) has a probability of 0. An event that is sure to occur (the certain even has a probability of 1.

There are three types of probability:

- · A priori
- · Empirical
- · Subjective

In the simplest case, where each outcome is equally likely, the chance of occurrence of event is defined in Equation (4.1).

PROBABILITY OF OCCURRENCE Probability of occurrence = $\frac{X}{T}$ X = number of ways in which the event occurs T = total number of possible outcomes

In a priori probability, the probability of an occurrence is based on prior knowledge of process involved. The probability of an occurrence is based on prior knowledge of process involved. the process involved. Consider a standard deck of cards that has 26 red cards and 26 black cards. The probability of an accurrence is based on prior known and 26 black cards. The probability of an accurrence is based on prior known are X = 26cards. The probability of selecting a black card is 26/52 = 0.50 because there are X = 26 black cards and T = 50. black cards and T = 52 total cards. What does this probability mean? If each card is replaced after it is selected at after it is selected, does it mean that I out of the next 2 cards selected will be black? No, because you cannot say for certain what will happen on the next several selections. However, you can say that in the 1 can say that in the long run, if this selection process is continually repeated, the proportion of black cards selected will approach 0.50. Example 4.1 shows another example of computing an

EXAMPLE 4.1

Finding A Priori Probabilities

A standard six-sided die has six faces. Each face of the die contains either one, two, three, four, five, or six dots. If you roll a die, what is the probability that you will get a face with five dots?

SOLUTION Each face is equally likely to occur. Because there are six faces, the probability of getting a face with five dots is 1/6.

The preceding examples use the a priori probability approach because the number of ways the event occurs and the total number of possible outcomes are known from the composition of the deck of cards or the faces of the die.

In the empirical probability approach, the probabilities are based on observed data, not on prior knowledge of a process. Surveys are often used to generate empirical probabilities. Examples of this type of probability are the proportion of individuals in the Using Statistics scenario who actually purchase large-screen HDTVs, the proportion of registered voters who prefer a certain political candidate, and the proportion of students who have part-time jobs. For example, if you take a survey of students, and 60% state that they have part-time jobs, then there is a 0.60 probability that an individual student has a part-time job.

The third approach to probability, subjective probability, differs from the other two approaches because subjective probability differs from person to person. For example, the development team for a new product may assign a probability of 0.60 to the chance of success for the product, while the president of the company may be less optimistic and assign a probability of 0.30. The assignment of subjective probabilities to various outcomes is usually based on a combination of an individual's past experience, personal opinion, and analysis of a particular situation. Subjective probability is especially useful in making decisions in situations in which you cannot use a priori probability or empirical probability.

Events and Sample Spaces

The basic elements of probability theory are the individual outcomes of a variable under study. You need the following definitions to understand probabilities.

Student Tip

aents are represented letters of the alphabet

EVENT

Each possible outcome of a variable is referred to as an event. A simple event is described by a single characteristic.

For example, when you toss a coin, the two possible outcomes are heads and tails. Each of these represents a simple event. When you roll a standard six-sided die in which the six faces of the die contain either one, two, three, four, five, or six dots, there are six possible simple events. An event can be any one of these simple events, a set of them, or a subset of all of them. For example, the event of an even number of dats consists of three simple events (i.e., two, four, or six dots).

Student Tip

The key word when is and.

A joint event is an event that has two or more characteristics.

Getting two heads when you toss a coin twice is an example of a joint event because consists of heads on the first toss and heads on the second toss.

The complement of event A (represented by the symbol A') includes all events that are m_0 part of A.

The complement of a head is a tail because that is the only event that is not a head, The complement of five dots on a die is not getting five dots. Not getting five dots consists of ting one, two, three, four, or six dots.

SAMPLE SPACE

The collection of all the possible events is called the sample space.

The sample space for tossing a coin consists of heads and tails. The sample space when rolling a die consists of one, two, three, four, five, and six dots. Example 4.2 demonstrate events and sample spaces.

EXAMPLE 4.2

Events and Sample Spaces

TABLE 4.1

Purchase Behavior for Large-screen HDTVs

The Using Statistics scenario on page 149 concerns M&R Electronics World. Table 4.1 preents the results of the sample of 1,000 households in terms of purchase behavior for large

PLANNED TO	ACTUALLY P	PURCHASED	
PURCHASE	Yes	No	
Yes	200		Total
No		50	250
Total	100	650	
	300	700	_750
		700	1.000

What is the sample space? Give examples of simple events and joint events.

SOLUTION The sample space consists of the 1,000 respondents. Simple events are "planted purchase." "did not plan to purchase." "out of the 1,000 respondents. to purchase," "did not plan to purchase," "purchased," and "did not purchase." The complement of the event "planned to purchase" is "did not plan to purchase." The event "planned" purchase and actually purchased" is a joint event barrend to purchase." The event "planned" purchase and actually purchased is did not plan to purchase." The event "plans must plan to purchase the television and actually purchase in this joint event, the respondent

Contingency Tables and Venn Diagrams

There are several ways in which you can view a particular sample space. One way involves using a continuous to the voice of the voice o using a contingency table (see Section 2.1) such as the one displayed in Table 4.1. You get the values in the cells of the values in the cells of the table by subdividing the sample space of 1,000 households according to whether some subdividing the sample space of 1,000 households according to whether someone planned to purchase and actually purchased a large-screen HDTV. For example, 200 of the respondents planned to purchase a large-screen HDTV and subsequently did purchase the large-screen HDTV.

A second way to present the sample space is by using a Venn diagram. This diagram graphically represents the various events as "unions" and "intersections" of circles. Figure 4.1 presents a typical Venn diagram for a two-variable situation, with each variable having only two events (A and A', B and B'). The circle on the left (the red one) represents all events that

GURE 4.1 an diagram for events and B





FIGURE 4.2 Venn diagram for the M&R Electronics World example



The circle on the right (the yellow one) represents all events that are part of B. The area contained within circle A and circle B (center area) is the intersection of A and B (written as $A \cap B$), since it is part of A and also part of B. The total area of the two circles is the union of A and B (written as $A \cup B$) and contains all outcomes that are just part of event A, just part of event B, or part of both A and B. The area in the diagram outside of $A \cup B$ contains outcomes that are neither part of A nor part of B.

You must define A and B in order to develop a Venn diagram. You can define either event as A or B, as long as you are consistent in evaluating the various events. For the large-screen HDTV example, you can define the events as follows:

> A =planned to purchase B = actually purchasedA' = did not plan to purchase B' = did not actually purchase

In drawing the Venn diagram (see Figure 4.2), you must first determine the value of the intersection of A and B so that the sample space can be divided into its parts. $A \cap B$ consists of all 200 households who planned to purchase and actually purchased a large-screen HDTV. The remainder of event A (planned to purchase) consists of the 50 households who planned to purchase a large-screen HDTV but did not actually purchase one. The remainder of event B (actually purchased) consists of the 100 households who did not plan to purchase a largescreen HDTV but actually purchased one. The remaining 650 households represent those who neither planned to purchase nor actually purchased a large-screen HDTV.

Simple Probability

Now you can answer some of the questions posed in the Using Statistics scenario. Because the results are based on data collected in a survey (refer to Table 4.1), you can use the empirical probability approach.

As stated previously, the most fundamental rule for probabilities is that they range in value from 0 to 1. An impossible event has a probability of 0, and an event that is certain to occur has a probability of 1.

Simple probability refers to the probability of occurrence of a simple event, P(A). A Simple probability in the Using Statistics scenario is the probability of planning to purchase a large-screen HDTV. How can you determine the probability of selecting a household the planned to purchase a large-screen HDTV? Using Equation (4.1) on page 150:

Probability of occurrence =
$$\frac{X}{T}$$

P(Planned to purchase) = $\frac{\text{Number who planned to purchase}}{\text{Total number of households}}$

= $\frac{250}{1,000} = 0.25$

Thus, there is a 0.25 (or 25%) chance that a household planned to purchase a large-screen HDTV.

Example 4.3 illustrates another application of simple probability.

EXAMPLE 4.3

Computing the Probability That the Large-Screen HDTV Purchased Had a Faster Refresh Rate In the Using Statistics follow-up survey, additional questions were asked of the 300 household that actually purchased large-screen HDTVs. Table 4.2 indicates the consumers' responses to whether the television purchased had a faster refresh rate and whether they also purchased a streaming media box in the past 12 months.

Find the probability that if a household that purchased a large-screen HDTV is randomly selected, the television purchased had a faster refresh rate.

TABLE 4.2

Purchase Behavior Regarding Purchasing a Faster Refresh Rate Television and a Streaming Media Box

REFRESH RATE OF		NG MEDIA DX	
TELEVISION PURCHASED	Yes	No	Total
Faster	38	42	80
Standard	70	150	220
Total	108	192	300

SOLUTION Using the following definitions:

A =purchased a television with a faster refresh rate

A' = purchased a television with a standard refresh rate

B = purchased a streaming media box

B' = did not purchase a streaming media box

 $P(\text{Faster refresh rate}) = \frac{\text{Number of faster refresh rate televisions purchased}}{\text{Total refresh rate}}$

Total number of televisions $\frac{80}{00} = 0.267$

There is a 26.7% chance that a randomly selected large-screen HDTV purchased has a fastor

Joint Probability

Whereas simple probability refers to the probability of occurrence of simple events, joint ample of joint probability is the probability that you will get heads on the second toss of a coin.

In Table 4.1 on page 152, the group of individuals who planned to purchase and actually chased a large screen III. purchased a large-screen HDTV consist only of the outcomes in the single cell "yes—planned to purchase and was to purchase and yes—actually purchased." Because this group consists of 200 households, the probability of picking a household that planned to purchase and actually purchased a large-

$$P(\text{Planned to purchase } and \text{ actually purchased}) = \frac{\text{Planned to purchase } and \text{ actually purchased}}{\text{Total number of respondents}}$$
$$= \frac{200}{1000} = 0.20$$

Example 4.4 also demonstrates how to determine joint probability.

XAMPLE 4.4

etermining the int Probability at a Household rchased a Largereen HDTV with aster Refresh te and Purchased treaming Media

In Table 4.2 on page 154, the purchases are cross-classified as having a faster refresh rate or having a standard refresh rate and whether the household purchased a streaming media box. Find the probability that a randomly selected household that purchased a large-screen HDTV also purchased a television that had a faster refresh rate and purchased a streaming media box.

SOLUTION Using Equation (4.1) on page 150,

P(Television with a faster refresh rate and streaming media box)
$$= \frac{\text{Number that purchased a television with a faster refresh rate and purchased a streaming media box}{\text{Total number of large-screen HDTV purchasers}}$$

$$= \frac{38}{300} = 0.127$$

Therefore, there is a 12.7% chance that a randomly selected household that purchased a largescreen HDTV purchased a television that had a faster refresh rate and purchased a streaming media box.

Marginal Probability

The marginal probability of an event consists of a set of joint probabilities. You can determine the marginal probability of a particular event by using the concept of joint probability just discussed. For example, if B consists of two events, B_1 and B_2 , then P(A), the probability of event A, consists of the joint probability of event A occurring with event B_1 and the joint probability of event A occurring with event B_2 . You use Equation (4.2) to compute marginal probabilities.

$$P(A) = P(A \text{ and } B_1) + P(A \text{ and } B_2) + \cdots + P(A \text{ and } B_k)$$
 (4.2)

where B_1, B_2, \ldots, B_k are k mutually exclusive and collectively exhaustive events, defined as follows:

Two events are mutually exclusive if both the events cannot occur simultaneously. A set of events is collectively exhaustive if one of the events must occur.

Heads and tails in a coin toss are mutually exclusive events. The result of a coin toss collectively. Heads and tails in a coin toss are mutually exclusive simultaneously be a head and a tail. Heads and tails in a coin toss are also collectively experienced occur. It was not occur, tails must occur. If their simultaneously be a head and a tail. Heads and tails the cocur, tails must occur. If tails does not occur, tails must occur. If heads does not occur, tails must occur and the heads does not occur. If heads does not occur, tails must occur and the heads does not occur. If heads does not occur are not occur and the heads does not occur and the h tive events. One of them must occur. If neads uoes not occur, heads must occur. Being male and being female are mutually exclusive) and events occur, heads must occur. Being male and being female are mutually exclusive) and events. occur, neads must occur. Being maie and being terman and everyone is one exhaustive events. No person is both (the two are mutually exclusive), and everyone is one the other (the two are collectively exhaustive).

You can use Equation (4.2) to compute the marginal probability of "planned to purch." a large-screen HDTV:

een HDTV:
$$P(\text{Planned to purchase}) = P(\text{Planned to purchase } and \text{ purchased}) + P(\text{Planned to purchase } and \text{ did not purchase})$$

$$= \frac{200}{1,000} + \frac{50}{1,000}$$

$$= \frac{250}{1,000} = 0.25$$

You get the same result if you add the number of outcomes that make up the simple even "planned to purchase."

Student Tip

The key word when using the addition rule is or.

General Addition Rule

How do you find the probability of event "A or B"? You need to consider the occurrences either event A or event B or both A and B. For example, how can you determine the probabile that a household planned to purchase or actually purchased a large-screen HDTV?

The event "planned to purchase or actually purchased" includes all households that plane. to purchase and all households that actually purchased a large-screen HDTV. You examine and cell of the contingency table (Table 4.1 on page 152) to determine whether it is part of this even From Table 4.1, the cell "planned to purchase and did not actually purchase" is part of the even because it includes respondents who planned to purchase. The cell "did not plan to purchase." actually purchased" is included because it contains respondents who actually purchased. Final the cell "planned to purchase and actually purchased" has both characteristics of interest. The fore, one way to calculate the probability of "planned to purchase or actually purchased" is

P(Planned to purchase or actually purchased) = P(Planned to purchase and did not actuallypurchase) + P(Did not plan topurchase and actually purchased) + P(Planned to purchase and actually purchase $=\frac{50}{1,000}+\frac{100}{1,000}+\frac{200}{1,000}$ $=\frac{350}{1,000}=0.35$

Often, it is easier to determine P(A or B), the probability of the event A or B, by using the general A or B, and A or B is the general A or B.

GENERAL ADDITION RULE

The probability of A or B is equal to the probability of A plus the probability of B matrix the probability of A and B.

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$

EXAMPLE 4.5

Using the General Addition Rule for the Households That Purchased Large-Screen HDTVs In Example 4.3 on page 154, the purchases were cross-classified in Table 4.2 as televisions that had a faster refresh rate or televisions that had a standard refresh rate and whether the household purchased a streaming media box. Find the probability that among households that purchased a large-screen HDTV, they purchased a television that had a faster refresh rate or purchased a streaming media box.

SOLUTION Using Equation (4.3),

P(Television had a faster refresh = P(Television had a faster refresh rate)rate or purchased a streaming media box) + P(purchased a streaming media box)

 P(Television had a faster refresh rate and purchased a streaming media box)

$$= \frac{80}{300} + \frac{108}{300} - \frac{38}{300}$$
$$= \frac{150}{300} = 0.50$$

Therefore, of households that purchased a large-screen HDTV, there is a 50% chance that a randomly selected household purchased a television that had a faster refresh rate or purchased a streaming media box.

Problems for Section 4.1

LEARNING THE BASICS

- 4.1 Two coins are tossed.
- a. Give an example of a simple event.
- h. Give an example of a joint event.
- What is the complement of a head on the first toss?
- d. What does the sample space consist of?

- **4.2** An urn contains 12 red balls and 8 white balls. One ball is to be selected from the urn.
- a. Give an example of a simple event.
- b. What is the complement of a red ball?
- c. What does the sample space consist of?

4.3 Consider the following contingency table:

	В	B'
A	10	20
A'	20	40

What is the probability of event

- b. A'?
- c. A and B?
- d. A or B?
- 4.4 Consider the following contingency table:

	В	B'
A	10	30
A'	25	35

What is the probability of event

- a. A'?
- b. A and B?
- c. A' and B'?
- d. A' or B'?

APPLYING THE CONCEPTS

- 4.5 For each of the following, indicate whether the type of probability involved is an example of a priori probability, empirical probability, or subjective probability.
- a. The next toss of a fair coin will land on heads.
- b. Italy will win soccer's World Cup the next time the competition is held.
- c. The sum of the faces of two dice will be seven.
- d. The train taking a commuter to work will be more than 10 minutes late.
- 4.6 For each of the following, state whether the events created are mutually exclusive and whether they are collectively exhaustive.
- a. Undergraduate business students were asked whether they were sophomores or juniors.
- b. Each respondent was classified by the type of car he or she drives: sedan, SUV, American, European, Asian, or none.
- c. People were asked, "Do you currently live in (i) an apartment or (ii) a house?"
- d. A product was classified as defective or not defective.
- 4.7 Which of the following events occur with a probability of zero? For each, state why or why not.
- a. A company is listed on the New York Stock Exchange and NASDAO.
- b. A consumer owns a smartphone and a tablet.
- c. A cellphone is a Motorola and a Samsung.
- d. An automobile is a Toyota and was manufactured in the United States.
- 4.8 Do males or females feel more tense or stressed out at work? A survey of employed adults conducted online by Harris Interactive

on behalf of the American Psychological Association revealed following:

FELT 1	TENS	E O	R
STRESSED	OUT	AT	WORK

GENDER	Yes	No
Male	244	495
Female	282	480

Source: Data extracted from "The 2013 Work and Well-Being Survey," American Psychological Association and Harris Interactive, March 2013. p. 5, bit.ly/11JGcPf.

- a. Give an example of a simple event.
- b. Give an example of a joint event.
- c. What is the complement of "Felt tense or stressed out at wor
- d. Why is "Male and felt tense or stressed out at work" a joint ex
- 4.9 Referring to the contingency table in Problem 4.8, if any ployed adult is selected at random, what is the probability that
- a. the employed adult felt tense or stressed out at work?
- b. the employed adult was a male who felt tense or stressed on
- c. the employed adult was a male or felt tense or stressed ou work?
- d. Explain the difference in the results in (b) and (c).
- 4.10 How will marketers change their social media use in them future? A survey by Social Media Examiner reported that 789 B2B marketers (marketers that focus primarily on attracting b nesses) plan to increase their use of LinkedIn, as compared to of B2C marketers (marketers that primarily target consume The survey was based on 1,331 B2B marketers and 1,694 B. marketers. The following table summarizes the results:

INCREASE USE	BU	ISINESS FOO	US
OF LINKEDIN?	B2B	B2C	Total
Yes	1,038	915	1.953
No Total	_ 293	779	1,072
Course	1,331	1,694	3,025

ce; Data extracted from "2013 Social Media Marketing Industry Report," May 2013, bit.ly/Ig5vMQN.

- a. Give an example of a simple event.
- b. Give an example of a joint event.
- c. What is the complement of a marketer who plans to incle use of LinkedIn?
- d. Why is a marketer who plans to increase use of Linkedin and a B2C marketer a joint event?
- 4.11 Referring to the contingency table in Problem 4.10.1 marketer is selected at random, what is the probability that
- a. he or she plans to increase use of LinkedIn?
- b. he or she is a B2C marketer?
- c. he or she plans to increase use of LinkedIn or is a B. marketer?
- d. Explain the difference in the results in (b) and (c).

SELF 4.12 What business and technical skills are critical for today's business intelligence/analytics and informaion management professionals? As part of InformationWeek's 2013 U.S. IT Salary Survey, business intelligence/analytics and 2013 C.S. and a second staff and managers, both staff and managers, were asked to indicate what business and technical skills are critito their job. The list of business and technical skills included analyzing Data. The following table summarizes the responses to his skill:

PROFESS		
Staff		Total
4,374		
3,436		8,007
7,810	6.264	
	Staff 4,374 3,436	4,374 3,633 3,436 2,631

InformationWeek Reports, April 2013, p. 40. ubm.io/lewjKT5.

f a professional is selected at random, what is the probability that ne or she

- a. indicates analyzing data as critical to his or her job?
- b. is a manager?
- indicates analyzing data as critical to his or her job or is a manager?
- i. Explain the difference in the results in (b) and (c).
- 4.13 Do Americans prefer Coke or Pepsi? A survey was conducted by Public Policy Polling (PPP) in 2013; the results were as follows:

	GENE	ER	
PREFERENCE	Female	Male	Total
Coke	120	95	215
Pepsi	95	80	175
Neither/Unsure	65	45	110
Total	280	220	500

Source: Data extracted from "Public Policy Polling," Report 2013. bit.ly/YKXfzN.

If an American is selected at random, what is the probability that he or she

- a. prefers Pepsi?
- b. is male and prefers Pepsi?
- c. is male or prefers Pepsi?
- d. Explain the difference in the results in (b) and (c).
- 4.14 A survey of 1,085 adults asked, "Do you enjoy shopping for clothing for yourself?" The results (data extracted from "Split Decision on Clothes Shopping," USA Today, January 28, 2011, p. 1B) indicated that 51% of the females enjoyed shopping for clothing for themselves as compared to 44% of the males. The sample sizes of males and females were not provided. Suppose that the results indicated that of 542 males, 238 answered yes. Of 543 females, 276 answered yes. Construct a contingency table to evaluate the probabilities. What is the probability that a respondent chosen at
- a. enjoys shopping for clothing for himself or herself?
- b. is a female and enjoys shopping for clothing for herself?
- c. is a female or is a person who enjoys shopping for clothing?
- d. is a male or a female?
- 4.15 Each year, ratings are compiled concerning the performance of new cars during the first 90 days of use. Suppose that the cars have been categorized according to whether a car needs warrantyrelated repair (yes or no) and the country in which the company manufacturing a car is based (United States or not United States). Based on the data collected, the probability that the new car needs a warranty repair is 0.04, the probability that the car was manufactured by a U.S.-based company is 0.60, and the probability that the new car needs a warranty repair and was manufactured by a U.S.based company is 0.025. Construct a contingency table to evaluate the probabilities of a warranty-related repair. What is the probability that a new car selected at random
- a. needs a warranty repair?
- b. needs a warranty repair and was manufactured by a U.S.-based company?
- c. needs a warranty repair or was manufactured by a U.S.-based
- d. needs a warranty repair or was not manufactured by a U.S.based company?

4.2 Conditional Probability

Each example in Section 4.1 involves finding the probability of an event when sampling from the entire sample space. How do you determine the probability of an event if you know certain information about the events involved?

Computing Conditional Probabilities

Conditional probability refers to the probability of event A, given information about the occurrence of another event, B.

The probability of A given B is equal to the probability of A and B divided by the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A and B divided by the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A given B is equal to the probability of A giv ity of B.

$$P(A|B) = \frac{P(A \text{ and } B)}{P(B)}$$
(4.4a)

The probability of B given A is equal to the probability of A and B divided by the probability of A.

$$P(B|A) = \frac{P(A \text{ and } B)}{P(A)}$$
(4.4b)

where

$$P(A \text{ and } B) = \text{ joint probability of } A \text{ and } B$$

 $P(A) = \text{marginal probability of } A$
 $P(B) = \text{marginal probability of } B$

-Student Tip

The variable that is given goes in the denominator of Equation (4.4). Since you were given planned to purchase, planned to purchase is in the denominator.

Referring to the Using Statistics scenario involving the purchase of large-screen HDTVs suppose you were told that a household planned to purchase a large-screen HDTV. Now, what is the probability that the household actually purchased the television?

In this example, the objective is to find P(Actually purchased | Planned to purchase). Her you are given the information that the household planned to purchase the large-screen HDTV Therefore, the sample space does not consist of all 1,000 households in the survey. It consists of only those households that planned to purchase the large-screen HDTV. Of 250 such households 200 actually purchased the large-screen HDTV. Therefore, based on Table 4.1 on page 152.1 probability that a household actually purchased the large-screen HDTV given that they planned to purchase is

$$P(\text{Actually purchased} | \text{Planned to purchase}) = \frac{\text{Planned to purchase } and \text{ actually purchased}}{\text{Planned to purchase}}$$

$$= \frac{200}{250} = 0.80$$

You can also use Equation (4.4b) to compute this result:

$$P(B|A) = \frac{P(A \text{ and } B)}{P(A)}$$

where

A =planned to purchase

B = actually purchased

then

$$P(\text{Actually purchased} | \text{Planned to purchase}) = \frac{200/1,000}{250/1,000}$$
$$= \frac{200}{250} = 0.80$$

Example 4.6 further illustrates conditional probability.

EXAMPLE 4.6 finding the Conditional Probability of Purchasing a Streaming Media Box

Table 4.2 on page 154 is a contingency table for whether a household purchased a television with a faster refrank and a contingency table for whether a household purchased a television with a faster refrank and a contingency table for whether a household purchased a television with a faster refrank and a contingency table for whether a household purchased a television with a faster refrank and a contingency table for whether a household purchased a television with a faster refrank and a contingency table for whether a household purchased a television whether the contingency table for the contingency table for the contingency table and the contingency table for the contingency table and table at the contingency table and table at the contingency table at the co with a faster refresh rate and whether the household purchased a streaming media box. If a household purchased a streaming media box. If a household purchased a streaming media purchased media purchase

SOLUTION Because you know that the household purchased a television with a faster refresh rate, the sample space is reduced to 80 households. Of these 80 households, 38 also purchased a streaming media box. Therefore, the probability that a household purchased a streaming media box, given that the household purchased a television with a faster refresh rate, is

P(Purchased streaming media box | Purchased = faster refresh rate and streaming media box | television with 6 media box | Purchased = faster refresh rate and streaming media box television with faster refresh rate)

Number purchasing television with Number purchasing television

with faster refresh rate
$$= \frac{38}{80} = 0.475$$

If you use Equation (4.4b) on page 160:

A =purchased a television with a faster refresh rate

B =purchased a streaming media box

then


$$P(B|A) = \frac{P(A \text{ and } B)}{P(A)} = \frac{38/300}{80/300} = 0.475$$

Therefore, given that the household purchased a television with a faster refresh rate, there is a 47.5% chance that the household also purchased a streaming media box. You can compare this conditional probability to the marginal probability of purchasing a streaming media box, which is 108/300 = 0.36, or 36%. These results tell you that households that purchased televisions with a faster refresh rate are more likely to purchase a streaming media box than are households that purchased large-screen HDTVs that have a standard refresh rate,

Decision Trees

In Table 4.1 on page 152, households are classified according to whether they planned to purchase and whether they actually purchased large-screen HDTVs. A decision tree is an alternative to the contingency table. Figure 4.3 represents the decision tree for this example.

GURE 4.3 rision tree for planned Purchase and actually achased



In Figure 4.3, beginning at the left with the entire set of households, there are hard planned to purchase a large-screen HDT. In Figure 4.3, beginning at the left with the strong to whether the household activity the branches for whether or not the household planned to whether the household activity the branches. "branches" for whether or not the household planned to whether the household actually plan of these branches has two subbranches, corresponding to whether the household actually plan of these branches has two subbranches, corresponding to whether the household actually plan of these branches has two subbranches, corresponding to whether the household actually plan of these branches has two subbranches, corresponding to whether the household actually plan of these branches has two subbranches, corresponding to whether the household actually plan of these branches has two subbranches, corresponding to whether the household actually plan of these branches has two subbranches, corresponding to whether the household actually plan of these branches has two subbranches, corresponding to whether the household actually plan of these branches has two subbranches, corresponding to whether the household actually plan to the probabilities at the probabi of these branches has two subbranches, corresponding to the probabilities at the end of chased or did not actually purchase the large-screen HDTV. The probabilities of A and A'. The probabilities of A and A'. chased or did not actually purchase the large-screen 122 of A and A'. The probabilities at the initial branches represent the marginal probabilities of A and A'. The probabilities at the the initial branches represent the marginar probability for each combination of each of the four subbranches represent the joint probability for each combination of end of each of the four subtranches represent the joint probability by dividing the joint probability b_1 events A and B. You compute the conditional probability by appropriate marginal probability.

For example, to compute the probability that the household actually purchased, given the the appropriate marginal probability.

For example, to compute the probability that the household planned to purchase the large-screen HDTV, you take P(Planned to purchase the large-screen HDTV, you take <math>P(Planned to purchase the large-screen HDTV)and actually purchased) and divide by P(Planned to purchase). From Figure 4.3,

$$P(\text{Actually purchased} | \text{Planned to purchase}) = \frac{200/1,000}{250/1,000}$$
$$= \frac{200}{250} = 0.80$$

Example 4.7 illustrates how to construct a decision tree.

EXAMPLE 4.7


Constructing the Decision Tree for the Households That Purchased Large-Screen HDTVs

Using the cross-classified data in Table 4.2 on page 154, construct the decision tree. Use the decision tree to find the probability that a household purchased a streaming media box, given that the household purchased a television with a faster refresh rate.

SOLUTION The decision tree for purchased a streaming media box and a television with faster refresh rate is displayed in Figure 4.4.

FIGURE 4.4

Decision tree for purchased a television with a faster refresh rate and a streaming media



Using Equation (4.4b) on page 160 and the following definitions, A = purchased a television with a faster refresh rate

B = purchased a streaming media box

$$P(B|A) = \frac{P(A \text{ and } B)}{P(A)} = \frac{38/300}{80/300} = 0.475$$

Independence

In the example concerning the purchase of large-screen HDTVs, the conditional probability is 200/250 = 0.80 that the selected household actually purchased the large-screen HDTV, given that the household planned to purchase. The simple probability of selecting a household that actually planned to purchase affected the probability that the prior knowledge that the household In other words, the outcome of one matrix of the probability of the probability purchased the television.

In other words, the outcome of one event is *dependent* on the outcome of a second event.

When the outcome of one event does *not* affect the probability of occurrence of another event, the events are said to be independent. **Independence** can be determined by using Equation (4.5).

INDEPENDENCE

Two events, A and B, are independent if and only if

$$P(A|B) = P(A) \tag{4.5}$$

where

P(A|B) = conditional probability of A given BP(A) = marginal probability of A

Example 4.8 demonstrates the use of Equation (4.5).

XAMPLE 4.8

Determining Independence

ABLE 4.3

Purchase of Arge-Screen In the follow-up survey of the 300 households that actually purchased large-screen HDTVs, the households were asked if they were satisfied with their purchases. Table 4.3 cross-classifies the responses to the satisfaction question with the responses to whether the television had a faster refresh rate.

TELEVISION	SATISFIE PURCH		
REFRESH RATE	Yes	No	Total
Faster	64	16	80
Standard	176	44	220
Total	240	60	300

Determine whether being satisfied with the purchase and the refresh rate of the television purchased are independent.

SOLUTION For these data.

$$P(\text{Satisfied} | \text{Faster refresh rate}) = \frac{64/300}{80/300} = \frac{64}{80} = 0.80$$

which is equal to

$$P(\text{Satisfied}) = \frac{240}{300} = 0.80$$

Thus, being satisfied with the purchase and the refresh rate of the television purchased are independent. Knowledge of one event does not affect the probability of the other event.

Example 4.9 demonstrates the use of the general multiplication rule.

EXAMPLE 4.9

Using the General Multiplication Rule Consider the 80 households that purchased televisions that had a faster refresh rate. In Table 4.3 on page 163, you see that 64 households are satisfied with their purchase, and 16 households are dissatisfied. Suppose 2 households are randomly selected from the 80 households. Find the probability that both households are satisfied with their purchase.

SOLUTION Here you can use the multiplication rule in the following way. If

A = second household selected is satisfied

B =first household selected is satisfied

then, using Equation (4.6),

$$P(A \text{ and } B) = P(A | B)P(B)$$

The probability that the first household is satisfied with the purchase is 64/80. However, the probability that the second household is also satisfied with the purchase depends on the result of the first selection. If the first household is not returned to the sample after the satisfaction level is determined (i.e., sampling without replacement), the number of households remaining is 79. If the first household is satisfied, the probability that the second is also satisfied is 63/79 because 63 satisfied households remain in the sample. Therefore,

$$P(A \text{ and } B) = \left(\frac{63}{79}\right) \left(\frac{64}{80}\right) = 0.6380$$

There is a 63.80% chance that both of the households sampled will be satisfied with their purchase

The multiplication rule for independent events is derived by substituting $P(A \mid B)$ in Equation (4.6).

If this rule holds for two events, A and B, then A and B are independent. Therefore, there are two ways to determine independence:

1. Events A and B are independent if, and only if, P(A|B) = P(A).

2. Events A and B are independent if, and only if, P(A and B) = P(A)P(B).

Marginal Probability Using the General Multiplication Rule

In Section 4.1, marginal probability was defined using Equation (4.2) on page 155. You can state the equation for marginal probability by using the general multiplication rule. If

$$P(A) = P(A \text{ and } B_1) + P(A \text{ and } B_2) + \cdots + P(A \text{ and } B_k)$$

then, using the general multiplication rule, Equation (4.8) defines the marginal probability.

MARGINAL PROBABILITY USING THE GENERAL MULTIPLICATION RULE

$$P(A) = P(A|B_1)P(B_1) + P(A|B_2)P(B_2) + \cdots + P(A|B_k)P(B_k)$$
 (4.8)

where B_1, B_2, \ldots, B_k are k mutually exclusive and collectively exhaustive events.

To illustrate Equation (4.8), refer to Table 4.1 on page 152. Let

P(A) = probability of planned to purchase

 $P(B_1)$ = probability of actually purchased

 $P(B_2)$ = probability of did not actually purchase

Then, using Equation (4.8), the probability of planned to purchase is

$$P(A) = P(A|B_1)P(B_1) + P(A|B_2)P(B_2)$$

$$= \left(\frac{200}{300}\right)\left(\frac{300}{1,000}\right) + \left(\frac{50}{700}\right)\left(\frac{700}{1,000}\right)$$

$$= \frac{200}{1,000} + \frac{50}{1,000} = \frac{250}{1,000} = 0.25$$

Problems for Section 4.2

LEARNING THE BASICS

4.16 Consider the following contingency table:

	В	B'
A	10	20
A'	20	40

What is the probability of

a. A | B?

b. A B'?

C. A' B'?

d. Are events A and B independent?

4.17 Consider the following contingency table:

	В	B
A	10	30
A'	25	35

What is the probability of

a. A B?

b. A' | B'?

c. A B'?

d. Are events A and B independent?

4.18 If P(A and B) = 0.4 and P(B) = 0.8, find P(A|B).

4.19 If P(A) = 0.7, P(B) = 0.6, and A and B are independent, find P(A and B).

4.20 If P(A) = 0.3, P(B) = 0.4, and P(A and B) = 0.2, are Aand B independent?

APPLYING THE CONCEPTS

4.21 Do males or females feel more tense or stressed out at work? A survey of employed adults conducted online by Harris Interactive on behalf of the American Psychological Association revealed the following:

FELT TENSE OR	
STRESSED OUT AT WORK	

GENDER	Yes	No
Male	244	495
Female	282	480

Source: Data extracted from "The 2013 Work and Well-Being Survey," American Psychological Association and Harris Interactive, March 2013, p. 5, bit.ly/11JGcPf.

- a. Given that the employed adult felt tense or stressed out at work, what is the probability that the employed adult was a male?
- b. Given that the employed adult is male, what is the probability that he felt tense or stressed out at work?
- c. Explain the difference in the results in (a) and (b).
- d. Is feeling tense or stressed out at work and gender independent?
- 4.22 How will marketers change their social media use in the near future? A survey by Social Media Examiner of B2B marketers (marketers that focus primarily on attracting businesses) and B2C marketers (marketers that primarily target consumers) was based on 1,331 B2B marketers and 1,694 B2C marketers. The following table summarizes the results:

INCREASE USE	BUSINESS FOCUS				
OF LINKEDIN?	В2В	B2C	Total		
Yes	1,038	915	1,953		
No	293	779	1,072		
Total	1.331	1,694	3,025		

Source: Data extracted from "2013 Social Media Marketing Industry Report," May 2013, bit.ly/1g5vMQN.

- a. Suppose you know that the marketer is a B2B marketer. What is the probability that he or she plans to increase use of
- b. Suppose you know that the marketer is a B2C marketer. What is the probability that he or she plans to increase use of LinkedIn?
- c. Are the two events, increase use of LinkedIn and business focus, independent? Explain.

4.23 Do Americans prefer Coke or Pepsi? A survey was com 4.23 Do Americans producted by Public Policy Polling (PPP) in 2013; the results were as follows:

	GENE		
	Female	Male	Total
PREFERENCE	120	95	215
Coke	95	80	175
Pepsi	65	45	110
Neither/Unsure	280	220	500
Total Source: Data extracted fr	om "Public Poli	cy Polling	Report

2013, bit.ly/YKX

- a. Given that an American is a male, what is the probability that he prefers Pepsi?
- b. Given that an American is a female, what is the probability that she prefers Pepsi?
- c. Is preference independent of gender? Explain.

/ SELF 4.24 What business and technical skills are critical for today's business intelligence/analytics and information management professionals? As part of InformationWeek's 2013 U.S. IT Salary Survey, business intelligence/analytics and information management professionals, both staff and managers. were asked to indicate what business and technical skills are cntical to their job. The list of business and technical skills included Analyzing Data. The following table summarizes the responses to this skill:

	PROFES			
ANALYZING DATA	Staff	Management	Total	
Critical	4,374		8,007	
Not critical	3,436	3,633	6,067	
Total	-	2,631		
	7,810	6,264	14,07	
Information Week P	rom "IT Salarie	Show Slow Growth," 40, ubm.io/lewjKT5		

- a. Given that a professional is staff, what is the probability that the professional indicates analyzing data as critical to his or he
- b. Given that a professional is staff, what is the probability that the professional does not indicate analyzing data as critical to
- c. Given that a professional is a manager, what is the probability that the professional indicates analyzing data as critical to her indicates
- d. Given that a professional is a manager, what is the probability that the professional does not indicate analyzing data as critical

4.25 A survey of 1,085 adults asked, "Do you enjoy shopping for clothing for yourself?" The results (data extracted from "Split Decision on Clothes Shopping," USA Today, January 28, 2011, p. 1B) indicated that 51% of the females enjoyed shopping for clothing for themselves as compared to 44% of the males. The sample sizes of males and females were not provided. Suppose that the results were as shown in the following table:

ENJOYS SHOPPING	GE		
FOR CLOTHING	Male	Female	Total
Yes	238	276	514
No	304	267	571
Total	542	543	1,085

- a. Suppose that the respondent chosen is a female. What is the probability that she does not enjoy shopping for clothing?
- b. Suppose that the respondent chosen enjoys shopping for clothing. What is the probability that the individual is a male?
- c. Are enjoying shopping for clothing and the gender of the individual independent? Explain.
- 4.26 Each year, ratings are compiled concerning the performance of new cars during the first 90 days of use. Suppose that the cars have been categorized according to whether a car needs warranty-related repair (yes or no) and the country in which the company manufacturing a car is based (United States or not United States). Based on the data collected, the probability that the new car needs a warranty repair is 0.04, the probability that the car is manufactured by a U.S.-based company is 0.60, and the probability that the new car needs a warranty repair and was manufactured by a U.S.-based company is 0.025.
- a. Suppose you know that a company based in the United States manufactured a particular car. What is the probability that the car needs a warranty repair?
- b. Suppose you know that a company based in the United States did not manufacture a particular car. What is the probability that the car needs a warranty repair?
- c. Are need for a warranty repair and location of the company manufacturing the car independent?
- 4.27 In 41 of the 63 years from 1950 through 2013 (in 2011 there was virtually no change), the S&P 500 finished higher after the first five days of trading. In 36 out of 41 years, the S&P 500 finished higher for the year. Is a good first week a good omen for the upcoming year? The following table gives the first-week and annual performance over this 63-year period:

	S&P 500'S ANNUAL PERFORMAN		
FIRST WEEK	Higher	Lower	
Higher	36	5	
Lower	11	1.1	

- a. If a year is selected at random, what is the probability that the S&P 500 finished higher for the year?
- b. Given that the S&P 500 finished higher after the first five days of trading, what is the probability that it finished higher for the year?
- c. Are the two events "first-week performance" and "annual performance" independent? Explain.
- d. Look up the performance after the first five days of 2014 and the 2014 annual performance of the S&P 500 at finance.yahoo.com. Comment on the results.
- **4.28** A standard deck of cards is being used to play a game. There are four suits (hearts, diamonds, clubs, and spades), each having 13 faces (ace, 2, 3, 4, 5, 6, 7, 8, 9, 10, jack, queen, and king), making a total of 52 cards. This complete deck is thoroughly mixed, and you will receive the first 2 cards from the deck, without replacement (the first card is not returned to the deck after it is selected).
- a. What is the probability that both cards are queens?
- b. What is the probability that the first card is a 10 and the second card is a 5 or 6?
- c. If you were sampling with replacement (the first card is returned to the deck after it is selected), what would be the answer in (a)?
- d. In the game of blackjack, the face cards (jack, queen, king) count as 10 points, and the ace counts as either 1 or 11 points. All other cards are counted at their face value. Blackjack is achieved if 2 cards total 21 points. What is the probability of getting blackjack in this problem?
- **4.29** A box of nine iPhone 5C cellphones (the iPhone "for the colorful") contains two yellow cellphones and seven green cellphones.
- a. If two cellphones are randomly selected from the box, without replacement (the first cellphone is not returned to the box after it is selected), what is the probability that both cellphones selected will be green?
- b. If two cellphones are randomly selected from the box, without replacement (the first cellphone is not returned to the box after it is selected), what is the probability that there will be one yellow cellphone and one green cellphone selected?
- c. If three cellphones are selected, with replacement (the cellphones are returned to the box after they are selected), what is the probability that all three will be yellow?
- d. If you were sampling with replacement (the first cellphone is returned to the box after it is selected), what would be the answers to (a) and (b)?

4.3 Bayes' Theorem

Bayes' theorem is used to revise previously calculated probabilities based on new information. Developed by Thomas Bayes in the eighteenth century (see references 1, 2, 3, and 8), Bayes' theorem is an extension of what you previously learned about conditional probability.

You can apply Bayes' theorem to the situation in which M&R Electronics World is considering marketing a new model of televisions. In the past, 40% of the new-model televisions have been successful, and 60% have been unsuccessful. Before introducing the new-model

television, the marketing research department conducts an extensive study and releases television, the marketing research department contracts of the successful new-model a report, either favorable or unfavorable. In the past, 80% of the successful new-model a report, either favorable or unfavorable, the past, 80% of the successful new-model as the successful new-model n a report, either favorable or unfavorable. In the past, of the unsuccessful television(s) had received favorable market research reports, and 30% of the unsuccessful television(s) had received favorable market research reports. For the new model of television new-model television(s) had received favorable reports. For the new model of television unnew-model television(s) had received ravorable reports on the der consideration, the marketing research department has issued a favorable report. What is the probability that the television will be successful?

Bayes' theorem is developed from the definition of conditional probability. To find the bayes' theorem is developed from the definition of A, consider Equation (4.4b) (originally presented as conditional probability of B given A, consider Equation (4.4b) page 160 and shown again below):

$$P(B|A) = \frac{P(A \text{ and } B)}{P(A)} = \frac{P(A|B)P(B)}{P(A)}$$

Bayes' theorem is derived by substituting Equation (4.8) on page 165 for P(A) in the denominator of Equation (4.4b).

BAYES' THEOREM

$$P(B_i|A) = \frac{P(A|B_i)P(B_i)}{P(A|B_1)P(B_1) + P(A|B_2)P(B_2) + \cdots + P(A|B_k)P(B_k)}$$
(4.9)

where B_i is the *i*th event out of k mutually exclusive and collectively exhaustive events

To use Equation (4.9) for the television-marketing example, let

event S = successful television event F = favorable reportevent S' = unsuccessful television event F' = unfavorable report

and

$$P(S) = 0.40$$
 $P(F|S) = 0.80$
 $P(S') = 0.60$ $P(F|S') = 0.30$

Then, using Equation (4.9),

$$P(S|F) = \frac{P(F|S)P(S)}{P(F|S)P(S) + P(F|S')P(S')}$$

$$= \frac{(0.80)(0.40)}{(0.80)(0.40) + (0.30)(0.60)}$$

$$= \frac{0.32}{0.32 + 0.18} = \frac{0.32}{0.50}$$

$$= 0.64$$

The probability of a successful television, given that a favorable report was received 0.64. Thus, the probability of an unsuccessful television, given that a favorable report $\frac{1}{4}$ ceived, is 1 - 0.64 = 0.36.

Table 4.4 summarizes the computation of the probabilities, and Figure 4.5 present decision tree.

EXAMPLE 4.10

Using Bayes' Theorem in a Medical Diagnosis Problem The probability that a person has a certain disease is 0.03. Medical diagnostic tests are available to determine whether the person actually has the disease. If the disease is actually present, the probability that the medical diagnostic test will give a positive result (indicating that the disease is present) is 0.90. If the disease is not actually present, the probability of a positive test result (indicating that the disease is present) is 0.02. Suppose that the medical diagnostic test has given a positive result (indicating that the disease is present). What is the probability that the disease is actually present? What is the probability of a positive test result?

SOLUTION Let

event
$$D' = \text{has disease}$$
 event $T' = \text{test is positive}$
event $D' = \text{does not have disease}$ event $T' = \text{test is negative}$

and

$$P(D) = 0.03$$
 $P(T|D) = 0.90$
 $P(D') = 0.97$ $P(T|D') = 0.02$

Using Equation (4.9) on page 168,

$$P(D|T) = \frac{P(T|D)P(D)}{P(T|D)P(D) + P(T|D')P(D')}$$

$$= \frac{(0.90)(0.03)}{(0.90)(0.03) + (0.02)(0.97)}$$

$$= \frac{0.0270}{0.0270 + 0.0194} = \frac{0.0270}{0.0464}$$

$$= 0.582$$

(continued)

The probability that the disease is actually present, given that a positive result has occurred (indicating that the disease is present), is 0.582. Table 4.5 summarizes the computation of the (indicating that the disease is present), is 0.582. Table 4.5 summarizes the computation of the probabilities, and Figure 4.6 presents the decision tree. The denominator in Bayes' theorem probabilities, and Figure 4.6 presents the decision tree. The denominator in Bayes' theorem probabilities, and Figure 4.6 presents the decision tree. The denominator in Bayes' theorem probabilities, and Figure 4.6 presents the decision tree.


TABLE 4.5

Bayes' Theorem Computations for the Medical Diagnosis Problem

	Probability	Conditional Probability $P(T D_i)$	Joint Probability $P(T D_i)P(D_i)$	Probability $P(D_i T)$
Event D_l $D = \text{has disease}$	$\frac{P(D_i)}{0.03}$	0.90	0.0270	P(D T) = 0.0270/0.0464 = 0.582
D' = does not have disease	0.97	0.02	0.0194 0.0464	P(D' T) = 0.0194/0.0464 = 0.418

FIGURE 4.6

Decision tree for a medical diagnosis problem



THINK ABOUT THIS Divine Providence and Spam

Would you ever guess that the essays Divine Benevolence: Or, An Attempt to Prove That the Principal End of the Divine Providence and Government is the Happiness of His Creatures and An Essay Towards Solving a Problem in the Doctrine of Chances were written by the same person? Probably not, and in doing so, you illustrate a modernday application of Bayesian statistics: spam, or junk mail filters.

In not guessing correctly, you probably looked at the words in the titles of the essays and concluded that they were talking about two different things. An implicit rule you used was that word frequencies vary by subject matter. A statistics essay would very likely contain the word statistics as well as words such as chance, problem, and solving. An eighteenth-century essay about theology and religion would be more likely to contain the uppercase forms of Divine and Providence.

Likewise, there are words you would guess to be very unlikely to appear in either book, such as technical terms from finance, and words that are most likely to appear in both—common words such as *a*, *and*, and *the*. That words would be either likely or unlikely suggests an application of probability theory. Of course, likely and unlikely are fuzzy concepts, and we might occasionally misclassify an essay if we kept things too simple, such as relying solely on the occurrence of the words *Divine* and *Providence*.

For example, a profile of the late Harris Milstead, better known as *Divine*, the star of *Hairspray* and other films, visiting Providence (Rhode Island), would most certainly not be an essay about theology. But if we widened the number of words we examined and found such words as *movie* or the name John Waters (Divine's director in many films), we probably would quickly realize the essay had something to do with twentieth-century cinema and little to do with theology and religion.

We can use a similar process to try to classify a new email message in your in-box as either spam or a legitimate message (called "ham," in this context). We would first need to add to your email program a "spam filter" that has the ability to track word frequencies associated with spam and

ham messages as you identify them on a day-like day basis. This would allow the filter to constant update the prior probabilities necessary to us Bayes' theorem. With these probabilities, the filter can ask, "What is the probability that an email spam, given the presence of a certain word?"

Applying the terms of Equation (4.9) apage 168, such a Bayesian spam filter would mitiply the probability of finding the word in a sparemail, P(A|B), by the probability that the emalspam, P(B), and then divide by the probability finding the word in an email, the denominator Equation (4.9). Bayesian spam filters also use structure by focusing on a small set of words that has high probability of being found in a spam messar as well as on a small set of other words that has low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being found in a spam messar low probability of being

As spammers (people who send junk embedded of such new filters, they tried to dethem. Having learned that Bayesian filters no be assigning a high $P(A \mid B)$ value to words amonly found in spam, such as Viagra. Spammers thought they specified

USING STATISTICS

Possibilities at M&R Electronics World, Revisited

s the marketing manager for M&R Electronics World, you analyzed the survey results of an intent-to-purchase study. This study asked the heads of 1,000 households about their intentions to purchase a large-screen HDTV sometime during the next 12 months, and as a follow-up, M&R surveyed the same people 12 months later to see whether such a television was purchased. In addition, for households purchasing large-screen HDTVs, the survey asked whether the television they purchased had a faster refresh rate, whether they also purchased a streaming media box in the past 12 months, and whether they were satisfied with their purchase of the large-screen HDTV.

By analyzing the results of these surveys, you were able to uncover many pieces of valuable information that will help you plan a marketing strategy to enhance sales and better target those households likely to purchase multiple or more expensive products. Whereas only 30% of the households actually purchased a large-screen HDTV, if a household indicated that it planned to purchase a large-screen HDTV in the next 12 months, there was an 80% chance that the household actually made the purchase. Thus the marketing strategy



should target those households that have indicated an intertion to purchase.

You determined that for households that purchased television that had a faster refresh rate, there was a 47.50 chance that the household also purchased a streaming med box. You then compared this conditional probability to the marginal probability of purchasing a streaming media box which was 36%. Thus, households that purchased television that had a faster refresh rate are more likely to purchase streaming media box than are households that purchase large-screen HDTVs that have a standard refresh rate.

You were also able to apply Bayes' theorem to M&R Electronics World's market research reports. The reports investigate a potential new television model prior to its scheduled release. If a favorable report was received, then there was a 64% chance that the new television model would be successful However, if an unfavorable report was received, there is only a 16% chance that the model would be successful. Therefore the marketing strategy of M&R needs to pay close attention whether a report's conclusion is favorable or unfavorable.

SUMMARY

This chapter began by developing the basic concepts of probability. You learned that probability is a numeric value from 0 to 1 that represents the chance, likelihood, or possibility that a particular event will occur. In addition to simple probability, you learned about conditional probabilities and independent events. Bayes' theorem was used to revise previously calculated probabilities based on new inform tion. Throughout the chapter, contingency tables and dec sion trees were used to display information. You also learned about several counting rules. In the next chapter, imponi discrete probability distributions including the binomial Poisson distributions are developed.

REFERENCES

- 1. Anderson-Cook, C. M. "Unraveling Bayes' Theorem." Quality Progress, March 2014, p. 52-54.
- 2. Bellhouse, D. R. "The Reverend Thomas Bayes, FRS: A Biography to Celebrate the Tercentenary of His Birth," Statistical Science, 19 (2004), 3-43.
- 3. Hooper, W. "Probing Probabilities." Quality Progress, March 2014, pp. 18-22.
- 4. Lowd, D., and C. Meek. "Good Word Attacks on Statistical Spam Filters." Presented at the Second Conference on Email
- 5. Microsoft Excel 2013. Redmond, WA: Microsoft Corp. 1010
- 6. Minitab Release 16. State College, PA: Minitab, Inc. 2010.
- 7. Paulos, J. A. *Innumeracy*. New York: Hill and Wang, 1988. Silberman, S. 1971. 8. Silberman, S. "The Quest for Meaning," Wired 8.02, Februa 2000.
- 9. Zeller, T. "The Fight Against V1@gra (and Other Spam) New York Times, May 21, 2006, pp. B1, B6.

Probability of Occurrence		Marginal Probability Using the General	
probability of occurrence $=\frac{X}{T}$	(4.1)	Multiplication Rule $P(A) = P(A B_1)P(B_1) + P(A B_2)P(B_2) + \cdots + P(A B_k)P(B_k)$	(4.8)
Marginal Probability			
$P(A) = P(A \text{ and } B_1) + P(A \text{ and } B_2) + \cdots + P(A \text{ and } B_k)$	(4.2)	Bayes' Theorem $P(B_i A) = P(A B_i)P(B_i)$	
General Addition Rule		$\frac{P(A B_i)P(B_i)}{P(A B_1)P(B_1) + P(A B_2)P(B_2) + \cdots + P(A B_n)P(B_n)}$	
p(A or B) = P(A) + P(B) - P(A and B)	(4.3)		(4.9)
Conditional Probability P(A and B)		Counting Rule 1	(4.10)
$P(A B) = \frac{P(A \text{ and } B)}{P(B)}$	(4.4a)	Counting Rule 2	
$P(B A) = \frac{P(A \text{ and } B)}{P(A)}$	(4.4b)	$(k_1)(k_2)\ldots(k_n)$	(4.11)
		Counting Rule 3	
Independence		$n! = (n)(n-1)\dots(1)$	(4.12)
P(A B) = P(A)	(4.5)	Counting Rule 4: Permutations	
General Multiplication Rule $P(A \text{ and } B) = P(A B)P(B)$	(4.6)	${}_{n}P_{x}=\frac{n!}{(n-x)!}$	(4.13)
Multiplication Rule for Independent Events		Counting Rule 5: Combinations	
manpheation Rule for Independent Events	(4.7)	${}_{n}C_{x} = \frac{n!}{x!(n-x)!}$	(4.14)