

CS512 s23

HW#0

Khalid Saifullah
ID: A20423546

Answer to the Question No. A

Given,

$$a = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad b = \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}, \quad c = \begin{bmatrix} -1 \\ 1 \\ 3 \end{bmatrix}$$

A.1 $2a - b = 2 \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} - \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} = \begin{bmatrix} -2 \\ -1 \\ 0 \end{bmatrix}$

A.2 $\hat{a} = \frac{a}{\|a\|} = \frac{1}{\sqrt{1^2 + 2^2 + 3^2}} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \frac{1}{\sqrt{14}} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$

A.3 $\|a\| = \sqrt{1^2 + 2^2 + 3^2} = \sqrt{14}$

angle, $\theta = \cos^{-1}\left(\frac{1}{\sqrt{14}}\right) = 74.50^\circ$

A.4 The direction cosines of a are:

$$\alpha = \frac{1}{\sqrt{14}}, \quad \beta = \frac{2}{\sqrt{14}}, \quad \gamma = \frac{3}{\sqrt{14}}$$

A.5 The angle between a and $b = \cos^{-1}\left(\frac{a \cdot b}{\|a\| \|b\|}\right)$

$$= \cos^{-1}\left(\frac{1 \times 4 + 2 \times 5 + 3 \times 6}{\sqrt{14} \times \sqrt{77}}\right)$$
$$= 12.93^\circ$$

A.6 $a \cdot b = 1 \times 4 + 2 \times 5 + 3 \times 6 = 32$

$$b \cdot a = 4 \times 1 + 5 \times 2 + 6 \times 3 = 32$$

A.7

$$\cos \theta = \frac{a \cdot b}{\|a\| \|b\|}$$

$$\therefore a \cdot b = \|a\| \|b\| \cos \theta = \sqrt{14} \sqrt{77} \cos (12.93^\circ)$$

A.8

The scalar projection of b onto $\hat{a} = \frac{b \cdot a}{\|a\|}$

$$= b \cdot \frac{a}{\|a\|}$$

$$= b \cdot \hat{a}$$

$$= \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} \cdot \frac{1}{\sqrt{14}} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

$$= \frac{4 \times 1 + 5 \times 2 + 6 \times 3}{\sqrt{14}}$$

$$= \frac{32}{\sqrt{14}}$$

A.9 Let $V = [V_1 \ V_2 \ V_3]$ is a vector perpendicular to a . We know that the dot product of two orthogonal vector is 0.

$$\text{So, } 1 \times V_1 + 2 \times V_2 + 3 \times V_3 = 0$$

If $V_1 = 1$ and $V_2 = 1$, then

$$1 \times 1 + 2 \times 1 + 3 \times V_3 = 0$$

$$\therefore V_3 = -1$$

Hence, $V = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}$ is perpendicular to a .

A.10

$$a \times b = \begin{vmatrix} i & j & k \\ 1 & 2 & 3 \\ 4 & 5 & 6 \end{vmatrix} = (12 - 15)i - (6 - 12)j + (5 - 8)k$$
$$= \begin{bmatrix} -3 \\ 6 \\ -3 \end{bmatrix}$$

$$b \times a = \begin{vmatrix} i & j & k \\ 4 & 5 & 6 \\ 1 & 2 & 3 \end{vmatrix} = (15 - 12)i - (12 - 6)j + (8 - 5)k$$
$$= \begin{bmatrix} 3 \\ -6 \\ 3 \end{bmatrix}$$

11
The cross product of a and b produces a vector Q that is both perpendicular to the plane containing a and b . So,

$$Q = a \times b = \begin{bmatrix} -3 \\ 6 \\ -3 \end{bmatrix}$$

A.12

The linear dependence of a, b and c follows the relation $ax + by + cz = 0$

$$\begin{bmatrix} 1 & 4 & -1 \\ 2 & 5 & 1 \\ 3 & 6 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Forming the row-echelon form

$$\begin{bmatrix} 1 & 4 & -1 & 0 \\ 2 & 5 & 1 & 0 \\ 3 & 6 & 3 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 4 & -1 & 0 \\ 0 & -3 & 3 & 0 \\ 0 & -6 & 6 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 4 & -1 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 1 & -1 & 0 \end{bmatrix}$$

Therefore, a, b and c are not linearly independent.

$$\begin{bmatrix} 1 & 0 & 3 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \leftarrow \begin{bmatrix} 1 & 4 & -1 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

A.13

$$a^T b = \begin{bmatrix} 1 & 2 & 3 \end{bmatrix} \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} = 4 + 10 + 18 = 32$$

$$a b^T = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \begin{bmatrix} 4 & 5 & 6 \end{bmatrix} = \begin{bmatrix} 4 & 5 & 6 \\ 8 & 10 & 12 \\ 12 & 15 & 18 \end{bmatrix}$$

Answer to the Question B

Given,

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & -2 & 3 \\ 0 & 5 & -1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & -4 \\ 3 & -2 & 1 \end{bmatrix}, \quad C = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ -1 & 1 & 3 \end{bmatrix}$$

$$\text{and } d = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

$$\underline{\underline{B.1}} \quad 2A - B = \begin{bmatrix} 2 & 4 & 6 \\ 8 & -4 & 6 \\ 0 & 10 & -2 \end{bmatrix} - \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & -4 \\ 3 & -2 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 5 \\ 6 & -5 & 10 \\ -3 & 12 & -3 \end{bmatrix}$$

$$\underline{\underline{B.2}} \quad AB = \begin{bmatrix} 1 & 2 & 3 \\ 4 & -2 & 3 \\ 0 & 5 & -1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & -4 \\ 3 & -2 & 1 \end{bmatrix} = \begin{bmatrix} 14 & -2 & -4 \\ 9 & 0 & 15 \\ 7 & 7 & -21 \end{bmatrix}$$

$$BA = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & -4 \\ 3 & -2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & -2 & 3 \\ 0 & 5 & -1 \end{bmatrix} = \begin{bmatrix} 9 & 3 & 8 \\ 6 & -18 & 13 \\ -5 & 15 & 2 \end{bmatrix}$$

$$\underline{\underline{B.3}} \quad (AB)^T = \begin{bmatrix} 14 & 9 & 7 \\ -2 & 0 & 7 \\ -4 & 15 & -21 \end{bmatrix}$$

$$\underline{\underline{B.4}} \quad B^T A^T = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & -2 \\ 1 & -4 & 1 \end{bmatrix} \begin{bmatrix} 1 & 4 & 0 \\ 2 & -2 & 5 \\ 3 & 3 & -1 \end{bmatrix} = \begin{bmatrix} 14 & 9 & 7 \\ -2 & 0 & 7 \\ -4 & 15 & -21 \end{bmatrix}$$

$$\underline{\underline{B.4}} \quad |A| = \begin{vmatrix} 1 & 2 & 3 \\ 4 & -2 & 3 \\ 0 & 5 & -1 \end{vmatrix} = 1(2-15) - 2(-4-0) + 3(20-0) = 55$$

$$|C| = \begin{vmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ -1 & 1 & 3 \end{vmatrix} = 1(15-6) - 2(12+6) + 3(4+5) = 0$$

B.5 Two vectors are orthogonal if their inner product is zero.
For matrix A, row 1 (dot) row 2 = $\begin{bmatrix} 1 & 2 & 3 \end{bmatrix} \begin{bmatrix} 4 \\ -2 \\ 3 \end{bmatrix} = 9$
row 2 (dot) row 3 = $\begin{bmatrix} 4 & -2 & 3 \end{bmatrix} \begin{bmatrix} 0 \\ 5 \\ -1 \end{bmatrix} = -13$
row 1 (dot) row 3 = $\begin{bmatrix} 1 & 2 & 3 \end{bmatrix} \begin{bmatrix} 0 \\ 5 \\ -1 \end{bmatrix} = 7$

for matrix B, dot product of the rows pairwise yield 0.
So, the rows of the matrix B are normal to one another.

For the matrix C, dot product of the rows pairwise does not yield zero. So, the rows of the matrix C are not orthogonal to one another.

$$\underline{B.6} \quad A^{-1} = \frac{\text{adj}(A)}{|A|} = \frac{1}{55} \begin{bmatrix} -13 & 17 & 12 \\ 4 & -1 & 9 \\ 20 & -5 & -10 \end{bmatrix}$$

$$B^{-1} = \frac{\text{adj}(B)}{|B|} = \frac{1}{42} \begin{bmatrix} 7 & 4 & 9 \\ 14 & 2 & -6 \\ 7 & -8 & 3 \end{bmatrix}$$

$$\underline{B.7} \quad C^{-1} = \frac{\text{adj}(C)}{|C|}$$

As $|C| = 0$, the inverse of C doesn't exist.

$$\underline{B.8} \quad Ad = \begin{bmatrix} 1 & 2 & 3 \\ 4 & -2 & 3 \\ 0 & 5 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 14 \\ 9 \\ 7 \end{bmatrix}$$

$$\underline{B.9} \quad \text{row 1 (dot) } d = [1 \ 2 \ 3] \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = 14$$

$$\text{row 2 (dot) } d = [4 \ -2 \ 3] \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = 9$$

$$\text{row 3 (dot) } d = [0 \ 5 \ -1] \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = 7$$

$$\underline{B.10} \quad \begin{bmatrix} 1 \\ 4 \\ 0 \end{bmatrix} a_1 + \begin{bmatrix} 2 \\ -2 \\ 5 \end{bmatrix} a_2 + \begin{bmatrix} 3 \\ 3 \\ -1 \end{bmatrix} a_3 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & -2 & 3 \\ 0 & 5 & -1 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

Augmented matrix,

$$\left[\begin{array}{ccc|c} 1 & 2 & 3 & 1 \\ 4 & -2 & 3 & 2 \\ 0 & 5 & -1 & 3 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 2 & 3 & 1 \\ 0 & -10 & -9 & -2 \\ 0 & 0 & 11 & -4 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 2 & 3 & 1 \\ 0 & 1 & 9/10 & 1/5 \\ 0 & 0 & 1 & -4/11 \end{array} \right]$$

$$\beta_0, \quad a_1 = 24/25 \\ a_2 = 29/55 \\ a_3 = -4/11$$

$$\downarrow$$

$$\left[\begin{array}{ccc|c} 1 & 0 & 1.2 & 3/5 \\ 0 & 1 & 0.9 & 1/5 \\ 0 & 0 & 1 & -4/11 \end{array} \right]$$

$$\downarrow$$

$$\left[\begin{array}{ccc|c} 1 & 0 & 0 & 24/25 \\ 0 & 1 & 0 & 29/55 \\ 0 & 0 & 1 & -4/11 \end{array} \right]$$

B.11

$$Bx = d$$

$$\left[\begin{array}{ccc|c} 1 & 2 & 1 & 1 \\ 2 & 1 & -4 & 2 \\ 3 & -2 & 1 & 3 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 2 & 1 & 1 \\ 0 & -3 & -6 & 0 \\ 0 & -8 & -2 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 2 & 1 & 1 \\ 0 & 1 & 2 & 0 \\ 0 & 4 & 1 & 0 \end{array} \right]$$

$$\beta_0, \quad x = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

$$\downarrow$$

$$\left[\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right]$$

B.12

Matrix C is a non invertible matrix or singular matrix, so there is no solution.

Answer to the Question No. C

Given,

$$D = \begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix}, E = \begin{bmatrix} 2 & -2 \\ -2 & 5 \end{bmatrix}, F = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix}$$

C.1 To find the eigenvalues of D,

$$Dx = \lambda x$$

$$(D - \lambda I)x = 0$$

$$\text{So, } |D - \lambda I| = 0$$

$$\begin{vmatrix} 1-\lambda & 2 \\ 3 & 2-\lambda \end{vmatrix} = 0$$

$$(1-\lambda)(2-\lambda) - 6 = 0$$

$$\lambda^2 - 3\lambda - 4 = 0$$

$$\lambda^2 + \lambda - 4\lambda - 4 = 0$$

$$\lambda(\lambda+1) - 4(\lambda+1) = 0$$

$$(\lambda+1)(\lambda-4) = 0$$

$$\therefore \lambda = -1, 4$$

For $\lambda = -1$,

$$\begin{bmatrix} 2 & 2 \\ 3 & 3 \end{bmatrix}$$

$$\text{So, } u_1 = \begin{bmatrix} c_1 \\ -c_1 \end{bmatrix}$$

$$\text{let } c_1 = 1$$

$$\text{So, } u_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

For $\lambda = 4$,

$$\begin{bmatrix} -3 & 2 \\ 3 & -2 \end{bmatrix}$$

$$\text{So, } u_2 = \begin{bmatrix} \frac{2}{3}c_2 \\ c_2 \end{bmatrix}$$

$$\text{let } c_2 = 3$$

$$\text{So, } u_2 = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$$

$$\text{So, } x = [u_1 \ u_2] = \begin{bmatrix} 1 & 2 \\ -1 & 3 \end{bmatrix}$$

(Ans)

C.2

$$\begin{bmatrix} 1 & -1 \end{bmatrix} \begin{bmatrix} 2 \\ 3 \end{bmatrix} = -1$$

C.3

$$\begin{bmatrix} -0.89442719 & -0.4472136 \end{bmatrix} \begin{bmatrix} 0.4472136 \\ -0.89442719 \end{bmatrix} = 0$$

C.4 Since the dot product of the eigenvectors of \mathbb{E} is 0, they are orthogonal.

C.5
$$\begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Augmented matrix,

$$\left[\begin{array}{cc|c} 1 & 2 & 0 \\ 2 & 4 & 0 \end{array} \right] \rightarrow \left[\begin{array}{cc|c} 1 & 2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

The trivial solution is $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$

C.6 The two non trivial solutions are

$$u_1 = \begin{bmatrix} -2c_1 \\ c_1 \end{bmatrix} = \begin{bmatrix} -2 \\ 1 \end{bmatrix}$$

$$u_2 = \begin{bmatrix} c_2 \\ -\frac{1}{2}c_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$$

$$\text{So, } x = [u_1 \ u_2] = \begin{bmatrix} -2 & 2 \\ 1 & -1 \end{bmatrix}$$

C.7 The solution is $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$

As the determinant of D is non-zero, the homogeneous system of linear equations must have a unique (trivial) solution.

Answer to the Question No. D

Given, $f(x) = x^4 + 3$, $g(x) = x^2$, $q(x, y) = x^2 + y^2$

D.1

$$f'(x) = 2x$$

$$f''(x) = 2$$

D.2

$$\frac{\partial q}{\partial x} = 2x$$

$$\frac{\partial q}{\partial y} = 2y$$

D.3

$$\nabla q(x, y) = \begin{bmatrix} \frac{\partial q}{\partial x} \\ \frac{\partial q}{\partial y} \end{bmatrix} = \begin{bmatrix} 2x \\ 2y \end{bmatrix}$$

D.4

$$\frac{d}{dx} f(g(x)) = f'(g(x)) \times g'(x)$$

$$= f'(x^2) \times 2x$$

$$= \frac{d}{dx} (f(x^2)) \times 2x$$

$$= \frac{d}{dx} (x^4 + 3) \times 2x$$

$$= 4x^3 \times 2x$$

$$= 8x^4$$

$$\text{and } \frac{d}{dx} f(g(x)) = \frac{d}{dx} f(x^2)$$

$$= \frac{d}{dx} (x^4 + 3)$$

$$= 4x^3$$