

Isomeric states in ^{273}Ds and ^{269}Hs populated in the $^{238}\text{U}(^{40}\text{Ar}, 5n)$ reaction

Z. R. Sun,^{1,2} M. H. Huang,^{1,2,*} Z. G. Gan,^{1,2,3,†} H. B. Yang,^{1,2} J. G. Wang,¹ M. M. Zhang,^{1,4} Z. Y. Zhang,^{1,2} L. Ma,^{1,3} C. L. Yang,¹ X. L. Wu,^{1,3} Y. L. Tian,^{1,3} Y. S. Wang,^{1,3} J. Y. Wang,^{1,3} Y. H. Qiang,¹ L. C. Sun,^{1,5} X. Zhang,^{1,2} X. Y. Huang,^{1,2} Z. C. Li,^{1,2} L. Zhu,^{1,2} H. Zhou,^{1,2} G. Xie,^{1,2} J. H. Zheng,^{1,2} Z. L. Cheng,^{1,2} Z. H. Li,¹ W. X. Huang,^{1,2,3} Z. W. Lu,¹ Y. He,^{1,2} L. T. Sun,^{1,2} Z. Z. Ren,⁶ S. G. Zhou,^{2,7} V. K. Utyonkov,⁴ A. A. Voinov,⁴ Yu. S. Tsyganov,⁴ A. N. Polyakov,⁴ D. I. Solov'yev,⁴ N. D. Kovrizhnykh,⁴ and M. V. Shumeiko⁴

¹State Key Laboratory of Heavy Ion Science and Technology,
Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, China

²School of Nuclear Science and Technology, University of Chinese Academy of Sciences, Beijing 100049, China

³Advanced Energy Science and Technology Guangdong Laboratory, Huizhou 516000, China

⁴Joint Institute for Nuclear Research, RU-141980 Dubna, Russian Federation

⁵Shandong Provincial Key Laboratory of Nuclear Science,
Nuclear Energy Technology and Comprehensive Utilization,
Weihai Frontier Innovation Institute of Nuclear Technology,
School of Innovation Science, Energy and Power Engineering,
Shandong University, Shandong 250061, People's Republic of China

⁶School of Physics Science and Engineering, Tongji University, Shanghai 200092, China

⁷CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics,
Chinese Academy of Sciences, Beijing 100190, China

(Dated: April 22, 2026)

The $^{238}\text{U}(^{40}\text{Ar}, 5n)^{273}\text{Ds}$ reaction was investigated across an excitation-energy range of 40 to 51 MeV using the SHANS-2 separator. Upper cross-section limits of 833, 1118, and 511 fb were established at excitation energies of 40, 45, and 50 MeV, respectively. At an excitation energy of 49.5 MeV, two decay chains of ^{273}Ds were observed, yielding a production cross section of 219_{-147}^{+292} fb. The transitions observed in these new chains exhibit distinct energies and lifetimes, corroborating previous findings and confirming the existence of an isomeric state in ^{273}Ds in addition to the ground state. The measured cross section is in excellent agreement with the previously reported value of 180_{-120}^{+440} fb obtained at the DGFRS-2 separator at 49 MeV.

I. INTRODUCTION

Superactinides are generally considered to be elements with $Z \geq 104$. According to the liquid drop model, these elements should not exist [1]. Currently, the academic community believes that the existence of such superheavy elements is related to the nuclear shell structure. The predicted limits of nuclear survival lie at the closed nuclear shells of protons $Z = 108$ and neutrons $N = 162$ [2], as well as the emergence of heavier superheavy nuclei with $Z = 114 - 126$ and $N = 184$ [3]. As an odd-neutron nucleus with a single particle on the $N = 162$ shell, ^{273}Ds and its decay product ^{269}Hs , which is an odd-neutron nucleus with a hole below the $N = 161$ shell, are of great importance for gaining deeper insights into the $N = 162$ shell.

^{273}Ds was first unambiguously observed as a decay product in cold fusion experiments at the SHIP facility at GSI [4]. It is worth noting that in the cold fusion reaction $^{208}\text{Pb}(^{70}\text{Zn}, n)^{277}\text{Cn}$ conducted at GSI [4, 5] and RIKEN [6, 7], the lifetime of ^{273}Ds as a decay product is on the order of 0.1 ms. Last year, in the

$^{238}\text{U}(^{40}\text{Ar}, 5n)^{273}\text{Ds}$ experiment conducted by Yu. Ts. Oganessian *et al.* [8] at the DGFRS-2 separator [9], two decay chains were discovered where ^{273}Ds was produced directly. These were identified using the genetic correlation method [10]. One chain is consistent with historical data, while the other was assigned to an isomeric state of ^{273}Ds . Unfortunately, both chains are incomplete, and little is known about the decay information of ^{269}Hs .

Subsequently, Z. Wang *et al.* performed theoretical calculations on the decay pathways of these two chains using the density-dependent cluster model (DDCM) and the unified decay formula (UDF) [11], proposing minor modifications to these two chains. The present work was carried out at SHANS-2 [12], where two additional distinct decay pathways were observed using the same reaction. The pathway information is complete and consistent with historical data through the genetic correlation method [10], which fully and independently confirms the authenticity of the two states of ^{273}Ds ($^{273}\text{Ds}^a$ and $^{273}\text{Ds}^b$) observed by Yu. Ts. Oganessian *et al.* [8].

The experiment is described below in Table I, and the experimental data are presented.

* Corresponding author: hmh@impcas.ac.cn

† Corresponding author: zggan@impcas.ac.cn

TABLE I. Experimental parameters and cross sections for the $^{238}\text{U} + ^{40}\text{Ar}$ reaction.

Reaction	Target thickness (mg/cm ²)	E_{lab} (MeV)	E^* (MeV)	Beam dose $\times 10^{19}$	No. of chains 4n/5n	σ_{4n} (pb)	σ_{5n} (pb)
$^{238}\text{U} + ^{40}\text{Ar}$	0.47 / 0.60	212.7	50.6	1.95 / 1.45	0/2	< 0.43	$0.15^{+0.20}_{-0.10}$
	0.47	207.4	46.2	0.82	0/0	< 0.9	< 0.9
	0.60	204.0	43.6	2.04	0/0	< 0.3	< 0.3
	0.60	214.3	50.9	1.527	0/0	< 0.4	< 0.4

- [1] O. Hahn and F. Strassmann, Über den nachweis und das verhalten der bei der bestrahlung des urans mittels neutronen entstehenden erdalkalimetalle, in *Das Neutron*, edited by B. M. Kedrow (De Gruyter, 1939) pp. 257–262.
- [2] R. Smolańczuk, J. Skalski, and A. Sobiczewski, Spontaneous-fission half-lives of deformed superheavy nuclei, *Physical Review C* **52**, 1871 (1995).
- [3] Y. T. Oganessian and K. P. Rykaczewski, A beachhead on the island of stability, *Physics Today* **68**, 32 (2015).
- [4] S. Hofmann, V. Ninov, F. P. Heßberger, P. Armbruster, H. Folger, G. Münzenberg, H. J. Schött, A. G. Popeko, A. V. Yereimin, S. Saro, R. Janik, and M. Leino, The new element 112, *Zeitschrift für Physik A Hadrons and Nuclei* **354**, 229 (1996).
- [5] S. Hofmann, F. Heßberger, D. Ackermann, G. Münzenberg, S. Antalic, P. Cagarda, B. Kindler, J. Kojouharova, M. Leino, B. Lommel, R. Mann, A. Popeko, S. Reshitko, S. Šaro, J. Uusitalo, and A. Yereimin, New results on elements 111 and 112, *The European Physical Journal A* **14**, 147 (2002).
- [6] K. Morita, K. Morimoto, D. Kaji, T. Akiyama, S.-i. Goto, H. Haba, E. Ideguchi, K. Katori, H. Koura, H. Kudo, T. Ohnishi, A. Ozawa, T. Suda, K. Sueki, F. Tokanai, T. Yamaguchi, A. Yoneda, and A. Yoshida, Experiment on Synthesis of an Isotope²⁷⁷ 112 by²⁰⁸ Pb+⁷⁰ Zn Reaction, *Journal of the Physical Society of Japan* **76**, 043201 (2007).
- [7] T. Sumita, K. Morimoto, D. Kaji, H. Haba, K. Ozeki, R. Sakai, A. Yoneda, A. Yoshida, H. Hasebe, K. Katori, N. Sato, Y. Wakabayashi, S.-i. Mitsuoka, S.-i. Goto, M. Murakami, Y. Kariya, F. Tokanai, K. Mayama, M. Takeyama, T. Moriya, E. Ideguchi, T. Yamaguchi, H. Kikunaga, J. Chiba, and K. Morita, New Result on the Production of²⁷⁷ Cn by the²⁰⁸ Pb +⁷⁰ Zn Reaction, *Journal of the Physical Society of Japan* **82**, 024202 (2013).
- [8] Yu. Ts. Oganessian, V. K. Utyonkov, M. V. Shumeiko, F. Sh. Abdullin, G. G. Adamian, S. N. Dmitriev, D. Ibadullayev, M. G. Itkis, N. D. Kovrizhnykh, D. A. Kuznetsov, O. V. Petrushkin, A. V. Podshibiakin, A. N. Polyakov, A. G. Popeko, I. S. Rogov, R. N. Sagaidak, L. Schlattauer, V. D. Shubin, D. I. Soloviyev, Yu. S. Tsyganov, A. A. Voinov, V. G. Subbotin, N. S. Bublikova, M. G. Voronyuk, A. V. Sabelnikov, A. Yu. Broduv, N. V. Aksenov, A. V. Khalkin, Z. G. Gan, Z. Y. Zhang, M. H. Huang, and H. B. Yang, Synthesis and decay properties of isotopes of element 110: Ds 273 and Ds 275, *Physical Review C* **109**, 054307 (2024).
- [9] Yu. Ts. Oganessian, V. Utyonkov, A. Popeko, D. Soloviyev, F. Sh. Abdullin, S. Dmitriev, D. Ibadullayev, M. Itkis, N. Kovrizhnykh, D. Kuznetsov, O. Petrushkin, A. Podshibiakin, A. Polyakov, R. Sagaidak, L. Schlattauer, I. Shirokovsky, V. Shubin, M. Shumeiko, Yu. S. Tsyganov, A. Voinov, V. Subbotin, V. Bekhterev, N. Belykh, O. Chernyshev, K. Gikal, G. Ivanov, A. Khalkin, V. Konstantinov, N. Osipov, S. Paschenko, A. Protasov, V. Semin, V. Sorokoumov, K. Sychev, V. Verevchkin, B. Yakovlev, S. Antoine, W. Beeckman, P. Jehanno, M. Yavor, A. Shcherbakov, K. Rykaczewski, T. King, J. Roberto, N. Brewer, R. Grzywacz, Z. Gan, Z. Zhang, M. Huang, and H. Yang, DGFRS-2—A gas-filled recoil separator for the Dubna Super Heavy Element Factory, *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* **1033**, 166640 (2022).
- [10] S. Hofmann, S. N. Dmitriev, C. Fahlander, J. M. Gates, J. B. Roberto, and H. Sakai, On the discovery of new elements (IUPAC/IUPAP Provisional Report): Provisional Report of the 2017 Joint Working Group of IUPAC and IUPAP, *Pure and Applied Chemistry* **90**, 1773 (2018).
- [11] Z. Wang and Z. Ren, Investigating α decays of recently discovered heavy and superheavy nuclei within the improved density-dependent cluster model, *Physical Review C* **110**, 064307 (2024).
- [12] L. Sheng, Q. Hu, H. Jia, Z. Zhang, Z. Chai, B. Zhao, Y. Zhang, Z. Gan, Y. He, and J. Yang, Ion-optical design and multiparticle tracking in 3D magnetic field of the gas-filled recoil separator SHANS2 at CAFE2, *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* **1004**, 165348 (2021).