

AMC 12/AHSME 1963
www.artofproblemsolving.com/community/c4827

by TheMaskedMagician, rrusczyk

1 Which one of the following points is not on the graph of $y = \frac{x}{x+1}$?

- (A) $(0, 0)$ (B) $\left(-\frac{1}{2}, -1\right)$ (C) $\left(\frac{1}{2}, \frac{1}{3}\right)$ (D) $(-1, 1)$ (E) $(-2, 2)$

2 Let $n = x - y^{x-y}$. Find n when $x = 2$ and $y = -2$.

- (A) -14 (B) 0 (C) 1 (D) 18 (E) 256

3 If the reciprocal of $x + 1$ is $x - 1$, then x equals:

- (A) 0 (B) 1 (C) -1 (D) ± 1 (E) none of these

4 For what value(s) of k does the pair of equations $y = x^2$ and $y = 3x + k$ have two identical solutions?

- (A) $\frac{4}{9}$ (B) $-\frac{4}{9}$ (C) $\frac{9}{4}$ (D) $-\frac{9}{4}$ (E) $\pm \frac{9}{4}$

5 If x and $\log_{10} x$ are real numbers and $\log_{10} x < 0$, then:

- (A) $x < 0$ (B) $-1 < x < 1$ (C) $0 < x \leq 1$
 (D) $-1 < x < 0$ (E) $0 < x < 1$

6 Triangle BAD is right-angled at B . On AD there is a point C for which $AC = CD$ and $AB = BC$. The magnitude of angle DAB , in degrees, is:

- (A) $67\frac{1}{2}$ (B) 60 (C) 45 (D) 30 (E) $22\frac{1}{2}$

7 Given the four equations:

- (1) $3y - 2x = 12$ (2) $-2x - 3y = 10$ (3) $3y + 2x = 12$ (4) $2y + 3x = 10$

The pair representing the perpendicular lines is:

- (A) (1) and (4) (B) (1) and (3) (C) (1) and (2) (D) (2) and (4) (E) (2) and (3)


8 The smallest positive integer x for which $1260x = N^3$, where N is an integer, is:

- (A) 1050 (B) 1260 (C) 1260^2 (D) 7350 (E) 44100

- 9 In the expansion of $\left(a - \frac{1}{\sqrt{a}}\right)^7$ the coefficient of $a^{-\frac{1}{2}}$ is:
 (A) -7 (B) 7 (C) -21 (D) 21 (E) 35
-
- 10 Point P is taken interior to a square with side-length a and such that it is equally distant from two consecutive vertices and from the side opposite these vertices. If d represents the common distance, then d equals:
 (A) $\frac{3a}{5}$ (B) $\frac{5a}{8}$ (C) $\frac{3a}{8}$ (D) $\frac{a\sqrt{2}}{2}$ (E) $\frac{a}{2}$
-
- 11 The arithmetic mean of a set of 50 numbers is 38. If two numbers of the set, namely 45 and 55, are discarded, the arithmetic mean of the remaining set of numbers is:
 (A) 38.5 (B) 37.5 (C) 37 (D) 36.5 (E) 36
-
- 12 Three vertices of parallelogram $PQRS$ are $P(-3, -2)$, $Q(1, -5)$, $R(9, 1)$ with P and R diagonally opposite. The sum of the coordinates of vertex S is:
 (A) 13 (B) 12 (C) 11 (D) 10 (E) 9
-
- 13 If $2^a + 2^b = 3^c + 3^d$, the number of integers a, b, c, d which can possibly be negative, is, at most:
 (A) 4 (B) 3 (C) 2 (D) 1 (E) 0
-
- 14 Given the equations $x^2 + kx + 6 = 0$ and $x^2 - kx + 6 = 0$. If, when the roots of the equation are suitably listed, each root of the second equation is 5 more than the corresponding root of the first equation, then k equals:
 (A) 5 (B) -5 (C) 7 (D) -7 (E) none of these
-
- 15 A circle is inscribed in an equilateral triangle, and a square is inscribed in the circle. The ratio of the area of the triangle to the area of the square is:
 (A) $\sqrt{3} : 1$ (B) $\sqrt{3} : \sqrt{2}$ (C) $3\sqrt{3} : 2$ (D) $3 : \sqrt{2}$ (E) $3 : 2\sqrt{2}$
-
- 16 Three numbers a, b, c , none zero, form an arithmetic progression. Increasing a by 1 or increasing c by 2 results in a geometric progression. Then b equals:
 (A) 16 (B) 14 (C) 12 (D) 10 (E) 8
-
- 17 The expression $\frac{\frac{a}{a+y} + \frac{y}{a-y}}{\frac{y}{a+y} - \frac{a}{a-y}}$, a real, $a \neq 0$, has the value -1 for:

- (A) all but two real values of y (B) only two real values of y
 (C) all real values of y (D) only one real value of y (E) no real values of y

- 18 Chord EF is the perpendicular bisector of chord BC , intersecting it in M . Between B and M point U is taken, and EU extended meets the circle in A . Then, for any selection of U , as described, triangle EUM is similar to triangle:



- (A) EFA (B) EFC (C) ABM (D) ABU (E) FMC

- 19 In counting n colored balls, some red and some black, it was found that 49 of the first 50 counted were red. Thereafter, 7 out of every 8 counted were red. If, in all, 90% or more of the balls counted were red, the maximum value of n is:

- (A) 225 (B) 210 (C) 200 (D) 180 (E) 175

- 20 Two men at points R and S , 76 miles apart, set out at the same time to walk towards each other. The man at R walks uniformly at the rate of $4\frac{1}{2}$ miles per hour; the man at S walks at the constant rate of $3\frac{1}{4}$ miles per hour for the first hour, at $3\frac{3}{4}$ miles per hour for the second hour, and so on, in arithmetic progression. If the men meet x miles nearer R than S in an integral number of hours, then x is:

- (A) 10 (B) 8 (C) 6 (D) 4 (E) 2

- 21 The expression $x^2 - y^2 - z^2 + 2yz + x + y - z$ has:

- (A) no linear factor with integer coefficients and integer exponents
 (B) the factor $-x + y + z$
 (C) the factor $x - y - z + 1$

(D) the factor $x + y - z + 1$

(E) the factor $x - y + z + 1$

- 22 Acute-angled triangle ABC is inscribed in a circle with center at O ; $\widehat{AB} = 120$ and $\widehat{BC} = 72$. A point E is taken in minor arc AC such that OE is perpendicular to AC . Then the ratio of the magnitudes of angles OBE and BAC is:

(A) $\frac{5}{18}$ (B) $\frac{2}{9}$ (C) $\frac{1}{4}$ (D) $\frac{1}{3}$ (E) $\frac{4}{9}$


- 23 A gives B as many cents as B has and C as many cents as C has. Similarly, B then gives A and C as many cents as each then has. C , similarly, then gives A and B as many cents as each then has. If each finally has 16 cents, with how many cents does A start?

(A) 24 (B) 26 (C) 28 (D) 30 (E) 32

- 24 Consider equations of the form $x^2 + bx + c = 0$. How many such equations have real roots and have coefficients b and c selected from the set of integers $\{1, 2, 3, 4, 5, 6\}$?

(A) 20 (B) 19 (C) 18 (D) 17 (E) 16

- 25 Point F is taken in side AD of square $ABCD$. At C a perpendicular is drawn to CF , meeting AB extended at E . The area of $ABCD$ is 256 square inches and the area of triangle CEF is 200 square inches. Then the number of inches in BE is:



(A) 12 (B) 14 (C) 15 (D) 16 (E) 20

- 26 **Form 1**
Consider the statements:

(1) $p \wedge \sim q \wedge r$ (2) $\sim p \wedge \sim q \wedge r$ (3) $p \wedge \sim q \wedge \sim r$ (4) $\sim p \wedge q \wedge r$,

where p, q , and r are propositions. How many of these imply the truth of $(p \rightarrow q) \rightarrow r$?

- (A) 0 (B) 1 (C) 2 (D) 3 (E) 4

Form 2

Consider the statements (1) p and r are true and q is false (2) r is true and p and q are false (3) p is true and q and r are false (4) q and r are true and p is false. How many of these imply the truth of the statement

" r is implied by the statement that p implies q "?

- (A) 0 (B) 1 (C) 2 (D) 3 (E) 4

-
- 27** Six straight lines are drawn in a plane with no two parallel and no three concurrent. The number of regions into which they divide the plane is:

- (A) 16 (B) 20 (C) 22 (D) 24 (E) 26

-
- 28** Given the equation $3x^2 - 4x + k = 0$ with real roots. The value of k for which the product of the roots of the equation is a maximum is:

- (A) $\frac{16}{9}$ (B) $\frac{16}{3}$ (C) $\frac{4}{9}$ (D) $\frac{4}{3}$ (E) $-\frac{4}{3}$

-
- 29** A particle projected vertically upward reaches, at the end of t seconds, an elevation of s feet where $s = 160t - 16t^2$. The highest elevation is:

- (A) 800 (B) 640 (C) 400 (D) 320 (E) 160

-
- 30** Let

$$F = \log \frac{1+x}{1-x}.$$

Find a new function G by replacing each x in F by

$$\frac{3x+x^3}{1+3x^2},$$

and simplify. The simplified expression G is equal to:

- (A) $-F$ (B) F (C) $3F$ (D) F^3 (E) $F^3 - F$

-
- 31** The number of solutions in positive integers of $2x + 3y = 763$ is:

- (A) 255 (B) 254 (C) 128 (D) 127 (E) 0

-
- 32** The dimensions of a rectangle R are a and b , $a < b$. It is required to obtain a rectangle with dimensions x and y , $x < a$, $y < a$, so that its perimeter is one-third that of R , and its area is one-third that of R . The number of such (different) rectangles is:

(A) 0 (B) 1 (C) 2 (D) 4 (E) infinitely many

- 33 Given the line $y = \frac{3}{4}x + 6$ and a line L parallel to the given line and 4 units from it. A possible equation for L is:

(A) $y = \frac{3}{4}x + 1$ (B) $y = \frac{3}{4}x$ (C) $y = \frac{3}{4}x - \frac{2}{3}$

(D) $y = \frac{3}{4}x - 1$ (E) $y = \frac{3}{4}x + 2$

- 34 In triangle ABC, side $a = \sqrt{3}$, side $b = \sqrt{3}$, and side $c > 3$. Let x be the largest number such that the magnitude, in degrees, of the angle opposite side c exceeds x . Then x equals:

(A) 150 (B) 120 (C) 105 (D) 90 (E) 60

- 35 The lengths of the sides of a triangle are integers, and its area is also an integer. One side is 21 and the perimeter is 48. The shortest side is:

(A) 8 (B) 10 (C) 12 (D) 14 (E) 16

- 36 A person starting with 64 cents and making 6 bets, wins three times and loses three times, the wins and losses occurring in random order. The chance for a win is equal to the chance for a loss. If each wager is for half the money remaining at the time of the bet, then the final result is:

(A) a loss of 27 (B) a gain of 27 (C) a loss of 37

(D) neither a gain nor a loss (E) a gain or a loss depending upon the order in which the wins and losses occur

Note: Due to the lack of \LaTeX packages, the numbers in the answer choices are in cents

- 37 Given points P_1, P_2, \dots, P_7 on a straight line, in the order stated (not necessarily evenly spaced). Let P be an arbitrarily selected point on the line and let s be the sum of the undirected lengths


$$PP_1, PP_2, \dots, PP_7.$$

Then s is smallest if and only if the point P is:

(A) midway between P_1 and P_7 (B) midway between P_2 and P_6 (C) midway between P_3 and P_5

(D) at P_4 (E) at P_1

- 38 Point F is taken on the extension of side AD of parallelogram $ABCD$. BF intersects diagonal AC at E and side DC at G . If $EF = 32$ and $GF = 24$, then BE equals:




- (A) 4 (B) 8 (C) 10 (D) 12 (E) 16

- 39 In triangle ABC lines CE and AD are drawn so that

$$\frac{CD}{DB} = \frac{3}{1} \text{ and } \frac{AE}{EB} = \frac{3}{2}. \text{ Let } r = \frac{CP}{PE}$$

where P is the intersection point of CE and AD . Then r equals:



- (A) 3 (B) $\frac{3}{2}$ (C) 4 (D) 5 (E) $\frac{5}{2}$

- 40 If x is a number satisfying the equation $\sqrt[3]{x+9} - \sqrt[3]{x-9} = 3$, then x^2 is between:

- (A) 55 and 65 (B) 65 and 75 (C) 75 and 85 (D) 85 and 95 (E) 95 and 105



— These problems are copyright © Mathematical Association of America (<http://maa.org>).
