

**AMC 12/AHSME 1975**
[www.artofproblemsolving.com/community/c4839](http://www.artofproblemsolving.com/community/c4839)

by ernie, dft, rrusczyk

- 1 The value of  $\frac{1}{2 - \frac{1}{2 - \frac{1}{2 - \frac{1}{2}}}}$  is
- (A)  $3/4$  (B)  $4/5$  (C)  $5/6$  (D)  $6/7$  (E)  $6/5$

- 2 For which real values of  $m$  are the simultaneous equations

$$y = mx + 3$$

$$y = (2m - 1)x + 4$$

 satisfied by at least one pair of real numbers  $(x, y)$ ?

- (A) all  $m$  (B) all  $m \neq 0$  (C) all  $m \neq 1/2$  (D) all  $m \neq 1$  (E) no values of  $m$
- 3 Which of the following inequalities are satisfied for all real numbers  $a, b, c, x, y, z$  which satisfy the conditions  $x < a, y < b$ , and  $z < c$ ?
- I.  $xy + yz + zx < ab + bc + ca$  II.  $x^2 + y^2 + z^2 < a^2 + b^2 + c^2$  III.  $xyz < abc$
- (A) None are satisfied. (B) I only (C) II only (D) III only (E) All are satisfied.

- 4 If the side of one square is the diagonal of a second square, what is the ratio of the area of the first square to the area of the second?
- (A) 2 (B)  $\sqrt{2}$  (C)  $1/2$  (D)  $2\sqrt{2}$  (E) 4

- 5 The polynomial  $(x + y)^9$  is expanded in decreasing powers of  $x$ . The second and third terms have equal values when evaluated at  $x = p$  and  $y = q$ , where  $p$  and  $q$  are positive numbers whose sum is one. What is the value of  $p$ ?
- (A)  $1/5$  (B)  $4/5$  (C)  $1/4$  (D)  $3/4$  (E)  $8/9$

- 6 The sum of the first eighty positive odd integers subtracted from the sum of the first eighty positive even integers is
- (A) 0 (B) 20 (C) 40 (D) 60 (E) 80

- 7 For which non-zero real numbers  $x$  is  $\frac{|x-|x||}{x}$  a positive integer?
- (A) for negative  $x$  only (B) for positive  $x$  only (C) only for  $x$  an even integer (D) for all non-zero  $x$  (E) for no non-zero real numbers  $x$

- 8 If the statement "All shirts in this store are on sale." is false, then which of the following statements must be true?

- I. All shirts in this store are at non-sale prices.  
 II. There is some shirt in this store not on sale.  
 III. No shirt in this store is on sale.  
 IV. Not all shirts in this store are on sale.

(A) II only    (B) IV only    (C) I and III only    (D) II and IV only    (E) I, II and IV only


- 
- 9 Let  $a_1, a_2, \dots$  and  $b_1, b_2, \dots$  be arithmetic progressions such that  $a_1 = 25$ ,  $b_1 = 75$ , and  $a_{100} + b_{100} = 100$ . Find the sum of the first hundred terms of the progression  $a_1 + b_1, a_2 + b_2, \dots$   
 (A) 0    (B) 100    (C) 10,000    (D) 505,000    (E) not enough information given to solve the problem
- 
- 10 The sum of the digits in base ten of  $(10^{4n^2+8} + 1)^2$ , where  $n$  is a positive integer, is  
 (A) 4    (B)  $4n$     (C)  $2 + 2n$     (D)  $4n^2$     (E)  $n^2 + n + 2$
- 
- 11 Let  $P$  be an interior point of circle  $K$  other than the center of  $K$ . Form all chords of  $K$  which pass through  $P$ , and determine their midpoints. The locus of these midpoints is  
 (A) a circle with one point deleted    (B) a circle if the distance from  $P$  to the center of  $K$  is less than one half the radius of  $K$ ; otherwise a circular arc of less than  $360^\circ$     (C) a semicircle with one point deleted  
 (D) a semicircle    (E) a circle
- 
- 12 If  $a \neq b$ ,  $a^3 - b^3 = 19x^3$ , and  $a - b = x$ , which of the following conclusions is correct?  
 (A)  $a = 3x$     (B)  $a = 3x$  or  $a = -2x$     (C)  $a = -3x$  or  $a = 2x$     (D)  $a = 3x$  or  $a = 2x$   
 (E)  $a = 2x$
- 
- 13 The equation  $x^6 - 3x^5 - 6x^3 - x + 8$  has  
 (A) no real roots    (B) exactly two distinct negative roots    (C) exactly one negative root  
 (D) no negative roots, but at least one positive root    (E) none of these
- 
- 14 If the *whatsis* is *so* when the *whosis* is *is* and the *so* and *so* is *is* · *so*, what is the *whosis* · *whatsis* when the *whosis* is *so*, the *so* and *so* is *so* · *so* and the *is* is two (*whatsis*, *whosis*, *is* and *so* are variables taking positive values)?  
 (A) *whosis* · *is* · *so*    (B) *whosis*    (C) *is*    (D) *so*    (E) *so* and *so*
- 
- 15 In the sequence of numbers 1, 3, 2, ... each term after the first two is equal to the term preceding it minus the term preceding that. The sum of the first one hundred terms of the sequence is  
 (A) 5    (B) 4    (C) 2    (D) 1    (E) -1
- 
- 16 If the first term of an infinite geometric series is a positive integer, the common ratio is the reciprocal of a positive integer, and the sum of the series is 3, then the sum of the first two terms of the series is  
 (A)  $1/3$     (B)  $2/3$     (C)  $8/3$     (D) 2    (E)  $9/2$
-

- 17 A man can commute either by train or by bus. If he goes to work on the train in the morning, he comes home on the bus in the afternoon; and if he comes home in the afternoon on the train, he took the bus in the morning. During a total of  $x$  working days, the man took the bus to work in the morning 8 times, came home by bus in the afternoon 15 times, and commuted by train (either morning or afternoon) 9 times. Find  $x$ .  
 (A) 19 (B) 18 (C) 17 (D) 16 (E) not enough information given to solve the problem

- 18 A positive integer  $N$  with three digits in its base ten representation is chosen at random, with each three digit number having an equal chance of being chosen. The probability that  $\log_2 N$  is an integer is  
 (A) 0 (B)  $3/899$  (C)  $1/225$  (D)  $1/300$  (E)  $1/450$

- 19 Which positive numbers  $x$  satisfy the equation  $(\log_3 x)(\log_x 5) = \log_3 5$ ?  
 (A) 3 and 5 only (B) 3, 5, and 15 only (C) only numbers of the form  $5^n \cdot 3^m$ , where  $n$  and  $m$  are positive integers (D) all positive  $x \neq 1$  (E) none of these


- 20 In the adjoining figure triangle  $ABC$  is such that  $AB = 4$  and  $AC = 8$ . If  $M$  is the midpoint of  $BC$  and  $AM = 3$ , what is the length of  $BC$ ?  
 (A)  $2\sqrt{26}$  (B)  $2\sqrt{31}$  (C) 9 (D)  $4+2\sqrt{13}$  (E) not enough information given to solve the problem



- 21 Suppose  $f(x)$  is defined for all real numbers  $x$ ;  $f(x) > 0$  for all  $x$ , and  $f(a)f(b) = f(a+b)$  for all  $a$  and  $b$ . Which of the following statements is true?  
 I.  $f(0) = 1$  II.  $f(-a) = 1/f(a)$  for all  $a$  III.  $f(a) = \sqrt[3]{f(3a)}$  for all  $a$  IV.  $f(b) > f(a)$  if  $b > a$   
 (A) III and IV only (B) I, III, and IV only (C) I, II, and IV only (D) I, II, and III only  
 (E) All are true.

- 22 If  $p$  and  $q$  are primes and  $x^2 - px + q = 0$  has distinct positive integral roots, then which of the following statements are true?  
 I. The difference of the roots is odd. II. At least one root is prime. III.  $p^2 - q$  is prime. IV.  $p+q$  is prime.  
 (A) I only (B) II only (C) II and III only (D) I, II and IV only (E) All are true.

- 23 In the adjoining figure  $AB$  and  $BC$  are adjacent sides of square  $ABCD$ ;  $M$  is the midpoint of  $AB$ ;  $N$  is the midpoint of  $BC$ ; and  $AN$  and  $CM$  intersect at  $O$ . The ratio of the area of  $AOCD$  to the area of  $ABCD$  is



- (A)  $\frac{5}{6}$     (B)  $\frac{3}{4}$     (C)  $\frac{2}{3}$     (D)  $\frac{\sqrt{3}}{2}$     (E)  $\frac{(\sqrt{3}-1)}{2}$

- 24** In triangle  $ABC$ ,  $\angle C = \theta$  and  $\angle B = 2\theta$ , where  $0^\circ < \theta < 60^\circ$ . The circle with center  $A$  and radius  $AB$  intersects  $AC$  at  $D$  and intersects  $BC$ , extended if necessary, at  $B$  and at  $E$  ( $E$  may coincide with  $B$ ). Then  $EC = AD$

- (A) for no values of  $\theta$     (B) only if  $\theta = 45^\circ$     (C) only if  $0^\circ < \theta \leq 45^\circ$   
 (D) only if  $45^\circ \leq \theta < 60^\circ$     (E) for all  $\theta$  such that  $0^\circ < \theta < 60^\circ$

- 25** A woman, her brother, her son and her daughter are chess players (all relations by birth). The worst player's twin (who is one of the four players) and the best player are of opposite sex. The worst player and the best player are the same age. Who is the worst player?

- (A) the woman    (B) her son    (C) her brother    (D) her daughter  
 (E) No solution is consistent with the given information


- 26** In acute triangle  $ABC$  the bisector of  $\angle A$  meets side  $BC$  at  $D$ . The circle with center  $B$  and radius  $BD$  intersects side  $AB$  at  $M$ ; and the circle with center  $C$  and radius  $CD$  intersects side  $AC$  at  $N$ . Then it is always true that

- (A)  $\angle CND + \angle BMD - \angle DAC = 120^\circ$     (B)  $AMDN$  is a trapezoid    (C)  $BC$  is parallel to  $MN$   
 (D)  $AM - AN = \frac{3(DB-DC)}{2}$     (E)  $AB - AC = \frac{3(DB-DC)}{2}$

- 27** If  $p, q$  and  $r$  are distinct roots of  $x^3 - x^2 + x - 2 = 0$ , then  $p^3 + q^3 + r^3$  equals

- (A)  $-1$     (B)  $1$     (C)  $3$     (D)  $5$     (E) none of these

- 28** In triangle  $ABC$  shown in the adjoining figure,  $M$  is the midpoint of side  $BC$ ,  $AB = 12$  and  $AC = 16$ . Points  $E$  and  $F$  are taken on  $AC$  and  $AB$ , respectively, and lines  $EF$  and  $AM$  intersect at  $G$ . If  $AE = 2AF$  then  $\frac{EG}{GF}$  equals



- (A)  $\frac{3}{2}$     (B)  $\frac{4}{3}$     (C)  $\frac{5}{4}$     (D)  $\frac{6}{5}$   
 (E) not enough information to solve the problem

29 What is the smallest integer larger than  $(\sqrt{3} + \sqrt{2})^6$ ?

- (A) 972    (B) 971    (C) 970    (D) 969    (E) 968

30 Let  $x = \cos 36^\circ - \cos 72^\circ$ . Then  $x$  equals

- (A)  $\frac{1}{3}$     (B)  $\frac{1}{2}$     (C)  $3 - \sqrt{6}$     (D)  $2\sqrt{3} - 3$     (E) none of these



These problems are copyright © Mathematical Association of America (<http://maa.org>).