

Controlled Release of Madecassoside and Asiaticoside of *Centella asiatica* L. Origin from Sustainable Cold-Processed Topical Formulations

Monika Krzyżośtan ^{1,2}, Agata Wawrzyńczak ^{1,*} and Izabela Nowak ^{1,*}

¹ Faculty of Chemistry, Adam Mickiewicz University in Poznań, Uniwersytetu Poznańskiego 8, 61-614 Poznań, Poland; monika.krzyzostan@amu.edu.pl

² Dr Koziej Instytut Badań Kosmetyków, Czerniakowska 58, 00-717 Warsaw, Poland

* Correspondence: agata.wawrzynczak@amu.edu.pl (A.W.), nowakiza@amu.edu.pl (I.N.)

Supporting Information




Figure S1. Photographs of the prepared topical formulations containing 10 wt.% of *Centella asiatica* L. extract.




A**B**

Figure S2. Calibration curves for HPLC analysis of madecassoside (A) and asiaticoside (B).

(A) Madecassoside



(B) Asiaticoside





Figure S3. Graphic representation of kinetic models applied to the release profiles from 2 % M+A – O/W sample.

(A) Madecassoside



(B) Asiaticoside





Figure S4. Graphic representation of kinetic models applied to the release profiles from 2 % M+A – W/O sample.

(A) Madecassoside



(B) Asiaticoside





Figure S5. Graphic representation of kinetic models applied to the release profiles from 2 % M+A – G sample.

(A) Madecassoside



(B) Asiaticoside





Figure S6. Graphic representation of kinetic models applied to the release profiles from 5 % M+A – O/W sample.

(A) Madecassoside



(B) Asiaticoside





Figure S7. Graphic representation of kinetic models applied to the release profiles from 5 % M+A – W/O sample.

(A) Madecassoside



(B) Asiaticoside




Figure S8. Graphic representation of kinetic models applied to the release profiles from 5 % M+A – G sample.