

LETTER

Tree diversity reduces herbivory by forest insects

Hervé Jactel¹* and Eckehard G. Brockerhoff²

¹INRA, UMR1202 Biodiversity, Genes & Communities, Laboratory of Forest Entomology and Biodiversity, 69 Route d'Arcachon, 33612 Cestas Cedex, France

²Ensis^a, PO Box 29237, Christchurch 850, New Zealand

*Correspondence: E-mail: herve.jactel@pierroton.inra.fr

Abstract

Biodiversity loss from plant communities is often acknowledged to affect primary production but little is known about effects on herbivores. We conducted a meta-analysis of a worldwide data set of 119 studies to compare herbivory in single-species and mixed forests. This showed a significant reduction of herbivory in more diverse forests but this varied with the host specificity of insects. In diverse forests, herbivory by oligophagous species was virtually always reduced, whereas the response of polyphagous species was variable. Further analyses revealed that the composition of tree mixtures may be more important than species richness *per se* because diversity effects on herbivory were greater when mixed forests comprised taxonomically more distant tree species, and when the proportion of non-host trees was greater than that of host trees. These findings provide new support for the role of biodiversity in ecosystem functioning across trophic levels.

Keywords

Biodiversity, ecosystem functioning, forest, herbivory, insect, meta-analysis.

Ecology Letters (2007) 10: 835–848

INTRODUCTION

The loss of biodiversity, mainly due to human activities, is a major environmental concern (United Nations 1992; Balmford & Bond 2005). Due to the combined effects of deforestation, forest fragmentation, declining habitat quality and climate change (Saunders *et al.* 1991; Lindenmayer & Franklin 2002; Brook *et al.* 2003; Thomas *et al.* 2004), biodiversity loss is particularly serious in forests, which appear to harbour more than half of the known terrestrial plant and animal species (Hassan *et al.* 2005). Apart from concerns for the conservation of biodiversity, the unprecedented loss of species may have dramatic detrimental effects on the functioning of ecosystems (Hooper *et al.* 2005). Although the large number of species involved in ecosystem processes and the complexity of their interactions make it difficult to draw general conclusions, a consensus is emerging that both species richness and community composition are important for ecosystem functioning (Hooper *et al.* 2005). However, this debate has focused on evidence from grasslands or microbial communities (Naeem *et al.* 1994; Tilman 1996; McCann 2000; Loreau *et al.* 2001, 2002; Mittelbach *et al.* 2001; Tilman *et al.* 2001; Hooper *et al.* 2005), whereas limited attention has been given to more

complex ecosystems such as forests (Pretzsch 2003; Vila *et al.* 2003). Recently, two reviews analysed biodiversity effects on ecosystem functioning but only two of 446 studies considered by Balvanera *et al.* (2006) and none of the 111 experiments reported by Cardinale *et al.* (2006) concerned forests.

Furthermore, most studies focused on the producer level and paid little or no attention to interactions among multiple trophic levels (e.g. Balvanera *et al.* 2006), which would be more representative of real ecosystems. This is critical because diversity at the consumer level can be positively or negatively related to primary production (Ives *et al.* 2005), which may lead to idiosyncratic responses of multitrophic ecosystems to biodiversity loss (Hooper *et al.* 2005). Better resource exploitation is considered to be a main driver that leads to the greater productivity of diverse plant communities (Tilman 1996; Loreau *et al.* 2001). The same principle is likely apply to herbivores, which would result in increased plant consumption (Mulder *et al.* 1999; Cardinale *et al.* 2006). Studies that assessed interactions among trophic levels also revealed contrasting relationships between food web complexity and stability, with higher diversity resulting in lower population stability (Tilman 1996; McGrady-Steed & Morin 2000; Fox & McGrady-Steed 2002) and higher total productivity (Petchey 2000; Aoki 2003; Hillebrand & Cardinale 2004; Balvanera *et al.* 2006).

There is an under-utilized body of evidence from studies on forest diversity and insect herbivory that is highly

^aEnsis is a joint venture between CSIRO and Scion/New Zealand Forest Research Institute.

relevant to this debate. The notion that boreal forests, single-species plantation forests and other low-diversity forests are particularly susceptible to pest outbreaks is widely cited in textbooks (e.g. Begon *et al.* 1996; Speight & Wylie 2001). However, previous reviews suggest that the evidence for this is equivocal (Gibson & Jones 1977; Schwerdtfeger 1981; Barthod 1994; Koricheva *et al.* 2006). Furthermore, pest problems in forest monocultures are not necessarily related to their simplified composition but could also be caused by their even-aged structure or intensive silvicultural practices (Watt 1992).

Here, we present the results from meta-analyses of 119 studies that compared insect herbivory on the same tree species growing in either pure or mixed stands. Meta-analysis is a powerful method (Gurevitch & Hedges 1993) to conduct quantitative, objective reviews of numerous independent studies. Our objectives were (i) to test the relationship between tree species diversity and forest insect herbivory, (ii) to examine whether such a relationship depends on species richness *per se* or more on tree species composition and (iii) to test whether responses of insect herbivores to tree diversity may depend on key functional traits of insect species. Our meta-analyses confirm that, overall, there is indeed an inverse relationship between forest diversity and herbivory. This applied virtually to all cases involving oligophagous species, whereas the relationship was weak when polyphagous species were considered. The qualitative composition of tree mixtures (e.g. whether mixtures of tree species composed of more or less closely related species) and the proportion of host and non-host tree species were also important. These findings are important for our understanding of forest ecology and for practical aspects of managed forests.

METHODS

Data collection

We searched for suitable studies using on-line bibliographic databases such as ISI Web of Science and CAB Abstracts and cited references in relevant publications. Our literature search yielded 119 individual studies, derived from 38 publications published from 1966 to 2006 and involved 33 tree species and 33 herbivorous insect taxa (in five additional cases the herbivore was not named; Appendix 1). We included studies for the meta-analyses if they met two conditions: (i) damage from herbivorous insect species (101 cases) or their abundance (18 cases) on a particular tree species (subsequently referred to as the 'focus tree species') was compared in pure vs. mixed stands, in the same area and time period and (ii) the mean of the response variable (herbivory), a measure of the variance and the sample size were reported in the text or available via graphic digitization. We chose the pure stand as the control and the mixed stand

as the treatment group. When results for a response variable were reported in the same paper for several insect species, data for each insect were included as an individual study. Likewise, when different tree mixtures, in terms of species composition or relative abundance, were compared to the pure stand treatment, data for each paired tree diversity comparison were included as individual studies. Results for several sampling dates were not used as replicates to calculate mean, variance and sample size values unless data were obtained in independent tree samples.

Meta-analyses

Analyses were carried out using METAWIN 2.0 software (Rosenberg *et al.* 2000). For each individual study an estimate of the magnitude of the treatment effect, Hedges' *d* effect size (eqn 1.1) was calculated as the difference between the mean herbivory magnitude of the experimental group (\bar{X}^E , mixed stand) and the control group (\bar{X}^C , pure stand) divided by the pooled standard deviation (*S*, eqn 1.2), and multiplied by a correction factor (*J*, eqn 1.3) that accounts for small sample sizes (Hedges & Olkin 1985).

$$d = j \frac{\bar{X}^E - \bar{X}^C}{S} \quad (1.1)$$

$$S = \sqrt{\frac{(N^E - 1)(S^E)^2 + (N^C - 1)(S^C)^2}{N^E + N^C - 2}} \quad (1.2)$$

$$J = 1 - \frac{3}{4(N^C + N^E - 2) - 1} \quad (1.3)$$

A negative value of *d* indicates lower herbivory in mixed stands than in pure stands. Effect sizes across all studies were combined using the random effects model (Gurevitch & Hedges 1993) to provide the grand mean effect size (d_{++}). The effect was considered as statistically significant if the bootstrap confidence interval, calculated with 999 iterations, did not bracket zero. The so-called 'file drawer problem', which refers to the potential publication bias resulting from the greater likelihood of studies with statistically significant results being published than studies that did not show any significant effects, was addressed by calculating a fail-safe sample size which represents an estimate of the number of non-significant, unpublished or missing studies that would need to be added to the analysis in order to make the overall test of an effect statistically non-significant. Rosenthal's method was used to calculate the fail-safe number for our data set and this number was then compared to Rosenthal's (1979) conservative critical value of $5n + 10$, where *n* is the sample size.

Because effect sizes were significantly heterogeneous (Q -statistics) we split the data set into subsets of classes of different categories of insects and forest stand composition. Herbivorous insects were classified according to their host specificity, whereby those that are host specific to trees within a genus or a family were categorized as 'oligophagous' and those that feed on trees from more than one family as 'polyphagous'. Forests were classified according to the focus tree class (broadleaf vs. conifer). The tree species composition of mixed stands was classified according to the presence of other palatable host species for polyphagous insect herbivores (no host vs. less palatable host vs. more palatable host). The mean effect size (d_+) and a bootstrap confidence interval were then calculated for each class and the between-classes heterogeneity was tested against a chi-square distribution to evaluate the significance of the class effect. Because multiple tests were conducted with the same data set, the resulting P -value was assessed against a Bonferroni-adjusted alpha level, a method previously used with meta-analyses (e.g. Etzel & Guerra 2002). We split the data into two subpopulations (polyphagous and oligophagous insects), and tested the effects of two factors on each subpopulation, giving four separate tests in all and a conservative adjusted alpha of 0.0125 (alpha = 0.05/4).

To examine the effects on herbivory of the relative proportion of the associated tree species, the composition of mixed stands was classified as 'fewer or as many ($\leq 50\%$)' vs. 'more ($> 50\%$) associated trees than trees of the focus species' and herbivory in both types of mixed stands was compared with herbivory in the pure stands. This was possible with 24 studies involving oligophagous insects. A Wilcoxon paired test was used to compare Hedges' effect sizes calculated for the two combinations (pure stand and mixed stand with $> 50\%$ associated trees) vs. (pure stand and mixed stand with $\leq 50\%$ associated trees). Furthermore, in eight case studies with oligophagous insect herbivores, the pure stand treatment was compared with several mixed stand treatments with an increasing proportion of associated tree species. We correlated Hedges' effect size for each comparison (pure stand vs. mixed stand) with the proportion of associated tree species in the corresponding mixed stand. The correlation coefficient r of individual studies was transformed to z correlations (Cooper & Hedges 1994) and the conditional variance of z was calculated using the METAWIN software. These estimates were included in a random model to test whether the overall mean of the correlations between the Hedges' effect size and the proportion of associated trees differed from zero. To facilitate the interpretation, the results (overall mean z) are presented as back-transformed correlation coefficients. The overall average coefficient r was analysed by calculating the 95% bootstrap confidence interval, as outlined in the text.

RESULTS

Our meta-analysis revealed that in forests of single tree species these species are significantly more affected by herbivory than the same species in more diverse forests (Table 1). The overall mean effect size equalled -0.67 and was significantly different from zero ($P = 0.0004$). According to Cohen (1988), an effect size of 0.5 – 0.8 shows a medium effect (as in this case) and an effect size > 0.8 shows a large effect. The fail-safe sample size was 3320, thus much greater than the conservative critical value of 605 ($5 \times 119 + 10$) and ≈ 28 times the number of studies included in the meta-analysis. Thus, this result is unlikely to be affected by the 'file drawer problem'.

For 115 of the 119 individual studies we were able to qualify the host specificity of the insect herbivore that was involved (Appendix 1). For oligophagous herbivores, there was a significant large and negative mean effect size, whereas for polyphagous insects, this was low and only marginally significant (Table 1). The difference between oligophagous and polyphagous insects was striking (Fig. 1) and highly significant (d.f. = 2, $Q_B = 11.5$, $P = 0.002$). (Note, for this and all following significance tests within this meta-analysis we used a Bonferroni-adjusted alpha level of 0.0125, as explained in the Methods.) Because of this dramatic difference, subsequent tests on the effects of additional variables on the diversity–herbivory relationship were conducted separately for oligophagous and polyphagous insect herbivores.

The qualitative composition of mixed stands had a significant overall effect for oligophagous insects. There was a consistently greater effect when the focus tree species and the associated tree species did not belong to the same class (i.e. broadleaved tree or conifer), with a mean effect size more than twice that for studies with mixtures that involved tree species from the same class (Table 1; d.f. = 1, $Q_B = 9.1$, $P = 0.004$). In contrast, for polyphagous insects there was no clear overall effect of the composition of mixed stands (Table 1; d.f. = 1, $Q_B = 0.7$, $P = 0.45$), although mixtures of trees from different classes had a marginally significant effect size, whereas the effect was not significant when mixtures comprised trees from the same class (Table 1).

Tests for the response in herbivory to the presence of other palatable host tree species in mixed stands also provided contrasting results for the oligophagous vs. polyphagous insect groups. Oligophagous insects showed a consistent reduction in herbivory in mixed stands regardless of the palatability of the other tree species (Table 1), although the effect size appeared to be somewhat larger when the associated trees were not palatable (d.f. = 1, $Q_B = 2.5$, $P = 0.10$). However, in studies involving polyphagous insects, the palatability of associated tree

Table 1 Meta-analyses of the influence of forest diversity on insect herbivory

Class variable*	Class	Sample size	Mean Hedges' effect size (d_+) [†]	Bootstrap confidence interval
All insects				
Overall		119	-0.67*	-0.86 to -0.49
Insect host specificity ($P = 0.002$)	Oligophagous [‡]	73	-0.95*	-1.19 to -0.75
	Polyphagous [§]	42	-0.34*	-0.62 to -0.06
Oligophagous insects ($n = 73$)				
Qualitative composition of mixed stands [¶] (associated + focus tree; $P = 0.004^{**}$)	Same classes (B + B or C + C)	36	-0.62*	-0.89 to -0.41
	Different classes (B + C or C + B)	37	-1.33*	-1.69 to -0.98
Functional composition of mixed stands [$P = 0.10$ (n.s.)]	No other palatable tree species	48	-1.11*	-1.44 to -0.82
	Other less palatable tree species	23	-0.70*	-0.98 to -0.45
Polyphagous insects ($n = 42$)				
Qualitative composition of mixed stands [associated + focus tree; $P = 0.45$ (n.s.)]	Same classes (B + B or C + C)	25	-0.25	-0.62 to +0.10
	Different classes (B + C or C + B)	17	-0.47*	-0.86 to -0.04
Functional composition of mixed stands ($P = 0.001^{**}$)	No other palatable tree species	5	-1.61*	-2.62 to -0.61
	Other less palatable tree species	16	-0.66*	-0.91 to -0.46
	Other more palatable tree species	21	+0.16	-0.27 to +0.63

Herbivory is compared between pure stands of a focus tree species and mixed stands composed of the focus species and associated tree species. Negative Hedges' effect size indicates that the rate of herbivory on focus trees decreased in mixed stands.

*Class effect (P -value).

[†]The asterisk denotes a significant effect based on a bootstrap test.

[‡]Insects with a host range that includes at most one family of trees.

[§]Insects with a host range that includes trees from more than one family of trees.

[¶]Tree class: B = broadleaf, C = conifer.

**Significant if $P < 0.0125$ (Bonferroni correction, $\alpha = 0.05/4 = 0.0125$).

species was a significant factor (Table 1; d.f. = 2, $Q_B = 18.5$, $P = 0.001$). For the individual studies where no other palatable trees ($n = 5$) or only less susceptible tree species ($n = 16$) were present in mixed stands, the size of the overall effect was large or medium and negative, indicating that these mixed forests had a lower risk of damage from polyphagous pests (Table 1). On the other hand, for the other 21 studies where more palatable trees were present in mixed stands the overall effect size was positive, although non-significant (Table 1). This indicates that there is a risk of greater damage from polyphagous pests when a more palatable host tree is associated with the focus tree species (Table 1).

We were also able to examine the effects of different proportions of associated tree species in the mixed stands. We compiled a subset of studies that met this requirement, all concerning oligophagous insects, and compared the mean effect sizes of 24 pairs of individual studies focusing on the same insect–tree interaction where the proportion of associated tree species in the mixture was either greater than, equal to or smaller than the proportion of the focus tree species (Fig. 2). In studies where the proportion of associated tree species was $> 50\%$, the mean effect size was on average approximately twice that (-2.7 vs. -1.2) of studies with a smaller proportion (Wilcoxon paired test,

$n = 24$, $Z = 4.14$, $P < 0.0001$). This indicated that an increased proportion of associated trees in mixed stands led to a decrease in herbivory sustained by the focus trees (Fig. 2). This pattern was even more obvious in eight case studies with oligophagous insect herbivores where the pure stand treatment was compared with several mixed stand treatments with an increasing proportion of associated tree species. Overall, there was a consistently increasing effect size when the proportion of associated tree species increased (Fig. 3). The mean correlation coefficient r between the decrease in herbivory in mixed stands compared to pure stands and the rate of associated tree species was 0.95 and significantly different from 0 (95% bootstrap confidence interval ± 0.02 ; back-transformed from $\hat{\tau}$ -transformed correlation for random model).

DISCUSSION

Based on the evidence from 119 comparative studies of 47 different insect–tree interactions, our quantitative review showed that overall tree species growing in mixtures suffer significantly less herbivory than those in pure stands. This diversity effect on herbivory appears to be generally applicable at this forest stand scale as the studies used in our meta-analysis include a wide range of insect taxa and




Figure 1 Hedges' d effect size and variance of individual studies on the response of forest insect herbivory to pure vs. mixed tree stand composition, for (a) oligophagous herbivores and (b) polyphagous herbivores. Negative effect sizes indicate that higher tree species diversity (mixed stand treatment) results in lower insect herbivore abundance or damage. More information on each study is given in Appendix 1.




Figure 2 Twenty-four paired comparisons of the effect of increased tree species diversity on herbivory by oligophagous insects in mixed stands, in studies where two relative abundance classes of associated trees were compared with pure stands. Note: Light grey bars show Hedges' d effect size of insect herbivory in pure stands vs. mixed stands where associated trees were more frequent ($> 50\%$) than the focus tree species. Dark grey bars show Hedges' d effect size of insect herbivory in pure stand vs. mixed stand where associated trees were less frequent ($\leq 50\%$) than the focus tree species.




Figure 3 Eight response patterns of Hedges' d to the relative abundance of associated tree species in mixed stands compared with pure stands. (1) Nichols *et al.* 1999; (2, 4, 5, 6) Fauss & Pierce 1969; (3) Jactel *et al.* 2006; (7) Su *et al.* 1996; (8) Katovich 1992.

feeding guilds that affect trees from numerous plant families and orders in boreal, temperate and tropical biomes (see also Jactel *et al.* 2005). Overall, our study provides new and convincing support for the theory that species diversity of producer assemblages may reduce the magnitude of consumer effects on producers. Three similar meta-analyses that examined diversity effects in crop plant and algal communities (Tonhasca & Byrne 1994; Hillebrand & Cardinale 2004; Balvanera *et al.* 2006) also showed that more diverse communities were less affected by herbivore consumption. Our analysis gives support to the extension of this global pattern to long-lived producers such as tree species. However, the magnitude of this effect varied greatly in relation to an additional important factor that has been overlooked in previous studies, the host specificity of the primary consumer. The contrast between the responses of oligophagous and polyphagous herbivores to increased tree diversity was striking and merits further exploration.

What distinguished the subpopulation of studies involving *polyphagous* insects in our meta-analysis was the significantly different effect size in those studies where other, more palatable species were present in mixed stands. In at least six of these studies (Brown *et al.* 1988; Gottschalk & Twery 1989; White & Whitham 2000) this result can most likely be attributed to 'associational susceptibility', an outcome of the occurrence of several host tree species in situations where polyphagous herbivores may be able to use a more palatable host to build up their populations, exploit those resources and then 'spill over' to the associated hosts (Brown & Ewel 1987; White & Whitham 2000). For example, gypsy moth (*Lymantria dispar*) can feed on conifers once its preferred hosts, oaks and other broadleaved species have been defoliated. As a result of this polyphagy, white pine growing in mixed stands with oaks is more likely to be attacked by *L. dispar* than white pine in pure stands (Brown *et al.* 1988). We were able to confirm this process as a general pattern by separating studies with polyphagous insects according to the presence or absence of other host tree species in the mixture. Our results clearly showed that such a detrimental effect of tree diversity is not uncommon but only when a more palatable host is present in the tree species assemblage. Similar effects may apply to generalist mammalian grazers which appear to prefer mixed stands in boreal forests (Koricheva *et al.* 2006).

It is noteworthy that our analysis considers the *overall* effects of many different studies of which each examined a particular herbivore species on a particular focus tree species growing either in a pure or a mixed stand. It is conceivable that the observed *overall* reduction in herbivory by a particular oligophagous herbivore on a particular tree species could be outweighed by a corresponding *overall* increase in total herbivory by polyphagous species on all tree

species present in mixed stands. To our knowledge, there has been no published study that specifically addressed this question in forest ecosystems; however, two observations can be offered that oppose this view: (i) collectively, overall herbivory by polyphagous species was not greater in mixed stands than in pure stands and (ii) oligophagous, i.e. specialist herbivores, are likely to exert a higher herbivory pressure than generalist, i.e. polyphagous species. This has been demonstrated particularly in tropical forests (Barone 1998, 2000).

In the subset of studies involving *oligophagous* insects, we observed a significant effect of the composition of mixed stands. Increased host availability is often cited as the main reason for the higher rates of herbivory in monocultures (Root 1973; Risch 1981; Russell 1989; Jones 2001). This process is well supported by the results of our meta-analyses which showed that trees in mixed stands consistently experienced less herbivory when the proportion of non-host trees increased (e.g. Fauss & Pierce 1969; Katovich 1992; Su *et al.* 1996; Nichols *et al.* 1999; Jactel *et al.* 2006). In more diverse plant communities, each particular plant species is relatively less abundant and its distribution is likely to be more patchy, making it less available to specialist herbivores (Kareiva 1983; Yamamura 2002). Apart from this 'spatial concentration' effect, an association with non-host species can also provide physical or chemical barriers to host location by foraging or dispersing herbivores. Numerous tree-feeding insects are wind-dispersed, and physical barriers are known to affect such insects in particular. For example, the larvae of *C. fumiferana* are passively dispersed by ballooning in wind currents, and their survival is higher in pure stands of balsam fir, their main host tree, than in mixed stands due to the reduced risk of landing on a non-host (Kemp & Simmons 1979). Hence, balsam fir in pure stands tends to be more defoliated than in mixed stands (Cappucino *et al.* 1998). Chemical barriers to host location are known to occur when stimuli from non-host trees disrupt olfactory host tree recognition. Several non-host volatiles have been identified from angiosperm trees, which showed repellent effects in conifer-feeding bark beetles (Byers *et al.* 1998; Schlyter *et al.* 2000; Huber & Borden 2001; Jactel *et al.* 2001; Zhang *et al.* 2001). Our analysis supports the relevance of this 'chemical barrier' effect on herbivory because associations of trees from two different classes (broadleaved trees and conifers) reduced herbivory in mixed stands more than associations of trees from the same class (e.g. broadleaved plus broadleaved or conifers plus conifers), which would have volatiles that are more similar.

Another likely mechanism is generally known as the 'natural enemy' hypothesis (Root 1973; Wilby & Thomas 2002; Cardinale *et al.* 2003; Jakel & Roth 2005; Riihimäki *et al.* 2005) but only a few of the studies included in our meta-analysis tested this directly. The density of

accompanying understorey vegetation appeared not to influence the rate of parasitism of the Nantucket pine tip (Miller & Stephen 1983). In contrast, both Bae *et al.* (1997) and Quayle *et al.* (2003) showed that the parasitism rate of herbivorous insects was significantly higher in mixed stands than in pure stands. *Trichogramma minutum* Riley, the main parasitoid in a study on spruce budworm (*Choristoneura fumiferana*), is a generalist species that appeared to benefit from eggs of other insect species feeding on non-host trees (Quayle *et al.* 2003). Generalist parasitoids and predators are likely to encounter a higher availability of alternative prey or hosts in mixed forests because diverse plant communities usually provide habitat for more herbivorous species (Lawton & Strong 1981; Siemann *et al.* 1998). Richer plant communities can also provide a better supply of complementary food such as pollen, nectar and honeydew that improve the fitness of specialized parasitoids (Simmons *et al.* 1975; Russell 1989; Cappucino *et al.* 1999). Finally, because of their higher structural complexity, mixed forests may provide natural enemies with more opportunities to shelter from adverse conditions, or provide nesting sites as in the case of birds that prey on spruce budworm (Zach & Falls 1975; Dickson 1979).

Thus, our results are consistent with two of the main points in relation to the effects of biodiversity on ecosystem functioning on which ecologists recently agreed (Hooper *et al.* 2005): (i) species composition can be more important than species richness *per se* and (ii) the strength or the sign of the relationship can vary with the functional traits of the species. Our results confirm this because (i) the beneficial effects of tree diversity on herbivory increase with the relative proportion of associated trees, (ii) the association of phylogenetically less similar tree species, e.g. angiosperms associated with gymnosperms, is more effective in preventing herbivory in mixed stands and (iii) the association of more palatable host trees is likely to increase damage from polyphagous herbivores in the mixture. These findings also highlight the fact that insufficient consideration of such factors may lead to erroneous conclusions about the existence of such effects of biodiversity, and whether or not they are widely applicable. This also has important implications for the sustainable management of forests, particularly planted forests which are predominantly managed as monocultures and are expected to soon supply the majority of the world's demand for wood and fibre products (FAO 2001). Enhancing biodiversity by enriching forests with additional tree species that are less palatable for pest insects may increase food web stability while at the same time potentially offering conservation benefits. However, more comprehensive studies are needed to determine whether there is a concomitant effect of forest diversity on herbivory, productivity and conservation.

ACKNOWLEDGEMENTS

We would like to thank Mark Kimberley for assistance with statistical analyses and John Bain, Margaret Dick, Judy Griffiths and three anonymous referees for their comments on the manuscript. This paper was partly funded by a Co-operative Project under the framework of the France and New Zealand Cultural Agreement, the New Zealand Foundation for Research Science and Technology (C04X0302), and the French GIP Ecofor.

REFERENCES

- Aoki, I. (2003). Diversity-productivity-stability relationship in freshwater ecosystems: whole-systemic view of all trophic levels. *Ecol. Res.*, 18, 397–404.
- Bae, W.I., Shin, S.C. & Kim, Z.S. (1997). Difference in occurrence of pine needle gall midge and sucking insects in pure-pine and mixed-pine stands. *FRI J. For. Sci.*, 56, 37–43.
- Balmford, A. & Bond, W. (2005). Trends in the state of nature and their implications for human well-being. *Ecol. Lett.*, 8, 1218–1234.
- Balvanera, P., Pfisterer, A.B., Buchmann, N., He, J.S., Nakashizuka, T., Raffaelli, D. *et al.* (2006). Quantifying the evidence for biodiversity effects on ecosystem functioning and services. *Ecol. Lett.*, 9, 1146–1156.
- Barone, J.A. (1998). Host-specificity of folivorous insects in a moist tropical forest. *J. Anim. Ecol.*, 67, 400–409.
- Barone, J.A. (2000). Comparison of herbivores and herbivory in the canopy and understory for two tropical tree species. *Biotropica*, 32, 307–317.
- Barthod, C. (1994). Sylviculture et risques sanitaires dans les forêts tempérées – 1ère partie. *Rev. For. Fr.*, 46, 609–628.
- Batzler, H.O., Benzie, J.W. & Popp, M.P. (1987). Spruce budworm damage in aspen/balsam fir stands affected by cutting methods. *North. J. Appl. For.*, 4, 73–75.
- Begon, M., Harper, J.L. & Townsend, C.R. (1996). *Ecology: Individuals, Populations and Communities*. Blackwell Science Ltd, Oxford, UK, p. 840.
- Bergeron, Y., Leduc, A., Morin, H. & Joyal, C. (1995). Balsam fir mortality following the last spruce budworm outbreak in northwestern Quebec. *Can. J. For. Res.*, 25, 1375–1384.
- Berisford, C.W. & Kulman, H.M. (1967). Infestation rate and damage by the Nantucket pine tip moth in six loblolly pine stand categories. *For. Sci.*, 13, 429–438.
- Bigger, M. (1985). The effect of attack by *Amblyopelta cocophaga* (Hemiptera: Coreidae) on growth of *Eucalyptus deglupta* in the Solomon Islands. *Bull. Ent. Res.*, 75, 595–608.
- Brook, B.W., Sodhi, N.S. & Ng, P.K.L. (2003). Catastrophic extinctions follow deforestation in Singapore. *Nature*, 424, 420–423.
- Brown, B.J. & Ewel, J.J. (1987). Herbivory in complex and simple tropical successional ecosystems. *Ecology*, 68, 108–116.
- Brown, J.H., Cruickshank, V.B., Gould, W.P. & Husband, T.P. (1988). Impact of gypsy moth defoliation in stands containing white pine. *North. J. Appl. For.*, 5, 108–111.
- Byers, J.A., Zhang, Q.H., Schlyter, F. & Birgersson, G. (1998). Volatiles from non-host birch trees inhibit pheromone response in spruce bark beetles. *Naturwissenschaften*, 85, 557–561.

- Cappuccino, N., Lavertu, D., Bergeron, Y. & Regniere, J. (1998). Spruce budworm impact, abundance and parasitism rate in a patchy landscape. *Oecologia*, 114, 236–242.
- Cappuccino, N., Houle, M.J. & Stein, J. (1999). The influence of understory nectar resources on parasitism of the spruce budworm *Choristoneura fumiferana* in the field. *Agric. For. Entomol.*, 1, 33–36.
- Cardinale, B.J., Harvey, C.T., Gross, K. & Ives, A.R. (2003). Biodiversity and biocontrol: emergent impact of a multi-enemy assemblage on pest suppression and crop yield in an agroecosystem. *Ecol. Lett.*, 6, 857–865.
- Cardinale, B.J., Srivastava, D.S., Duffy, J.E., Wright, J.P., Downing, A.L., Sankaran, M. *et al.* (2006). Effects of biodiversity on the functioning of trophic groups and ecosystems. *Nature*, 443, 989–992.
- Cohen, J. (1988). *Statistical Power Analysis for the Behavioral Sciences*, 2nd edn. Lawrence Earlbaum Associates, Hillsdale, NJ, USA.
- Cooper, H. & Hedges, L. (1994). *The Handbook of Research Synthesis*. Russel Sage Foundation, New York, NY, USA.
- Dickson, J.G. (1979). Seasonal population of insectivorous birds in mature bottomland hardwood forest in south Louisiana. In: *The Role of Insectivorous Birds in Forest Ecosystems* (eds Dickson, J.G., Connor, R.N., Fleet, R.R., Jackson, J.A. & Kroll, J.C.). Academic Press, New York, pp. 261–269.
- Etzel, C.J. & Guerra, R. (2002). Meta-analysis of genetic-linkage analysis of quantitative-trait loci. *Am. J. Hum. Genet.*, 71, 56–65.
- FAO (2001). *Future Production from Forest Plantations*. Report based on the work of A. del Lungo. Forest Plantation Thematic Papers, Working Paper 14. Forest Resources Development Service, Forest Resources Division. FAO, Rome.
- Fauss, D.L. & Pierce, W.R. (1969). Stand conditions and spruce budworm damage in a western Montana forest. *J. For.*, 67, 322–325.
- Folgarait, P.J., Marquis, R.J., Ingvarsson, P., Bracker, H.E. & Arguedas, M. (1995). Pattern of attack by insect herbivores and fungus on saplings in a tropical tree plantation. *Environ. Entomol.*, 24, 1487–1494.
- Fox, J.W. & McGrady-Steed, J. (2002). Stability and complexity in microcosm communities. *J. Anim. Ecol.*, 71, 749–756.
- Gantner, M. (2000). Aphidofauna of hazel bushes (*Corylus L.*) on a protected plantation, an unprotected plantation and in a forest. *Ann. Univ. Mariae Curie Skłodowska, Sect. EE Hort.*, 8, 155–167.
- Géri, C. (1980). Application des méthodes d'études démécologiques aux insectes défoliateurs forestiers: cas de *Diprion pini* et dynamique des populations de la processionnaire du pin en Corse. Thesis, Université de Paris-Sud.
- Gibson, I.A.S. & Jones, T.M. (1977). Monoculture as the origin of major pests and diseases. In: *Origins of Pest, Parasite, Disease and Weed Problems* (eds Cherrett, J.M. & Sagar, G.R.). 8th Symposium of the British Ecological Society, Bangor, 12–14 April 1976. Blackwell Scientific Publication, Oxford, pp. 139–161.
- Gottschalk, K.W. & Twery, M.J. (1989). Gypsy moth in pine-hardwood mixtures. In: *Pine-hardwood Mixtures: a Symposium on Management and Ecology of the Type*. (ed. Waldrop, T.A.), USDA Forest Service General Technical Report SE 58, Southeastern Forest Experiment Station, Asheville, NC 28802, pp. 50–58.
- Greaves, R. (1966). Insect defoliation of *Eucalyptus* regrowth in the Florentine Valley, Tasmania. *Appita*, 19, 119–126.
- Gurevitch, J. & Hedges, L.V. (1993). Meta-analysis: Combining the results of independent experiments. In: *Design and Analysis of Ecological Experiments* (eds Scheiner, S.M. & Gurevitch, J.), Chapman & Hall, New York, pp. 378–425.
- Hassan, R., Scholes, R. & Ash, N. (2005). *Ecosystems and Human Well-being: Current State and Trends*, Vol. 1. Island Press, Washington.
- Hedges, L.V. & Olkin, I. (1985). *Statistical Methods for Meta-Analysis*. Academic Press, New York.
- Hillebrand, H. & Cardinale, B.J. (2004). Consumer effects decline with prey diversity. *Ecol. Lett.*, 7, 192–201.
- Hooper, D.U., Chapin, F.S., Ewel, J.J., Hector, A., Inchausti, P., Lavorel, S. *et al.* (2005). Effects of biodiversity on ecosystem functioning: a consensus of current knowledge. *Ecol. Monogr.*, 75, 3–35.
- Huber, D.P.W. & Borden, J.H. (2001). Angiosperm bark volatiles disrupt response of Douglas-fir beetle, *Dendroctonus pseudotsugae*, to attractant-baited traps. *J. Chem. Ecol.*, 27, 217–233.
- Ives, A.R., Cardinale, B.J. & Snyder W.E. (2005). A synthesis of subdisciplines: predator-prey interactions, and biodiversity and ecosystem functioning. *Ecol. Lett.*, 8, 102–116.
- Jactel, H., Van Halder, I., Menassieu, P., Zhang, Q.H. & Schlyter, F. (2001). Non-host volatiles disrupt the response of the steno-grapher bark beetle, *Ips sexdentatus* (Coleoptera: Scolytidae), to pheromone baited traps and maritime pine logs. *I.P.M. Rev.*, 6, 197–207.
- Jactel, H., Brockerhoff, E. & Duelli, P. (2005). A test of the biodiversity-stability theory: meta-analysis of tree species diversity effects on insect pest infestations, and re-examination of responsible factors. In: *Forest Diversity and Function – Temperate and Boreal Systems* (eds Scherer-Lorenzen, M., Körner, C. & Schulze, E.-D.). Springer Verlag, Berlin Heidelberg, Ecological studies, Vol. 176, pp. 235–262.
- Jactel, H., Menassieu, P., Vétillard, F., Gaulier, A., Samalens, J.C. & Brockerhoff, E.G. (2006). Tree species diversity reduces the invasibility of maritime pine stands by the bark scale, *Matsucoccus feytaudi* (Homoptera: Margarodidae). *Can. J. For. Res.*, 36, 314–323.
- Jakel, A. & Roth, M. (2005). Conversion of single layered Scots pine monocultures into close-to-nature mixed hardwood forests: effects of parasitoid wasps as pest antagonists. *Eur. J. For. Res.*, 123, 203–212.
- Jones, R.E. (2001). Mechanisms for locating resources in space and time: impacts on the abundance of insect herbivores. *Austral. Ecol.*, 26, 518–524.
- Kareiva, P. (1983). Influence of vegetation texture on herbivore populations: resource concentration and herbivore movement. In: *Variable Plants and Herbivores in Natural and Managed Systems* (eds Denno, R.F. & McClure, M.S.), Academic Press, New York, pp. 259–289.
- Katovich, S.A. (1992). White pine weevil response to oak overstorey girdling – results from a 16-year-old study. *North. J. Appl. For.*, 9, 51–54.
- Keenan, R., Lamb, D. & Sexton, G. (1995). Experience with mixed species rainforest plantations in North Queensland. *Commonw. For. Rev.*, 74, 315–321.
- Kemp, W.P. & Simmons, G.A. (1979). Influence of stand factors on survival of early instar spruce budworm. *Environ. Entomol.*, 8, 993–996.
- Koricheva, J., Vehviläinen, H., Riihimäki, J., Ruohomäki, K., Kaitaniemi, P. & Ranta, H. (2006). Diversification of tree stands as a means to manage pests and diseases in boreal forests: myth or reality? *Can. J. For. Res.*, 36, 324–336.

- Lawton, J.H. & Strong, D.R. (1981). Community patterns and competition in folivorous insects. *Am. Nat.*, 118, 317–338.
- Lindenmayer, D.B. & Franklin, J.F. (2002). *Conserving Forest Biodiversity*. Island Press, Washington, DC.
- Loreau, M., Naeem, S., Inchausti, P., Bengtsson, J., Grime, J.P., Hector, A. *et al.* (2001). Biodiversity and ecosystem functioning: current knowledge and future challenges. *Science*, 294, 804–808.
- Loreau, M., Naeem, S. & Inchausti, P. (2002). *Biodiversity and Ecosystem Functioning. Synthesis and Perspectives*. Oxford University Press, Oxford, UK.
- McCann, K.S. (2000). The diversity-stability debate. *Nature*, 405, 228–233.
- McGrady-Steed, J. & Morin, P.J. (2000). Biodiversity, density compensation, and the dynamics of populations and functional groups. *Ecology*, 81, 361–373.
- McLean, D.A. (1980). Vulnerability of fir-spruce stands during uncontrolled spruce budworm outbreaks: a review and discussion. *For. Chron.*, 56, 213–221.
- McLean, J.A. (1989). Effect of red alder overstorey on the occurrence of *Pissodes strobi* during the establishment of a Sitka spruce plot. In: *Insects Affecting Reforestation: Biology and Damage* (eds Alfaro, R.I. & Glover, S.G.), Proceedings of a meeting of the IUFRO working group on insects affecting reforestation, Vancouver, 3–9 July 1988, Forestry Canada, Victoria. pp. 167–176.
- Miller, F.D. & Stephen, F.M. (1983). Effects of competing vegetation on Nantucket pine tip moth (Lepidoptera: Tortricidae) populations in loblolly pine plantations in Arkansas. *Environ. Entomol.*, 12, 101–105.
- Mittelbach, G.G., Steiner, C.F., Scheiner, S.M., Gross, K.L., Reynolds, H.L., Waide, R.B. *et al.* (2001). What is the observed relationship between species richness and productivity? *Ecology*, 82, 2381–2396.
- Montagnini, F., Gonzalez, E., Porras, C. & Rheingans, R. (1995). Mixed and pure forest plantations in the humid neotropics: a comparison of early growth, pest damage and establishment costs. *Commonw. For. Rev.*, 74, 306–314.
- Moore, R., Warrington, S. & Whittaker, J.B. (1991). Herbivory by insects on oak trees in pure stands compared with paired mixtures. *J. Appl. Ecol.*, 28, 290–304.
- Mulder, C.P.H., Koricheva, J., Huss-Danell, K., Hogberg, P. & Joshi, J. (1999). Insects affect relationships between plant species richness and ecosystem processes. *Ecol. Lett.*, 2, 237–246.
- Naeem, S., Thompson, L.J., Lawler, S.P., Lawton, J.H. & Woodfin, R.M. (1994). Declining biodiversity can alter the performance of ecosystems. *Nature*, 368, 734–736.
- Nichols, D.J., Ofori, D.A., Wagner, M.R., Bosu, P. & Cobbinah, J.R. (1999). Survival, growth and gall formation by *Phytolyma lata* on *Milicia excelsa* established in mixed-species tropical plantations in Ghana. *Agric. For. Entomol.*, 1, 137–141.
- Nordlander, G., Bylund, H., Örlander, G. & Wallertz, K. (2003). Pine weevil population density and damage to coniferous seedlings in a regeneration area with and without shelterwood. *Scand. J. For. Res.*, 18, 438–448.
- Nowak, J.T., Harrington, T.B. & Berisford, C.W. (2003). Nantucket pine tip moth development and population dynamics: influence of nitrogen fertilization and vegetation control. *For. Sci.*, 49, 731–737.
- Örlander, G., Nordlander, G. & Wallertz, K. (2001). Extra food supply decreases damage by the pine weevil *Hyllobius abietis*. *Scand. J. For. Res.*, 16, 450–454.
- Petchey, O.L. (2000). Prey diversity, prey composition, and predator population dynamics in experimental microcosms. *J. Anim. Ecol.*, 69, 874–882.
- Petersson, M. & Örlander, G. (2003). Effectiveness of combinations of shelterwood, scarification, and feeding barriers to reduce pine weevil damage. *Can. J. For. Res.*, 33, 64–73.
- Pretzsch, H. (2003). Diversity and productivity of forests. *Allgem. For. Jagdzeit.*, 174, 88–98.
- Quayle, D., Régnière, J., Cappucino, N. & Dupont, A. (2003). Forest composition, host population density, and parasitism of spruce budworm *Choristoneura fumiferana* eggs by *Trichogramma minutum*. *Entomol. Exp. Appl.*, 107, 215–227.
- Riihimäki, J., Kaitaniemi, P., Koricheva, J. & Vehviläinen, H. (2005). Testing the enemies hypothesis in forest stands: the important role of tree species composition. *Oecologia*, 142, 90–97.
- Risch, S.J. (1981). Insect herbivore abundance in tropical monocultures and polycultures: an experimental test of two hypotheses. *Ecology*, 62, 1325–1340.
- Root, R.B. (1973). Organisation of a plant-arthropod association in simple and diverse habitats: the fauna of collards (*Brassica oleracea*). *Ecol. Monogr.*, 43, 94–125.
- Rosenberg, M.S., Adams, D.C. & Gurevitch, J. (2000). *MetaWin: Statistical Software for Meta-analysis. Version 2.0*. Sinauer Associates, Sunderland, Massachusetts.
- Rosenthal, R. (1979). The “file drawer problem” and tolerance for null results. *Psychol. Rep.*, 12, 491–511.
- Russell, E.P. (1989). Enemies hypothesis: a review of the effect of vegetational diversity on predatory insects and parasitoids. *Environ. Entomol.*, 18, 590–599.
- Saunders, D.A., Hobbs, R.J. & Margules, C.R. (1991). Biological consequences of ecosystem fragmentation: a review. *Conserv. Biol.*, 5, 18–32.
- Schlyter, F., Zhang, Q.H., Andersson, P., Byers, J.A., Wadhams, L., Löfqvist, J. *et al.* (2000). Electrophysiological and behavioral responses of pine shoot beetles, *Tomicus piniperda* and *T. minor* (Coleoptera: Scolytidae), to non-host leaf and bark volatiles. *Can. Entomol.*, 132, 965–981.
- Schwalter, T.D. & Turchin, P. (1993). Southern pine beetle infestation development: interaction between pine and hardwood basal areas. *For. Sci.*, 39, 201–210.
- Schwerdtfeger, F. (1981) *Waldkrankheiten*. Paul Parey Verlag, Hamburg and Berlin.
- Siemann, E., Tilman, D., Haarstad, J. & Ritchie, M. (1998). Experimental tests of the dependence of arthropod diversity on plant diversity. *Am. Nat.*, 152, 738–750.
- Simmons, G.A., Leonard, D.E. & Chen, C.W. (1975). Influence of tree species density and composition on parasitism of the spruce budworm, *Choristoneura fumiferana*. *Environ. Entomol.*, 4, 832–836.
- Smith, I.W., Marks, G.C., Featherston, G.R. & Geary, P.W. (1989). Effects of inter-planted wattles on the establishment of *Eucalyptus* planted on forest sites affected by *Phytophthora cinnamomi*. *Aust. For.*, 52, 74–81.
- Soria, F.J., Villagran, R., DelTio, R. & Ocete, M.E. (1995). Incidencia de *Curculio elephas* (Coleoptera: Curculionidae) en alcornoques y encinares del parque natural Sierra Norte de Sevilla. *Bol. Sanid. Veg., Plagas.*, 21, 195–201.
- Speight, M.R. & Wylie, F.R. (2001). *Insect Pests in Tropical Forestry*. CABI Publishing, Oxon, UK.

- Stadniskii, G.V. (1978). Effect of the living ground vegetation on damage to Scots pine plantations by *Hyllobius abietis*. *Lesnoi Z.*, 6, 26–29.
- Su, Q., McLean, D.A. & Needham, T.D. (1996). The influence of hardwood content on balsam fir defoliation by spruce budworm. *Can. J. For. Res.*, 26, 1620–1628.
- Taylor, S.P., Alfaro, R.I., DeLong, C. & Rankin, L. (1996). The effect of overstory shading on white pine weevil damage to white spruce and its effects on spruce growth rates. *Can. J. For. Res.*, 26, 306–312.
- Thomas, C.D., Cameron, A., Green, R.E., Bakkenes, M., Beaumont, L.J., Collingham, Y.C. *et al.* (2004). Extinction risk from climate change. *Nature*, 427, 145–148.
- Tilman, D. (1996). Biodiversity: population versus ecosystem stability. *Ecology*, 77, 350–363.
- Tilman, D., Reich, P.B., Knops, J., Wedin, D., Mielke, T. & Lehmann, C. (2001). Diversity and productivity in a long-term grassland experiment. *Science*, 294, 843–845.
- Tonhasca, A. & Byrne, D.N. (1994). The effect of crop diversification on herbivorous insects: a meta-analysis approach. *Ecol. Entomol.*, 19, 239–244.
- United Nations (1992). *Convention on Biological Diversity*, 17 July 1992. United Nations, New York, Reprinted in: *Environ. Policy Law*, 22, 251–258.
- Vehviläinen, H., Koricheva, J., Ruohomäki, K., Johansson, T. & Valkonen, S. (2006). Effects of tree stand species composition on insect herbivory of silver birch in boreal forests. *Basic Appl. Ecol.*, 7, 1–11.
- Vila, M., Vayreda, J., Gracia, C. & Ibanez, J.J. (2003). Does tree diversity increase wood production in pine forests? *Oecologia*, 135, 299–303.
- Von Sydow, F. & Örlander, G. (1994). The influence of shelterwood density on *Hyllobius abietis* (L.) occurrence and feeding on planted conifers. *Scand. J. For. Res.*, 9, 367–375.
- Watt, A.D. (1992). Insect pest population dynamics: effects of tree species diversity. In: *The Ecology of Mixed-Species Stands of Trees* (eds Cannell, M.G.R., Malcolm, D.C. & Robertson, P.A.), Blackwell Scientific Publications, Oxford, pp. 267–275.
- Wazihullah, A.K.M., Islam, S.S., Rahman, F. & Das, S. (1996). Reduced attack of keora (*Sonneratia apetala*) by stem borer in mixed-species plantations in coastal Bangladesh. *J. Trop. For. Sci.*, 8, 476–480.
- White, J.A. & Whitham, T.G. (2000). Associational susceptibility of cottonwood to a box elder herbivore. *Ecology*, 81, 1795–1803.
- Wilby, A. & Thomas, M.B. (2002). Natural enemy diversity and pest control: patterns of pest emergence with agricultural intensification. *Ecol. Lett.*, 5, 353–360.
- Yamamura, K. (2002). Biodiversity and stability of herbivore populations: influences of the spatial sparseness of food plants. *Popul. Ecol.*, 44, 33–40.
- Zach, R. & Falls, J.B. (1975). Response of the ovenbird (Aves: Parulidae) to an outbreak of the spruce budworm. *Can. J. Zool.*, 53, 1669–1672.
- Zanuncio, J.C., Mezzomo, J.A., Guedes, R.N.C. & Oliveira, A.C. (1998). Influence of strips of native vegetation on Lepidoptera associated with *Eucalyptus cloeziana* in Brazil. *For. Ecol. Manag.*, 108, 85–90.
- Zhang, Q.H., Liu, G.T., Schlyter, F., Birgersson, G., Anderson, P. & Valeur, P. (2001). Olfactory responses of *Ips duplicatus* from Inner Mongolia, China to nonhost leaf and bark volatiles. *J. Chem. Ecol.*, 27, 995–1009.

Editor, Michael Hochberg

Manuscript received 25 April 2007

First decision made 20 May 2007

Manuscript accepted 29 May 2007

Appendix 1 Summary of data included in the meta-analyses

Individual study	Authors	Tree species	Insect species	Insect host specificity	RAH*	Hedges' d	N^E	X^E	N^C	X^C	S
1	Su <i>et al.</i> (1996)	<i>Abies balsamea</i>	<i>Choristoneura fumiferana</i>	O	More	-5.38	5	0.14	5	0.72	0.10
2	Su <i>et al.</i> (1996)	<i>Abies balsamea</i>	<i>Choristoneura fumiferana</i>	O	More	-4.46	5	0.25	5	0.72	0.10
3	Su <i>et al.</i> (1996)	<i>Abies balsamea</i>	<i>Choristoneura fumiferana</i>	O	Equal	-2.89	5	0.32	5	0.72	0.12
4	Su <i>et al.</i> (1996)	<i>Abies balsamea</i>	<i>Choristoneura fumiferana</i>	O	Less	-0.46	5	0.65	5	0.72	0.13
5	Bergeron <i>et al.</i> (1995)	<i>Abies balsamea</i>	<i>Choristoneura fumiferana</i>	O	More	-1.21	3	57.57	3	74.00	10.87
6	Bergeron <i>et al.</i> (1995)	<i>Abies balsamea</i>	<i>Choristoneura fumiferana</i>	O	Less	-0.84	3	63.20	3	74.00	10.25
7	McLean (1980)	<i>Abies balsamea</i>	<i>Choristoneura fumiferana</i>	O		-0.39	3	0.27	3	0.37	0.21
8	Batzler <i>et al.</i> (1987)	<i>Abies balsamea</i>	<i>Choristoneura fumiferana</i>	O	More	-3.22	15	20.00	15	93.00	22.08
9	Batzler <i>et al.</i> (1987)	<i>Abies balsamea</i>	<i>Choristoneura fumiferana</i>	O	Less	-2.61	10	32.00	15	93.00	22.58
10	Quayle <i>et al.</i> (2003)	<i>Abies balsamea</i>	<i>Choristoneura fumiferana</i>	O		-0.67	4	0.25	4	0.33	0.11
11	Quayle <i>et al.</i> (2003)	<i>Abies balsamea</i>	<i>Choristoneura fumiferana</i>	O		-1.75	4	0.16	4	0.33	0.09
12	Montagnini <i>et al.</i> (1995)	<i>Abies balsamea</i>	<i>Choristoneura fumiferana</i>	O		0.13	4	0.71	4	0.70	0.04
13	Vehviläinen <i>et al.</i> (2006)	<i>Abies balsamea</i>	<i>Attia cephalotes</i>	P		-0.96	8	0.82	5	1.00	0.17
14	Vehviläinen <i>et al.</i> (2006)	<i>Betula pendula</i>	Insect defoliators	O	Equal	-1.80	5	0.68	5	1.00	0.16
15	Vehviläinen <i>et al.</i> (2006)	<i>Betula pendula</i>	<i>Acallitus rudis</i>	O	Equal	-1.72	8	1.96	5	4.29	1.26
16	Vehviläinen <i>et al.</i> (2006)	<i>Betula pendula</i>	<i>Acallitus rudis</i>	O	More	-1.19	5	2.50	5	4.29	1.36
17	Vehviläinen <i>et al.</i> (2006)	<i>Betula pendula</i>	Leaf rollers	O	Equal	-1.35	8	0.36	5	0.54	0.12
18	Vehviläinen <i>et al.</i> (2006)	<i>Betula pendula</i>	Leaf rollers	O	More	-1.48	5	0.36	5	0.54	0.11
19	Montagnini <i>et al.</i> (1995)	<i>Calophyllum brasiliense</i>	<i>Attia cephalotes</i>	P		0.36	4	0.15	4	0.14	0.03
20	Brown & Ewel (1987)	<i>Cordia alliodora</i>	?	?		0.93	20	5.10	180	2.10	3.20
21	Gantner (2000)	<i>Corylus avellana</i>	<i>Myzocallis coryli</i>	O		-0.52	28	3.25	28	8.88	10.65
22	Montagnini <i>et al.</i> (1995)	<i>Dipteryx panamensis</i>	<i>Attia cephalotes</i>	P		-0.56	4	0.10	4	0.12	0.02
23	Smith <i>et al.</i> (1989)	<i>Eucalyptus botryoides</i>	<i>Cardiaspina fuscata</i>	O		-9.96	3	17.70	3	81.30	5.11
24	Zanuncio <i>et al.</i> (1998)	<i>Eucalyptus cloeziana</i>	<i>Lepidoptera</i> sp.	O		-2.06	5	1136.84	5	3326.32	959.35
25	Bigger (1985)	<i>Eucalyptus deglupta</i>	<i>Amblypelta cocophaga</i>	P		-0.28	13	0.33	13	0.41	0.25
26	Greaves (1966)	<i>Eucalyptus regnans</i>	<i>Chrysopharta bimaculata</i>	O	Equal	-0.19	2	7.00	2	9.25	6.83
27	Greaves (1966)	<i>Eucalyptus regnans</i>	<i>Chrysopharta bimaculata</i>	O	More	-0.59	2	3.90	2	9.25	5.20
28	Montagnini <i>et al.</i> (1995)	<i>Gonipa americana</i>	<i>Attia cephalotes</i>	P		-1.05	4	0.07	4	0.09	0.02
29	Nichols <i>et al.</i> (1999)	<i>Militia excelsa</i>	<i>Phytomyia lata</i>	O	Equal	-0.04	2	11.36	2	12.26	11.91
30	Nichols <i>et al.</i> (1999)	<i>Militia excelsa</i>	<i>Phytomyia lata</i>	O	More	-0.37	2	6.69	2	12.26	8.71
31	Nichols <i>et al.</i> (1999)	<i>Militia excelsa</i>	<i>Phytomyia lata</i>	O	More	-0.76	2	2.86	2	12.26	7.09
32	Peterson & Örlander (2003)	<i>Picea abies</i>	<i>Hyllobius abietis</i>	P		-0.43	4	10.15	4	17.60	15.06
33	Peterson & Örlander (2003)	<i>Picea abies</i>	<i>Hyllobius abietis</i>	P		-0.30	4	3.35	4	5.08	5.01
34	Örlander <i>et al.</i> (2001)	<i>Picea abies</i>	<i>Hyllobius abietis</i>	P		-1.84	4	54.00	4	140.00	40.73
35	Örlander <i>et al.</i> (2001)	<i>Picea abies</i>	<i>Hyllobius abietis</i>	P		-1.72	4	67.30	4	82.10	7.47
36	Nordlander <i>et al.</i> (2003)	<i>Picea abies</i>	<i>Hyllobius abietis</i>	P		-1.12	50	4.16	50	9.04	4.33
37	Nordlander <i>et al.</i> (2003)	<i>Picea abies</i>	<i>Hyllobius abietis</i>	P		-0.87	50	3.22	50	6.41	3.63
38	Nordlander <i>et al.</i> (2003)	<i>Picea abies</i>	<i>Hyllobius abietis</i>	P		-0.69	50	0.94	50	2.62	2.42
39	Nordlander <i>et al.</i> (2003)	<i>Picea abies</i>	<i>Hyllobius abietis</i>	P		-0.37	50	3.63	50	4.96	3.54

Appendix 1 (Continued)

Individual study	Authors	Tree species	Insect species	Insect host specificity	RAH*	Hedges' <i>d</i>	N^E	X^E	N^C	X^C	<i>S</i>
40	Nordlander <i>et al.</i> (2003)	<i>Pinus abies</i>	<i>Hylobius abietis</i>	P		0.04	50	3.08	50	2.96	2.87
41	Nordlander <i>et al.</i> (2003)	<i>Pinus abies</i>	<i>Hylobius abietis</i>	P		-0.76	50	0.55	50	2.00	1.90
42	Nordlander <i>et al.</i> (2003)	<i>Pinus abies</i>	<i>Hylobius abietis</i>	P		-0.66	50	3.85	50	6.07	3.33
43	Taylor <i>et al.</i> (1996)	<i>Pinus glauca</i>	<i>Pissodes strobi</i>	O	More	-2.79	3	6.78	3	18.34	3.31
44	Taylor <i>et al.</i> (1996)	<i>Pinus glauca</i>	<i>Pissodes strobi</i>	O	Equal	-1.24	2	12.65	3	18.34	3.34
45	McLean (1989)	<i>Pinus sitchensis</i>	<i>Pissodes strobi</i>	O		-0.79	6	5.33	6	17.00	13.58
46	McLean (1989)	<i>Pinus sitchensis</i>	<i>Pissodes strobi</i>	O		-0.48	6	2.00	6	9.33	14.10
47	Bae <i>et al.</i> (1997)	<i>Pinus densiflora</i>	<i>Thecodiplosis japonensis</i>	O		-0.29	30	39.45	30	44.75	18.06
48	Bae <i>et al.</i> (1997)	<i>Pinus densiflora</i>	Mite	?		-0.78	4	1.14	4	2.00	0.96
49	Bae <i>et al.</i> (1997)	<i>Pinus densiflora</i>	Scale insect	?		0.19	4	0.36	4	0.31	0.23
50	Bae <i>et al.</i> (1997)	<i>Pinus densiflora</i>	Leaf aphid	?		-0.53	4	0.21	4	0.39	0.29
51	Géri (1980)	<i>Pinus nigra laricio</i>	<i>Thaumetopoea pytiocampa</i>	O	Equal	-0.59	10	0.84	15	1.29	0.73
52	Géri (1980)	<i>Pinus nigra laricio</i>	<i>Thaumetopoea pytiocampa</i>	O	More	-1.91	10	0.05	15	1.29	0.63
53	Jactel <i>et al.</i> (2006)	<i>Pinus pinaster</i>	<i>Matsucoccus feytaudi</i>	O	Less	-0.18	25	10.68	25	13.16	13.91
54	Jactel <i>et al.</i> (2006)	<i>Pinus pinaster</i>	<i>Matsucoccus feytaudi</i>	O	Less	-0.26	25	15.32	25	18.84	13.53
55	Jactel <i>et al.</i> (2006)	<i>Pinus pinaster</i>	<i>Matsucoccus feytaudi</i>	O	More	-1.39	25	0.24	25	19.80	13.90
56	Jactel <i>et al.</i> (2006)	<i>Pinus pinaster</i>	<i>Matsucoccus feytaudi</i>	O	Less	-0.45	25	17.64	25	25.28	16.82
57	Jactel <i>et al.</i> (2006)	<i>Pinus pinaster</i>	<i>Matsucoccus feytaudi</i>	O	Less	-0.32	25	12.32	25	18.08	17.93
58	Gottschalk & Twery (1989)	<i>Pinus strobus</i>	<i>Lymantria dispar</i>	P		0.84	3	0.59	3	0.28	0.29
59	Gottschalk & Twery (1989)	<i>Pinus strobus</i>	<i>Lymantria dispar</i>	P		1.87	3	0.80	3	0.28	0.22
60	Brown <i>et al.</i> (1988)	<i>Pinus strobus</i>	<i>Lymantria dispar</i>	P		1.02	4	48.13	4	17.18	26.50
61	Brown <i>et al.</i> (1988)	<i>Pinus strobus</i>	<i>Lymantria dispar</i>	P		0.91	4	48.80	4	17.18	30.34
62	Katovich (1992)	<i>Pinus strobus</i>	<i>Pissodes strobi</i>	O	Less	-4.00	2	37.80	2	68.70	4.41
63	Katovich (1992)	<i>Pinus strobus</i>	<i>Pissodes strobi</i>	O	Less	-6.77	2	15.40	2	68.70	4.50
64	Katovich (1992)	<i>Pinus strobus</i>	<i>Pissodes strobi</i>	O	Equal	-9.47	2	5.70	2	68.70	3.80
65	Katovich (1992)	<i>Pinus strobus</i>	<i>Pissodes strobi</i>	O	More	-9.03	2	4.20	2	68.70	4.08
66	Stadniskii (1978)	<i>Pinus sylvestris</i>	<i>Hylobius abietis</i>	P		-3.40	4	0.24	6	0.68	0.12
67	Von Sydow & Ölander (1994)	<i>Pinus sylvestris</i>	<i>Hylobius abietis</i>	P		-1.64	3	13.16	4	57.89	22.94
68	Schowalter & Turchin (1993)	<i>Pinus taeda</i>	<i>Dendroctonus frontalis</i>	O		0.16	2	1.00	2	0.80	0.71
69	Schowalter & Turchin (1993)	<i>Pinus taeda</i>	<i>Dendroctonus frontalis</i>	O		-0.76	2	0.80	2	5.50	3.54
70	Miller & Stephen (1983)	<i>Pinus taeda</i>	<i>Rhyacionia frustrana</i>	O	More	-0.24	7	13.65	7	18.70	19.38
71	Miller & Stephen (1983)	<i>Pinus taeda</i>	<i>Rhyacionia frustrana</i>	O	Equal	-0.11	7	15.99	7	18.70	22.37
72	Berisford & Kulman (1967)	<i>Pinus taeda</i>	<i>Rhyacionia frustrana</i>	O		-0.39	9	36.33	9	43.56	17.77
73	Nowak <i>et al.</i> (2003)	<i>Pinus taeda</i>	<i>Rhyacionia frustrana</i>	O		-3.62	3	21.00	3	50.70	6.56
74	Nowak <i>et al.</i> (2003)	<i>Pinus taeda</i>	<i>Rhyacionia frustrana</i>	O		-1.62	3	37.10	3	50.60	6.68
75	White & Whitham (2000)	<i>Populus angustifolia</i>	<i>Alsophila pomataria</i>	P		1.09	8	32.00	8	13.00	16.49
76	White & Whitham (2000)	<i>Populus angustifolia</i>	<i>Alsophila pomataria</i>	P		0.81	19	48.00	19	26.00	26.51
77	Fauss & Pierce (1969)	<i>Pseudotsuga menziesii</i>	<i>Choristoneura occidentalis</i>	O	More	-1.70	4	34.90	4	60.83	13.26
78	Fauss & Pierce (1969)	<i>Pseudotsuga menziesii</i>	<i>Choristoneura occidentalis</i>	O	More	-1.86	4	39.35	4	60.83	10.03

Appendix 1 (Continued)

Individual study	Authors	Tree species	Insect species	Insect host specificity	RAH*	Hedges' d	N^E	X^E	N^C	X^C	S
79	Faust & Pierce (1969)	<i>Pseudotsuga menziesii</i>	<i>Choristoneura occidentalis</i>	O	Equal	-0.34	4	57.28	4	60.83	9.17
80	Faust & Pierce (1969)	<i>Pseudotsuga menziesii</i>	<i>Choristoneura occidentalis</i>	O	Less	-0.11	4	59.48	4	60.83	10.33
81	Faust & Pierce (1969)	<i>Pseudotsuga menziesii</i>	<i>Choristoneura occidentalis</i>	O	More	-2.77	4	8.08	4	42.75	10.88
82	Faust & Pierce (1969)	<i>Pseudotsuga menziesii</i>	<i>Choristoneura occidentalis</i>	O	More	-1.13	4	24.20	4	42.75	14.28
83	Faust & Pierce (1969)	<i>Pseudotsuga menziesii</i>	<i>Choristoneura occidentalis</i>	O	Equal	-0.33	4	37.50	4	42.75	13.63
84	Faust & Pierce (1969)	<i>Pseudotsuga menziesii</i>	<i>Choristoneura occidentalis</i>	O	Less	0.02	4	43.00	4	42.75	12.22
85	Faust & Pierce (1969)	<i>Pseudotsuga menziesii</i>	<i>Choristoneura occidentalis</i>	O	More	-1.01	4	41.70	4	61.95	17.44
86	Faust & Pierce (1969)	<i>Pseudotsuga menziesii</i>	<i>Choristoneura occidentalis</i>	O	More	-0.61	4	48.95	4	61.95	18.44
87	Faust & Pierce (1969)	<i>Pseudotsuga menziesii</i>	<i>Choristoneura occidentalis</i>	O	Equal	0.00	4	61.85	4	61.95	17.91
88	Faust & Pierce (1969)	<i>Pseudotsuga menziesii</i>	<i>Choristoneura occidentalis</i>	O	Less	0.16	4	65.10	4	61.95	17.18
89	Faust & Pierce (1969)	<i>Pseudotsuga menziesii</i>	<i>Choristoneura occidentalis</i>	O	More	-1.78	4	19.70	4	57.65	18.53
90	Faust & Pierce (1969)	<i>Pseudotsuga menziesii</i>	<i>Choristoneura occidentalis</i>	O	More	-0.63	4	40.60	4	57.65	23.42
91	Faust & Pierce (1969)	<i>Pseudotsuga menziesii</i>	<i>Choristoneura occidentalis</i>	O	Equal	-0.05	4	56.55	4	57.65	19.54
92	Faust & Pierce (1969)	<i>Pseudotsuga menziesii</i>	<i>Choristoneura occidentalis</i>	O	Less	0.26	4	63.30	4	57.65	18.93
93	Moore <i>et al.</i> (1991)	<i>Quercus petraea</i>	<i>Phylllobius argentatus</i>	P		0.78	3	231.00	3	125.33	108.94
94	Moore <i>et al.</i> (1991)	<i>Quercus petraea</i>	?	O		-0.47	3	488.33	3	538.67	86.58
95	Moore <i>et al.</i> (1991)	<i>Quercus petraea</i>	<i>Phyllonorycter</i> spp.	P		-0.21	3	11.00	3	12.33	5.02
96	Moore <i>et al.</i> (1991)	<i>Quercus petraea</i>	<i>Stigmella</i> spp.	P		3.30	3	18.00	3	2.00	3.87
97	Moore <i>et al.</i> (1991)	<i>Quercus petraea</i>	?	P		1.42	3	345.00	3	68.67	155.91
98	Moore <i>et al.</i> (1991)	<i>Quercus petraea</i>	<i>Neuroterus</i> spp.	O		-0.54	3	0.74	3	3.19	3.63
99	Moore <i>et al.</i> (1991)	<i>Quercus petraea</i>	<i>Phylllobius argentatus</i>	P		1.21	3	295.67	3	125.33	113.07
100	Moore <i>et al.</i> (1991)	<i>Quercus petraea</i>	?	O		-0.27	3	500.33	3	538.67	114.20
101	Moore <i>et al.</i> (1991)	<i>Quercus petraea</i>	<i>Phyllonorycter</i> spp.	P		-2.41	3	1.00	3	12.33	3.76
102	Moore <i>et al.</i> (1991)	<i>Quercus petraea</i>	<i>Stigmella</i> spp.	P		0.15	3	2.67	3	2.00	3.49
103	Moore <i>et al.</i> (1991)	<i>Quercus petraea</i>	?	P		-1.35	3	21.33	3	68.67	28.00
104	Moore <i>et al.</i> (1991)	<i>Quercus petraea</i>	<i>Neuroterus</i> spp.	O		-0.58	3	0.52	3	3.19	3.66
105	Soria <i>et al.</i> (1995)	<i>Quercus rotundifolia</i>	<i>Curculio elphas</i>	O		0.28	7	11.61	7	9.20	7.98
106	Soria <i>et al.</i> (1995)	<i>Quercus suber</i>	<i>Curculio elphas</i>	O		-0.36	7	9.29	7	12.10	7.35
107	Wazihullah <i>et al.</i> (1996)	<i>Somneratia apetala</i>	<i>Zeuzera conferta</i>	P		-0.67	20	15.00	29	27.00	17.66
108	Wazihullah <i>et al.</i> (1996)	<i>Somneratia apetala</i>	<i>Zeuzera conferta</i>	P		-0.40	36	42.00	34	54.00	29.56
109	Wazihullah <i>et al.</i> (1996)	<i>Somneratia apetala</i>	<i>Zeuzera conferta</i>	P		-0.77	31	32.00	24	51.00	24.40
110	Wazihullah <i>et al.</i> (1996)	<i>Somneratia apetala</i>	<i>Zeuzera conferta</i>	P		-2.06	13	22.00	13	56.00	15.98
111	Wazihullah <i>et al.</i> (1996)	<i>Somneratia apetala</i>	<i>Zeuzera conferta</i>	P		0.20	12	17.00	12	13.00	19.20
112	Folgarait <i>et al.</i> (1995)	<i>Strychnodendron microstachyum</i>	<i>Cecidomyiidae</i> sp.	O		-0.03	80	0.14	16	0.15	0.24
113	Folgarait <i>et al.</i> (1995)	<i>Strychnodendron microstachyum</i>	<i>Euclystis</i> sp.	P		0.31	80	0.02	16	0.01	0.02
114	Montagnini <i>et al.</i> (1995)	<i>Strychnodendron microstachyum</i>	<i>Atta cephalotes</i>	P		-5.49	4	0.41	4	0.87	0.07
115	Keenan <i>et al.</i> (1995)	<i>Toona ciliata</i>	<i>Hypsipyla robusta</i>	O	Equal	-4.58	2	0.23	2	0.71	0.06
116	Keenan <i>et al.</i> (1995)	<i>Toona ciliata</i>	<i>Hypsipyla robusta</i>	O	More	-7.44	3	0.10	2	0.71	0.06
117	Montagnini <i>et al.</i> (1995)	<i>Vitrola koschnyi</i>	<i>Atta cephalotes</i>	P		-1.39	4	0.14	4	0.25	0.07

Appendix 1 (Continued)

Individual study	Authors	Tree species	Insect species	Insect host specificity	RAH*	Hedges' d	N^E	X^E	N^C	X^C	S
118	Montagnini <i>et al.</i> (1995)	<i>Vochysia ferruginea</i>	<i>Atta cephalotes</i>	P		-0.49	4	0.06	4	0.06	0.01
119	Montagnini <i>et al.</i> (1995)	<i>Vochysia guatemalensis</i>	<i>Atta cephalotes</i>	P		0.81	4	0.12	4	0.09	0.03

Insect host specificity: O for oligophagous, P for polyphagous; RAH: relative abundance of host tree species in the mixture; Hedges' d : Hedges' unbiased effect size of insect herbivory; N^E : sample size of the experimental group (mixed stand); X^E : mean value of the experimental group (mixed stand); N^C : sample size of the control group (pure stand); X^C : mean value of the control group (pure stand); S : pooled standard deviation.

*RAH in bold indicates the studies for which at least two levels of host tree abundance could be compared (see Fig. 2).