


# **$t$ -tests**

ENTMLGY 6707 Entomological Techniques and Data Analysis



# Learning objectives

- 1) Identify the information required to conduct a  $t$ -test
- 2) Distinguish between types of  $t$ -tests
- 3) Interpret the outcome of  $t$ -tests

# ***t*-tests**

Used to determine if the means of two groups have statistically clear differences OR used to determine if a mean differs from a specified value (e.g., population mean or 0).

# ***t*-tests**

Used to determine if the means of two groups have statistically clear differences OR used to determine if a mean differs from a specified value (e.g., population mean or 0).

Assumptions:

- 1.
- 2.
- 3.
- 4.

**Activity:** Based on what you have learned so far about experimental design, sampling, and distributions, list assumptions of the standard t-test

# ***t*-tests**

Used to determine if the means of two groups have statistically clear differences OR used to determine if a mean differs from a specified value (e.g., population mean or 0).

## Assumptions:

1. Response (aka dependent) variable (the one we are comparing between groups) is continuous
- 2.
- 3.
- 4.

**Activity:** Based on what you have learned so far about experimental design, sampling, and distributions, list assumptions of the standard t-test

# ***t*-tests**

Used to determine if the means of two groups have statistically clear differences OR used to determine if a mean differs from a specified value (e.g., population mean or 0).

## Assumptions:

1. Response (aka dependent) variable (the one we are comparing between groups) is continuous
2. Observations are independent (typically meaning they comprise a random sample)
- 3.
- 4.

**Activity:** Based on what you have learned so far about experimental design, sampling, and distributions, list assumptions of the standard t-test

# ***t*-tests**

Used to determine if the means of two groups have statistically clear differences OR used to determine if a mean differs from a specified value (e.g., population mean or 0).

## Assumptions:

1. Response (aka dependent) variable (the one we are comparing between groups) is continuous
2. Observations are independent (typically meaning they comprise a random sample)
3. Response variable is normally distributed
- 4.

**Activity:** Based on what you have learned so far about experimental design, sampling, and distributions, list assumptions of the standard t-test

# ***t*-tests**


Used to determine if the means of two groups have statistically clear differences OR used to determine if a mean differs from a specified value (e.g., population mean or 0).

## Assumptions:


1. Response (aka dependent) variable (the one we are comparing between groups) is continuous
2. Observations are independent (typically meaning they comprise a random sample)
3. Response variable is normally distributed
4. Variances of the response variable are equal across groups (= homogeneity of variances)


**Activity:** Based on what you have learned so far about experimental design, sampling, and distributions, list assumptions of the standard t-test





Credit: Liz Thielen






Parametric Test	Non-Parametric Test
Paired t-test	Wilcoxon rank sum test
Unpaired t-test	Mann-Whitney U test
Pearson correlation	Spearman correlation
One-way Analysis of Variance	Kruskal-Wallis test

# William Sealy Gosset

- 1876 – 1937
- Head Experimental Brewer of Guinness
- Developed Student's t-distribution
- Argued – in the literature and probably at the pub – with Ronald Fisher
- Had awesome mustache



$$f(t) = \frac{\Gamma((r+1)/2)}{\sqrt{\pi r} \Gamma(r/2)} \cdot \frac{1}{(1+t^2/r)^{(r+1)/2}}$$

Note:  $r$  and  $t$  in the above equation are indicated by  $\nu$  and  $x$  on the figure

## One sample *t*-test

(includes paired *t*-tests)

$$t = \frac{\bar{x} - \mu_0}{s / \sqrt{n}}$$

$$s_N = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \bar{x})^2}$$

$$\text{d.f.} = n - 1$$


## Two sample *t*-test

These equations are for when your two groups have unequal sample sizes and variances


$$t = \frac{\bar{X}_1 - \bar{X}_2}{s_{\bar{\Delta}}}$$

$$s_{\bar{\Delta}} = \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$$

$$\text{d.f.} = \frac{\left( \frac{s_1^2}{n_1} + \frac{s_2^2}{n_2} \right)^2}{\frac{(s_1^2/n_1)^2}{n_1 - 1} + \frac{(s_2^2/n_2)^2}{n_2 - 1}}$$



$$p \leq 0.05$$



# Variable effects of temperature on insect herbivory

Nathan P. Lemoine<sup>1</sup>, Deron E. Burkepile<sup>1</sup> and John D. Parker<sup>2</sup>

<sup>1</sup> Department of Biological Sciences, Florida International University, Miami, FL, United States

<sup>2</sup> Smithsonian Environmental Research Center, Smithsonian Institution, Edgewater, MD, United States



Explore data | Search

Who we are | What we do | Join us | Help | Login

## Data from: Variable effects of temperature on insect herbivory

Lemoine, Nathan P., Florida International University

Burkepile, Deron E., Florida International University

Parker, John D., Smithsonian Institution

Published May 01, 2015 on Dryad. <https://doi.org/10.5061/dryad.9fd75>

Cite this dataset

Lemoine, Nathan P.; Burkepile, Deron E.; Parker, John D. (2015). Data from: Variable effects of temperature on insect herbivory [Dataset]. Dryad. <https://doi.org/10.5061/dryad.9fd75>

## Abstract

Rising temperatures can influence the top-down control of plant biomass by increasing herbivore metabolic demands. Unfortunately, we know relatively little about the effects of temperature on herbivory rates for most insect herbivores in a given community. Evolutionary history, adaptation to local environments, and dietary factors may lead to variable thermal response curves across different species. Here we characterized the effect of temperature on herbivory rates for 21 herbivore-plant pairs, encompassing 14 herbivore and 12 plant species. We show that overall consumption rates increase with temperature between 20 and 30 C but do not increase further with increasing temperature. However, there is substantial variation in thermal responses among individual herbivore-plant pairs at the highest temperatures. Over one third of the herbivore-plant pairs showed declining consumption rates at high temperatures, while an approximately equal number showed increasing consumption rates. Such variation existed even within herbivore species, as some species exhibited idiosyncratic thermal response curves on different host plants. Thus, rising temperatures, particularly with respect to climate change, may have highly variable effects on plant-herbivore interactions and, ultimately, top-down control of plant biomass.

## Data files

Download dataset

> May 01, 2015

## Related works

### Primary article

<https://doi.org/10.7717/peerj.376>

## Share



## Metrics

309 views

99 downloads

1 citations



**Paper:** <https://doi.org/10.7717/peerj.376>

**Data:** <https://doi.org/10.5061/dryad.9fd75>

# Variable effects of temperature on insect herbivory

Nathan P. Lemoine<sup>1</sup>, Deron E. Burkepile<sup>1</sup> and John D. Parker<sup>2</sup>

<sup>1</sup> Department of Biological Sciences, Florida International University, Miami, FL, United States

<sup>2</sup> Smithsonian Environmental Research Center, Smithsonian Institution, Edgewater, MD, United States

**Example of what NOT to do!!**



Explore data | Search

Who we are | What we do | Join us | Help | Login

## Data from: Variable effects of temperature on insect herbivory

Lemoine, Nathan P., Florida International University

Burkepile, Deron E., Florida International University

Parker, John D., Smithsonian Institution

Published May 01, 2015 on Dryad. <https://doi.org/10.5061/dryad.9fd75>

Cite this dataset

Lemoine, Nathan P.; Burkepile, Deron E.; Parker, John D. (2015). Data from: Variable effects of temperature on insect herbivory [Dataset]. Dryad. <https://doi.org/10.5061/dryad.9fd75>

## Abstract

Rising temperatures can influence the top-down control of plant biomass by increasing herbivore metabolic demands. Unfortunately, we know relatively little about the effects of temperature on herbivory rates for most insect herbivores in a given community. Evolutionary history, adaptation to local environments, and dietary factors may lead to variable thermal response curves across different species. Here we characterized the effect of temperature on herbivory rates for 21 herbivore-plant pairs, encompassing 14 herbivore and 12 plant species. We show that overall consumption rates increase with temperature between 20 and 30 C but do not increase further with increasing temperature. However, there is substantial variation in thermal responses among individual herbivore-plant pairs at the highest temperatures. Over one third of the herbivore-plant pairs showed declining consumption rates at high temperatures, while an approximately equal number showed increasing consumption rates. Such variation existed even within herbivore species, as some species exhibited idiosyncratic thermal response curves on different host plants. Thus, rising temperatures, particularly with respect to climate change, may have highly variable effects on plant-herbivore interactions and, ultimately, top-down control of plant biomass.

## Data files

Download dataset

> May 01, 2015

## Related works

Primary article

<https://doi.org/10.7717/peerj.376>

## Share



## Metrics

309 views

99 downloads

1 citations

## **NO METADATA INCLUDED**



**Paper:** <https://doi.org/10.7717/peerj.376>

**Data:** <https://doi.org/10.5061/dryad.9fd75>