

Sets, Relations and Functions

1 The Axiom of Extension

Sets (or collections) have elements (or members). Any object (concrete or otherwise), may be an element of a set. This extends to sets, which may be an element of some other set. In fact, it is generally sufficient to only discuss sets of sets, as the properties will extend to all other sets. There are some ways to relate sets: belonging, inclusion and equality.

If an object x is an element of set A , we say that it *belongs* to A , or " x is contained in A ". Denote this as $x \in A$.

If a set A is *included* in some set B , then A is a subset of B . Denote this as $A \subset B$ or $B \supset A$. Some related definitions: set inclusion is *reflexive* since a set is included in itself ($A \subset A$)¹; set inclusion is *transitive* as $A \subset B$ and $B \subset C$ implies $A \subset C$. (Conversely, belonging is neither reflexive or transitive).

Sets can also be related by equality ($A = B$), as defined by the Axiom of Extension.

Axiom 1 (Axiom of Extension). *Two sets are equal if and only if they have the same elements.*

In terms of inclusion, this can be rewritten as " $A \subset B$ and $B \subset A \Leftrightarrow A = B$ ".

A set is said to be a *proper* subset of another if $A \subset B$ and $A \neq B$; that is, all subsets are proper except from the set itself, which is a subset but not proper.

Equality is *symmetric*, that is $A = B \Leftrightarrow B = A$, whereas set inclusion is *antisymmetric* (that is, $A \subset B$ and $B \subset A \Rightarrow A = B$)².

2 The Axiom of Specification

The basis of most Set Theoretic principles is to construct new sets from old sets. The most important such principle is the so called "*Aussonderungsaxiom*":

Axiom 2 (Axiom of Specification). *To every set A and every condition $S(x)$ there corresponds a set B whose elements are exactly those elements x of A for which $S(x)$ holds.*

That is to say, anything one can say about the elements of a set defines a subset. The condition must form a valid *sentence*. A sentence is formed using the atomic sentences of belonging ($x \in A$) and equality ($A = B$), which are used to create more specific sentences using the following logical operators:

- and
- or
- if and only if
- not
- if - then - (equivalently *implies*)
- for some (equivalently *there exists*)
- for all.

¹Clearly, $A \subset B$ is the same as $A \subseteq B$ in this notation.

²Antisymmetry is defined by "*if aRb and bRa , then $a = b$* " for some binary relation R .

There are a few general rules for sentence construction³:

1. *and*, *or* and *if and only if* are placed between two sentences
2. *not* can be placed before a sentence enclosed by parentheses
3. the dashes in *if - then -* are replaced with sentences
4. follow *for some* or *for all* by a letter, in turn followed by a sentence

We can write the Axiom of Specification symbolically as: $B = \{x \in A : S(x)\}$. This set B is uniquely specified due to the Axiom of Extension.

Remark 1 (Russell paradox). *Nothing contains everything (or equivalently, there is no universe)*

Proof. Assume there is a set A which contains everything.

Consider the condition: $\text{not } (x \in x)$. Let us rewrite this as $(x \notin x)$ for ease of use.

Constructing the set B from A where this condition holds:

$$B = \{x \in A : x \notin x\}$$

Following from the Axiom of Extension,

$$y \in B \text{ if and only if } (y \in A \text{ and } y \notin y)$$

For A to contain everything, this must hold for any y . Consider $y = B$. If $B \in A$, then $B \in B \Leftrightarrow B \notin B$. This is clearly a contradiction so A does not contain B and by extension, A cannot contain everything. \square

The Russell Paradox gives us an example of a condition that produces an illegal set. Some texts name these illegal sets as "classes".

3 Unordered Pairs

In order to make any progress it is necessary for us to make an assumption: *there exists a set*. This will be formulated more carefully later on but for now this is sufficient.

Given the assumption that there exists a set, one can easily construct the *empty set* using the Axiom of Specification, using the condition $x \neq x$. Clearly there are no elements of such a set. This set is unique by the Axiom of Extension. Denote this as $\{x \in A : x \neq x\} = \emptyset$.

For any set A , $\emptyset \subset A$ as every element in \emptyset is in A . This is a vacuous truth.⁴

Axiom 3 (Axiom of Pairing). *For any two sets there exists a set that they both belong to.*

That is, for some sets a and b , there exists a set A such that $a \in A$ and $b \in A$.

Consider the set that contains a and b and nothing else⁵. This can be constructed from the Axiom of Specification using the condition $S(x) = "x = a \text{ or } x = b"$. By the Axiom of

³Some general notes: the set of a single object is not the same as that object itself; "for some y ($x \in A$)" is equivalent to " $x \in A$ "; "for some x ($x \in A$)" and "for some y ($y \in A$)" are equivalent.

⁴The argument can also be made from the other direction: the only case where $\emptyset \subset A$ is false is where there is some element in \emptyset that is not contained in A . Since \emptyset is empty, this doesn't make any sense and so the statement can never be false.

⁵The existence of such a set is actually equivalent to the Axiom of Pairing: if there is some set A such that $a \in A$ and $b \in A$, then we can construct a set with only a and b in using the sentence " $x = a \text{ or } x = b$ ".

Extension, this set is unique. We call this the *pair* or, more specifically, the *unordered pair* formed by a and b . Denote this set as

$$\{x \in A : x = a \text{ or } x = b\} = \{a, b\}. \quad (1)$$

Note that this isn't the same as the set of all elements in a or b but the set containing only two elements: a and b .

The *singleton* of a is a special case unordered pair $\{a, a\}$, denoted by $\{a\}$, where the set has a as its only element. $a \in A$ is equivalent to $\{a\} \subset A$.

Considering the unordered pair of any set and the empty set, it follows from the Axiom of Pairing that every set is an element of some other set. We can also infer the existence of infinitely many sets by construction: first consider the singletons $\emptyset, \{\emptyset\}, \{\{\emptyset\}\}$ etc.; now consider the pairs of such sets, and the pairs of these and the singletons and so on. Note that all of these sets are unique by the Axiom of Extension.

If $S(x)$ is a condition on x , then the set of all x that satisfy this condition is given by $\{x : S(x)\}$, granted that this is a valid set. Rewriting the generic set constructed by the Axiom of Specification,

$$\{x \in A : S(x)\} = \{x : x \in A \text{ and } S(x)\}.$$

4 Unions and Intersections