

# BESIII Analysis Memo

DocDB-497

BAM-228

June 18, 2019

## Amplitude analysis of $D_s^+ \rightarrow K^+ K^- \pi^+$

Meng Wang<sup>a</sup>, Yu Lu<sup>a</sup>, and Liaoyuan Dong<sup>a</sup>, and Huaimin Liu<sup>a</sup><sup>a</sup>*Institute of High Energy Physics, CAS*

## Internal Referee Committee

Ref1 xx (Chair)<sup>d</sup>, Ref2 xx<sup>e</sup>, and Ref3 xx<sup>f</sup><sup>d</sup>*Department of Computer Science and Engineering*<sup>e</sup>*Department of Electrical Engineering*<sup>f</sup>*Latex Univeristy*DocDB : <http://docbes3.ihep.ac.cn/cgi-bin/DocDB/ShowDocument?docid=497>Hypernews : <http://hnb3.ihep.ac.cn/HyperNews/get/paper228.html>

## Abstract

We have performed an amplitude analysis of  $D_s^+ \rightarrow K^+ K^- \pi^+$  using a data sample of  $3.19fb^{-1}$  recorded with BESIII detector at a center-of-mass energy of 4.178GeV. About 4000 events with %0.4 background are used in amplitude analysis. There are 6 intermediate resonances in the model which does the best fit to data. We tent to agree the Dalitz plot analysis of  $D_s^+ \rightarrow K^+ K^- \pi^+$  by BABAR. And we also measured the branching fraction of  $D_s^+ \rightarrow K^+ K^- \pi^+$  and the result is  $\mathcal{B}(D_s^+ \rightarrow K^+ K^- \pi^+) = (5.38 \pm 0.07_{stat.} \pm 0.11_{sys.})\%$ .

# Contents

<b>1</b>	<b>1 Introduction</b>	<b>2</b>
<b>2</b>	<b>2 Data Set and Monte Carlo Samples</b>	<b>3</b>
<b>3</b>	<b>3 Event Selection</b>	<b>4</b>
3.1	Tracking, PID, $\pi^0/\eta^{(\prime)}$ and $K_S^0$ Reconstruction . . . . .	4
3.2	Signal Selection . . . . .	5
<b>4</b>	<b>4 Model Independent Partial Wave Analysis</b>	<b>6</b>
4.1	Signal Candidate Selection . . . . .	6
4.2	Background Analysis . . . . .	6
4.3	MIPWA method . . . . .	8
4.4	S-wave parameterization at the $K^+K^-$ threshold . . . . .	11
<b>5</b>	<b>5 Amplitude Analysis</b>	<b>14</b>
5.1	Event Selection for Amplitude Analysis . . . . .	14
5.2	Background Analysis . . . . .	14
5.3	Fit Method . . . . .	14
5.4	Fit Fraction . . . . .	17
5.5	Fit Result . . . . .	17
5.6	Systematic Uncertainties . . . . .	19
<b>6</b>	<b>6 Branching Fraction Measurements</b>	<b>24</b>
6.1	Event Selection for Branching Fraction Measurements . . . . .	24
6.2	Analytic Strategy of Branching Fraction Measurement . . . . .	24
6.3	Results of Branching Fraction . . . . .	27
6.4	Systematic Uncertainties . . . . .	27
<b>7</b>	<b>7 Summary</b>	<b>29</b>
<b>Appendices</b>		<b>31</b>
<b>A</b>	<b>To be supplemented</b>	<b>32</b>

Table 1: previous analyses of this decay channel.

Amplitude	BABAR	CLEO-c	E687
$D_s^+ \rightarrow \bar{K}^*(892)^0 K^+$	$47.9 \pm 0.5 \pm 0.5$	$47.4 \pm 1.5 \pm 0.4$	$47.8 \pm 4.6 \pm 4.0$
$D_s^+ \rightarrow \phi(1020)\pi^+$	$41.4 \pm 0.8 \pm 0.5$	$42.2 \pm 1.6 \pm 0.3$	$39.6 \pm 3.3 \pm 4.7$
$D_s^+ \rightarrow f_0(980)\pi^+ / a_0(980)\pi^+$	$16.4 \pm 0.7 \pm 2.0$	$28.2 \pm 1.9 \pm 1.8$	$11.0 \pm 3.5 \pm 2.6$
$D_s^+ \rightarrow \bar{K}_0^*(1430)^0 K^+$	$2.4 \pm 0.3 \pm 1.0$	$3.9 \pm 0.5 \pm 0.5$	$9.3 \pm 3.2 \pm 3.2$
$D_s^+ \rightarrow f_0(1710)\pi^+$	$1.1 \pm 0.1 \pm 0.1$	$3.4 \pm 0.5 \pm 0.3$	$3.4 \pm 2.3 \pm 3.5$
$D_s^+ \rightarrow f_0(1370)\pi^+$	$1.1 \pm 0.1 \pm 0.2$	$4.3 \pm 0.6 \pm 0.5$	...
$\Sigma FF(\%)$	$110.2 \pm 0.6 \pm 2.0$	$129.5 \pm 4.4 \pm 2.0$	111.1
$\chi^2/NDF$	$\frac{2843}{2305-14} = 1.2$	$\frac{178}{117} = 1.5$	$\frac{50.2}{33} = 1.5$
Events	$96307 \pm 369$	$12226 \pm 22$	$701 \pm 36$

## 1 Introduction

The decay  $D_s^+ \rightarrow K^+ K^- \pi^+$  is a Cabibbo-favored (CF) channel and has a large branching fraction for the  $D_s$  meson. Thus, this decay channel is usually used to normalize measurements of decay chains involving charm quarks.

Knowledge of the decay amplitude allows us to properly account for interference effects when measuring absolute hadronic branching fractions of  $D_s$  mesons.

Below is the table of previous analyses of this decay channel.

From Tab.1 [1], we can see an obvious difference of decay fraction of  $f_0(980)\pi^+$  between BABAR and CLEO-c. E687 used about 700 events and the E687 model did not take  $f_0(1370)\pi^+$  into account. For CLEO-c, about 14400 events with purity about 84.9% were selected with single tag method. The analysis of BABAR used about 100000 events with purity about 95%. In this analysis with double tag method, we can get a nearly background free data sample, that will be good to perform the amplitude analysis.

## 2 Data Set and Monte Carlo Samples

We use  $3.195 \text{ fb}^{-1}$  data set collected at  $E_{cm} = 4.178 \text{ GeV}$  by BESIII detector in 2016. Both data sample and Monte Carlo samples are reconstructed under BOSS7.0.3. All samples were generated with run-dependent  $E_{cm}$  [3], except for the Bhabha,  $\mu - \text{pair}$  and Two-photon fusion events. For these thress types of events, we used a constant  $E_{cm}$  at  $4178.37 \text{ MeV}$  which is twice of a luminosity-weighted average of the measured beam energy in the center-of-mass frame. Totally 40 rounds of generic MC with each round equaling to data size are used for background study, tag efficiencies estimation(rounds 01-30) and input-output check for branching fraction measurement(rounds 31-40). They are available at `/besfs3/offline/data/703-1/4180/mc/`. For each round of generic MC, the detail component and corresponding size of each Monte Carlo sample are shown in Tab.2.

Table 2: Component and corresponding size, assume luminosity = 3195/pb.

Component	cross section (pb)	Size(M)	directory
$D^0 D^0$	179	0.5719	D0D0
$D^+ D^-$	197	0.6294	DpDm
$D^{*0} D^0$	1211	3.8691	DST0D0
$D^{*+} D^-$	1296	4.1407	DSTpDm
$D^{*0} D^{*0}$	2173	6.9427	DST0DST0
$D^{*+} D^{*-}$	2145	6.8533	DSTpDSTm
$D_s^+ D_s^-$	7	0.0225	DsDs
$D_s^{*+} D_s^-$	961	3.0700	DsSTDs
$DD^* \pi^+$	383	1.2237	DDSTPIp
$DD^* \pi^0$	192	0.6134	DDSTPI0
$DD \pi^+$	50	0.1598	DDPIp
$DD \pi^0$	25	0.0799	DDPI0
Component	cross section (nb)	Size(M)	
$q\bar{q}$	13.8	44.0910	qq
$\gamma J/\psi$	0.40	1.2780	RR1S
$\gamma \psi(2S)$	0.42	1.3419	RR2S
$\gamma \psi(3770)$	0.06	0.1917	RR3770
$\tau\tau$	3.45	11.0228	tt
$\mu\mu$	5.24	16.7418	mm
$ee$	423.99	13.5465(0.01 $\times$ )	ee
$\gamma\gamma$	1.7	5.4315	TwoGam
HCT	0.10178	0.3252	HCT

For the Signal MC, we generate the signal events with one  $D_s$  decaying to signal mode using the generator “DIY”, in which the parameters are obtained from the fit to data. PHSP MC and Signal MC are used in MC integration required for the amplitude fit. The Signal MC is also used in the input/output check.

### 3 Event Selection

At  $E_{cm} = 4.178\text{GeV}$ , pairs of  $D_s D_s^*$  are produced, and the  $D_s^*$  decays to either  $D_s \gamma$  or  $D_s \pi^0$ . We use double tag method to select our signal events.

#### 3.1 Tracking, PID, $\pi^0/\eta^{(\prime)}$ and $K_S^0$ Reconstruction

$D_s$  candidates are built from  $K^\pm$ ,  $\pi^\pm$ ,  $\pi^0/\eta^{(0)}$  and  $K_S^0$ . The selections of the particles to build  $D_s$  candidates are performed with DTagAlg-00-01-05 package with the default setting, which are summarized below.

- Tracking:

- The properties of charged tracks are determined based on the MDC information. Charged track candidates must satisfy:

- $|\cos\theta| < 0.93$ .

- $|dr| < 1\text{cm}$  and  $|dz| < 10\text{cm}$ ,

where  $|dr|$  and  $|dz|$  are defined as the one reconstructed minus the interaction point.

- Particle ID:

- Charged tracks are identified as pion or kaon with Particle Identification (PID), which is implemented by combining the information of the energy loss (dE/dx) in MDC and the time-of-flight measured from the TOF system. Kaon and Pion are identified with the requirements that

- $\text{Prob}(K) > 0.00$  and  $\text{Prob}(K) > \text{Prob}(\pi)$  for  $K$ ,

- $\text{Prob}(\pi) > 0.00$  and  $\text{Prob}(\pi) > \text{Prob}(K)$  for  $\pi$ , where  $\text{Prob}(X)$  is the probability of hypothesis X, X can be  $\pi$  or  $K$ .

- $\pi^0/\eta$  selection:  $\pi^0$  candidates are reconstructed through  $\pi^0 \rightarrow \gamma\gamma$  with package of PioEtaToGGRecAlg.

The photons are reconstructed as energy showers on the EMC. We require:

- Minimum energy for barrel showers ( $|\cos\theta| < 0.8$ ):  $E_{min} > 25\text{MeV}/c^2$ ,

- Minimum energy for endcap showers ( $0.86 < |\cos\theta| < 0.92$ ):  $E_{min} > 50\text{MeV}/c^2$ ,

- Shower within other  $|\cos\theta|$  regions are rejected.

- EMC time requirements for events with at least one charged track detected:  $0 \leq t \leq 14(50\text{ns})$ ,

Then we perform a constrained fit on the photon pairs to the nominal  $\pi^0/\eta$  mass and require:

- The unconstrained invariant mass for  $\pi^0$ :  $0.115 < M(\gamma\gamma) < 0.015\text{GeV}/c^2$ ,

- The unconstrained invariant mass for  $\eta$ :  $0.490 < M(\eta) < 0.580\text{GeV}/c^2$ ,

- Mass fit:  $\chi^2_{1c} < 30$ .

- $\eta'$  selection: The  $\eta'$  candidates are reconstructed with  $\pi^+\pi^-\eta$ , the invariant mass for  $\pi^+\pi^-\eta$  is required to fall into the range of  $[0.938, 0.978]\text{GeV}^2$ .

- $K_S^0$  selection:  $K_S^0$  candidates are reconstructed using VeeVertexAlg package with two opposite charged tracks with requiring:

- $|\cos\theta| < 0.93$

- $|dz| < 20\text{cm}$

For each pair of tracks, a constrained vertex fit is performed and the track parameters' results are used to get the invariant mass  $M(K_S^0)$ . Then the decay length of  $K_S^0$  is obtained with second vertex fit by the SecondVertexFit package. For  $K_S^0$  selection, we require:

- $0.487\text{GeV}/c^2 < M(K_S^0) < 0.511\text{GeV}/c^2$ .

- Significance of the  $K_S^0$  decay length has two standard deviations.

- $D_s$  selection: According to the MC study on  $D_s$  reconstruction (BESSIII-DocDB-380), the  $D_s$  candidates fall into the mass window of  $1.90 < m_{D_s} < 2.03\text{GeV}/c^2$  and the corresponding  $M_{rec}$  satisfied  $2.051 < M_{rec} < 2.180\text{GeV}/c^2$  are retained for further study. In which, the definition of  $M_{rec}$  is

$$M_{rec} = \sqrt{(E_{cm} - \sqrt{p_{D_s}^2 + m_{D_s}^2})^2 - |\vec{p}_{cm} - \vec{p}_{D_s}|^2}, \quad (1)$$

where  $E_{cm}$  is the energy of initial state, calculated from the beam energy [3],  $\vec{p}_{D_s}$  is the momentum of  $D_s$  candidate,  $m_{D_s}$  is  $D_s$  mass quoted from PDG [4] and  $\vec{p}_{cm}$  and  $\vec{p}_{D_s}$  are four-momentum of initial state and the decay products of a  $D_s$  candidate, respectively.

### 3.2 Sigal Selection

We use different methods to select events in model indepent analysis(Sec.4.1), amplitude analysis(Set.5.1) and branching fraction measurement(Set.6.1). And the details are in the correspondding section.

## 4 Model Independent Partial Wave Analysis

In the  $K^+K^-$  threshold both  $a_0(980)$  and  $f_0(980)$  can be present, and both resonances have very similar parameters which suffer from large uncertainties. In this section we obtain model-independent information on the  $K^+K^-S$  wave by performing a MIPPPWA(model independent partial wave analysis) in the  $K^+K^-$  threshold region.

### 4.1 Signal Candidate Selection

After the selection in Set.3, we continue to select signals for model independent partial wave analysis. As MIPWA need more data events, we do not require  $D_s^+$  and  $D_s^-$  appear in pairs. Before selecting the best candidate, we vote the candidates with  $\pi^\pm(\pi^0)$  whose momentum is less than  $0.1GeV$  to remove soft  $\pi^\pm(\pi^0)$  from  $D^*$  decays. For every candidates of  $D_s$  decays, all tracks at signal side are added to apply kinematic fitting. Only mass of  $D_s$  from signal side is constrained. Then we select the candidate with minimum  $\chi^2_{1c}$ .

### 4.2 Background Analysis

In order to further suppress the background, the multiple analysis method (MVA) is used. We train MVA methods separately with different sets of variables for the two event categories depending on the  $D_s^+$  origin. Sideband region used below is defined as the region of  $1.90 < M(D_s) < 1.952GeV$  and  $1.985 < M(D_s) < 2.03GeV$ ,  $M(D_s)$  is the invariant mass of  $D_s$ . And the signal region is  $1.952 < M(D_s) < 1.985GeV$ . These two categories of events are selected in a  $M_{rec} - \Delta M$  2D plane shown in Fig.1:

- Cat. #0: Direct  $D_s^+$ . We use the following variables whose distributions for signal and background are shown in Fig.2,

1.  $M_{rec}$ ,
2.  $P_{rest}$ , defined as the total momentum of the tracks and neutrals in the rest of event (not part of the  $D_s^+ \rightarrow K^+K^-\pi^+$  candidate),
3.  $E_\gamma$ , defined as the energy of gamma from  $D_s^*$ .

From Fig.2(generic MC) and Fig.3(data), we can see that the corresponding distributions of these variables of data and generic MC are roughly consistent.

- Cat. #1: Indirect  $D_s^+$ . We use the following variables whose distributions for signal and background are shown in Fig.4,




Figure 1: Two dimensional plane of  $\Delta M \equiv M(D_s^+ \gamma) - M(D_s^+)$  versus  $M_{rec}$  from the simulated  $D_s^+ \rightarrow K^+ K^- \pi^+$  decays. The red(green) dashed lines mark the mass window for the  $D_s^+$  Cat. #0( Cat. #1) around the  $M_{rec}(\Delta M)$  peak.

1.  $\Delta M$ ,

2.  $M'_{rec}$ , defined as  $M'_{rec} = \sqrt{(E_{cm} - \sqrt{p_{D_s \gamma}^2 + m_{D_s^*}^2})^2 - p_{D_s \gamma}^2}$ , with  $p_{D_s \gamma}$  as the momentum of the  $D_s \gamma$  combination,  $m_{D_s^*}$  as the nominal  $D_s^*$  mass,

3.  $N_{tracks}$ , defined as the total number of tracks and neutrals in an event.

From Fig.4(generic MC) and Fig.5(data), we can see that the corresponding distributions of these variables of data and generic MC are roughly consistent.




Figure 2: For event Cat. #0, distributions of MVA variables from simulated signal decays and background events.

As the results shown in Fig.6, this BDTG training and test samples are well matched. For event Cat.

#0 ( Cat. #1 ), the sample with BDTG value larger than 0.33(0.65) is retained for further study.






Figure 3: The distribution(Cat. #0) of these three observables ((a) and (d))  $M_{rec}$ , ((b) and (e))  $P_{rest}$  and ((c) and (f))  $E_\gamma$  for (a), (b) and (c)) Signal and (d), (e) and (f)) Sideband regions from data are shown.

After applying the BDTG requirement, the background shows no obviously peak around the region of  $[1.95, 1.982] \text{ GeV}$  (Signal region), which are shown in Fig.7. The fit to the signal  $D_s$  invariant mass ( $M_{D_s^2}$ ) spectrum gives the background yield in Signal region is  $73.6 \pm 18.7$ , shown as in Fig.8. In the fit, the signal shape is the MC shape convoluted with a Gaussian function and the background is described with 1<sup>st</sup>-order Chebychev polynomial.

The projections of the "Sideband" ( $1.90 < M(D_s) < 1.95 \text{ GeV}$  and  $1.985 < M(D_s) < 2.03 \text{ GeV}$ ) from data and generic MC after signal events removed are shown in Fig.9. The corresponding plots agree well.

Thus the generic MC sample with signal events removed is used to subtract the background in MIPWA.

### 4.3 MIPWA method

Let  $N$  be the number of events for a given mass interval  $I = [m_{K^+K^-}; m_{K^+K^-} + dm_{K^+K^-}]$ . We write the corresponding angular distributions in terms of the appropriate spherical harmonic functions as

$$\frac{dN}{d\cos\theta} = 2\pi \sum_{k=0}^L \langle Y_k^0 \rangle Y_k^0(\cos\theta), \quad (2)$$

where  $L - 2\ell_{max}$ , and  $\ell_{max}$  is the maximum orbital angular momentum quantum number required to describe the  $K^+K^-$  system at  $m_{K^+K^-}$  (e.g.  $\ell_{max}=1$  for S-, P-wave description);  $\theta$  is the angle between the




Figure 4: For event Cat. #1, distributions of MVA variables from simulated signal decays and background events.




Figure 5: The distribution (Cat. #1) of these three observables ((a) and (d))  $M_{rec}$ , ((b) and (e))  $P_{rest}$  and ((c) and (f))  $E_\gamma$  for ( (a), (b) and (c)) Signal and ( (d), (e) and (f)) Sideband regions from data are shown.

- 1  $K^+$  direction in the  $K^+K^-$  system in the  $D_s^+$  rest frame. The normalizations are such that

$$\int_{-1}^1 Y_k^0(\cos \theta) Y_j^0(\cos \theta) d \cos \theta = \frac{\delta_{kj}}{2\pi}, \quad (3)$$

- 2 and it is assumed that the distribution  $\frac{dN}{d \cos \theta}$  has been efficiency corrected and background subtracted.
- 3 Using this orthogonality condition, the coefficients in the expansion are obtained from

$$\langle Y_k^0 \rangle = \int_{-1}^1 \frac{dN}{d \cos \theta} d \cos \theta, \quad (4)$$

- 4 where the intergral is given, to a good approximation, by  $\sum_{n=1}^N Y_k^0(\cos \theta_n)$ , where  $\theta_n$  is the value of  $\theta$  for
- 5 the  $n$ -th event.




Figure 6: The comparisons between the training and test samples.




Figure 7: The signal and background distributions from generic MC after BDTG requirement.

Fig.10 shows the  $K^+K^-$  mass spectrum up to 1.15 GeV weighted by  $Y_k^0(\cos\theta) = \sqrt{(2k+1)/(4\pi)}P_k(\cos\theta)$  for  $k=0, 1$ , and  $2$ , where  $P_k$  is the Legendre polynomial function of order  $k$ . These distributions are corrected for efficiency and phase space, and background is subtracted using background from generic MC after BDTG requirement.

The number of events  $N$  for the mass interval  $I$  can be expressed also in terms of the partial-wave amplitudes describing the  $K^+K^-$  system. Assuming that only S- and P-wave amplitudes are necessary in this limited region, we can write:

$$\frac{dN}{d\cos\theta} = 2\pi \left| S Y_0^0(\cos\theta) + P Y_1^0(\cos\theta) \right|^2. \quad (5)$$

By comparing Eqa. 2 and 5, we obtain

$$\begin{aligned} \sqrt{4\pi} \langle Y_0^0 \rangle &= |S|^2 + |P|^2, \\ \sqrt{4\pi} \langle Y_2^0 \rangle &= \frac{2}{\sqrt{5}} |P|^2, \end{aligned} \quad (6)$$

The above system of equations can be solved in each interval of  $K^+K^-$  invariant mass for  $|S|$  and  $|P|$  and the resulting distributions are shown in Fig.11.




Figure 8: The fit to the signal  $D_s$  invariant mass ( $M_{D_s}$ ) spectrum after BDTG requirement, the area between the pink arrows is the signal area of the sample for MIPWA.

#### 4.4 S-wave parameterization at the $K^+K^-$ threshold

We empirically parameterize the  $f_0(980)$  with the following function:

$$A_{f_0(980)/a_0(980)} = \frac{1}{m_0^2 - m^2 - im_0\Gamma_0\rho_{KK}}, \quad (7)$$

where  $\rho_{KK} = 2p/m$ , and obtain the following parameter values:

$$\begin{aligned} m_0 &= (0.919 \pm 0.006_{stat}) GeV, \\ \Gamma_0 &= (0.272 \pm 0.040_{stat}) GeV. \end{aligned} \quad (8)$$

The errors are statistical only. The fit result are shown in Fig.12.




Figure 9: The projections of  $m(K^+K^-)$ ,  $m(K^-\pi^+)$ ,  $m(K^+\pi^+)$  from "Sideband" for data(dots with error bars) and generic MC after signal events removed (red histogram) after BDTG requirement.




Figure 10:  $K^+K^-$  mass spectrum in the threshold region weighted by (a)  $Y_0^0$ , (b)  $Y_1^0$  and (c)  $Y_2^0$ , corrected for efficiency and phase space, and background subtracted.




Figure 11: Squared (a) S- and (b) P-wave amplitudes




Figure 12: Fit of squared S-wave amplitudes. the curves result from the fit described in the text.

## 5 Amplitude Analysis

### 5.1 Event Selection for Amplitude Analysis

After  $K^\pm$ ,  $K_S^0$ ,  $\pi^\pm$  and  $\gamma$  are identified in Sec.3, hadronic  $D_s$  decays can be reconstructed with the DTag package. 8 tag modes are used:

$$D_s^- \rightarrow K^+ K^- \pi^-, D_s^- \rightarrow K_S^0 K^-, D_s^- \rightarrow K_S^0 K^- \pi^+ \pi^-, D_s^- \rightarrow K^- \pi^+ \pi^-, D_s^- \rightarrow K_S^0 K^+ \pi^- \pi^-, D_s^- \rightarrow \pi^+ \pi^- \pi^-, D_s^- \rightarrow \eta_{\pi^+ \pi^- \eta_{\gamma\gamma}}', D_s^- \rightarrow K^+ K^- \pi^- \pi^0.$$

With the tagged  $D_s$  meson, the signal  $D_s$  is reconstructed with the remaining good tracks and good showers. The momentum of  $\pi^\pm(\pi^0)$  is required to be larger than  $100 MeV/c^2$  to suppress the background of  $D^* \rightarrow D\pi$ . Only the  $D_s$  candidate with invariant mass falls into  $[1.87, 2.06] GeV/c^2$  are selected.

For every candidate of  $D_s$  decays, all tracks at signal side and tag side as well as gamma from  $D_s^*$  are added to apply kinematic fitting. 5 constrains are added in kinematic fitting: four-momentum of  $D_s D_s^*$ , mass of  $D_s^*$ . Then we select the candidate with minimum  $\chi_{5c}^2$ .

The candidates satisfy:

- $m_{sig}$  and  $m_{tag}$  falls in the mass regions shown in Tab.9.
- $\chi_{5c}^2 < 200$

are retained for amplitude analysis, where  $m_{sig}$  and  $m_{tag}$  refer to mass of  $D_s$  at signal side and tag side respectively.

### 5.2 Background Analysis

We use generic MC to estimate the background. The background and signal shape of generic MC is shown in Fig.13. According to the luminosities of the data and the generic MC, after scaling the background sample to the data size, the background yields in Signal region is 17.2. Then the fit to the signal  $D_s$  invariant mass ( $m_{sig}$ ) spectrum gives the background yield in Signal region is  $18.1 \pm 5.1$ , shown as in Fig.14. The background level in MC is then consistent with the data. In the fit, the signal shape is the MC shape convoluted with a Gaussian function and the background is the MC shape.

### 5.3 Fit Method

The method used in amplitude analysis is the same as the Ref. [6]. In this section, we briefly review the amplitude analysis method used in this analysis.

The relative magnitudes and phases of the partial waves and the mass and width of intermediate resonances are determined by a maximum-likelihood fit to the data selected. The formulas are constructed with covariant tensors [7].





Figure 13: The background and signal shape of generic MC(round 01-40)

Figure 14: The fit to  $m_{sig}$  for data after selections, the area between the purple arrows is the signal area of the sample for the amplitude analysis.

- 1 Since there are three final state particles, only one possible resonant state is allowed in any interme-  
 2 diate process. Thus the amplitude of the  $n^{th}$  intermediate state ( $A_n$ ) is,

$$A_n = P_n S_n F_n^r F_n^D, \quad (9)$$

- 3 where  $S_n$  and  $F_n^{r(D)}$  are the spin factor and the Blatt-Weisskopf barriers of the intermediate state (the  $D_s$   
 4 meson), respectively.  $P_n$  is the propagator of the intermediate resonance. For a decay process  $a \rightarrow bc$ ,  
 5  $s_{a/b/c}$  is denoted to be the invariant mass square of the particle a/b/c, and  $q$  is denoted as the magnitude  
 6 of the momentum of daughter particle in the rest system of a

$$q = \sqrt{\frac{(s_a + s_b + s_c)^2}{4s_a} - s_b}. \quad (10)$$



Then for  $K^*(892)^0$ ,  $\Phi(1020)^0$ ,  $K_0^*(1430)^0$  and  $f_0(1710)$ , they are parameterized as a RBW,

$$P = \frac{1}{(m_0^2 - s_a) - im_0\Gamma(m)},$$

$$\Gamma(m) = \Gamma_0 \left(\frac{q}{q_0}\right)^3 \left(\frac{m_0}{m}\right) (F^r)^2, \quad (11)$$

where  $m_0$  and  $\Gamma_0$  are the mass and the width of the intermediate resonances, and are fixed to the PDG values [8] except the mass and the width of  $f_0(1370)$ . The mass and width of  $f_0(1370)$  are fixed to Ref. [9].  $K_0^*(1430)^0$  is parameterized with Flatté formula:

$$P_{K_0^*(1430)^0} = \frac{1}{M^2 - s - i(g_1\rho_{K\pi}(s) + g_2\rho_{\eta'K}(s))}, \quad (12)$$

where  $s$  is the  $K^-\pi^+$  invariant mass squared,  $\rho_{K\pi}(s)$  and  $\rho_{\eta'K}(s)$  are Lorentz invariant PHSP factor, and  $g_{1,2}$  are coupling constants to the corresponding final state. The parameters of  $K_0^*(1430)^0$  are fixed to values measured by CLEO [10]. For resonance  $f_0(980)$  &  $a_0(980)$ , the RBW form Eq.7 is used to describe the propagator and the values of parameters are fixed to the values in Eq.8.

The Blatt-Weisskopf barriers are given by

$$F_n = 1, (S\text{ wave}),$$

$$F_n = \sqrt{\frac{z_0^2 + 1}{z^2 + 1}}, (P\text{ wave}), \quad (13)$$

$$F_n = \sqrt{\frac{z_0^4 + 3z_0^2 + 9}{z^4 + 3z^2 + 9}}, (D\text{ wave}),$$

where  $z$  and  $z_0$  are  $qR$  and  $q_0R$ , respectively. The  $R$  is the effective radius of the intermediate state or  $D_s$  meson and set to  $3.0\text{GeV}^{-1}$  for intermediate states and  $5.0\text{GeV}^{-1}$  for  $D_s$  meson [6], respectively.

As the limit of the phase space, we only consider the states with angular momenta no more than 2.

As discussed in [11], the spin factor is,

$$S_n = 1, (S\text{ wave}),$$

$$S_n = \tilde{T}^{(1)\mu}(D_s)\tilde{t}_\mu^{(1)}(R), (P\text{ wave}), \quad (14)$$

$$S_n = \tilde{T}^{(2)\mu}(D_s)\tilde{t}_\mu^{(2)}(R), (D\text{ wave}),$$

where the  $\tilde{T}^{(1)\mu}(D_s)(\tilde{T}^{(2)\mu}(D_s))$  and  $\tilde{t}_\mu^{(1)}(R)(\tilde{t}_\mu^{(2)}(R))$  are the same defined in Ref. [7].

The total amplitude  $M$  is then the coherent sum of the amplitudes of intermediate processes,  $M = \sum c_n A_n$ , where  $c_n = \rho_n e^{i\phi_n}$ , is the corresponding complex coefficient. The magnitude  $\rho_n$  and phase  $\phi_n$  are determined by the amplitude analysis. The signal PDF  $f_S(p_j)$  is given by

$$f_S(p_j) = \frac{\epsilon(p_j) |M(p_j)|^2 R_3(p_j)}{\int \epsilon(p_j) |M(p_j)|^2 R_3(p_j) dp_j}, \quad (15)$$

where  $\epsilon(p_j)$  is the detection efficiency parameterized in terms of the final four-momenta  $p_j$ . The index  $j$  refers to the different particles in the final states.  $R_3(p_j)dp_j$  is the standard element of the three-body

1 phase space. The normalization integral is determined by a MC technique,

$$\int \epsilon(p_j) |M(p_j)|^2 R_3(p_j) dp_j = \frac{1}{N_{MC}} \sum_{k_{MC}}^{N_{MC}} \frac{|M(p_j^{k_{MC}})|^2}{|M^{gen}(p_j^{k_{MC}})|^2}, \quad (16)$$

2 where  $k_{MC}$  is the index of the  $k_{MC}^{th}$  event of the MC sample and  $N_{MC}$  is the number of the selected MC  
 3 events. In this analysis, a signal MC sample with about 2 million events is used in the normalization  
 4 intergral calculation.  $M^{gen}(p_j)$  is the PDF function used to generate the MC samples in MC integration.  
 5 Firstly, the PHSP MC are used in MC integration.  $M^{gen}(p_j)$  is a constant overall the phase space. Then  
 6 with result obtained from the fit to data, the signal MC is then generated and used in MC integration. In  
 7 the numerator of Eq. 15,  $\epsilon(p_j)$  is independent of the fitted variables, so it is regarded as a constant term in  
 8 the fit.

9 Since there is only about 0.4% background in the data sample, the contribution from the background  
 10 is ignored in the likelihood calculation:

$$\ln L = \sum_k^{N_{data}} \ln f_S(p_j^k), \quad (17)$$

11 where  $N_{data}$  is the number of candidate events in data.

## 12 5.4 Fit Fraction

13 The fit fractions of the individual amplitudes are calculated according to the fit results and are com-  
 14 pared to the other measurements. In the calculation, a phase space(PHSP) MC with neither detector  
 15 acceptance nor resolution involved is used. The fit fraction for an amplitude is defined as

$$FF(n) = \frac{\sum_{k=1}^{N_{gen}} |A_n|^2}{\sum_{k=1}^{N_{gen}} |M(p_j^k)|^2}, \quad (18)$$

16 where  $N_{gen} = 2000000$ , is the number of the PHSP MC events in this analysis.

17 To estimate the statistical uncertainties of the fit fractions, we repeat the calculation of fit fractions  
 18 by randomly varying the fitted parameters according to the error matrix. Then, for each amplitude, we  
 19 fit the resulting distribution with a Gaussian function, whose width gives the corresponding statistical  
 20 uncertainty.

## 21 5.5 Fit Result

22 The Dalitz plot of  $m^2(K^+ K^-)$  versus  $m^2(K^- \pi^+)$  is shown in Fig. 15. In the plot, we can see a clear  
 23 peak of  $K^*(892)^0$  and  $\phi(1020)^0$ . In the fit, the magnitude and phase of the amplitude  $D_s^+ \rightarrow K^*(892)^0 K^+$   
 24 is fixed to 1.0 and 0.0, and the magnitudes and phases of the amplitudes are allowed to float.




Figure 15: The plot of  $m^2(K^+ K^-)$  versus  $m^2(K^- \pi^+)$  after event selection.

With the requiring the statistical significance larger than 5 standard deviations, there are 6 intermediate prcess,  $D_s^+ \rightarrow \bar{K}^*(892)^0 K^+$ ,  $D_s^+ \rightarrow \phi(1020)\pi^+$ ,  $D_s^+ \rightarrow f_0(980)\pi^+/a_0(980)\pi^+$ ,  $D_s^+ \rightarrow \bar{K}_0^*(1430)^0 K^+$ ,  $D_s^+ \rightarrow f_0(1370)\pi^+$ ,  $D_s^+ \rightarrow f_0(1710)\pi^+$  are retained in final result. The statistical significance of the three amplitudes in final result are also checked.

With one amplitude dropped and the fit repeated, compared with the nominal fit the likelihood shift( $\Delta \ln L$ ) and the number of freedom degree shift( $\Delta n_{par}$ ) are then corresponding to the statistical significance. The detail  $\Delta \ln L$ ,  $\Delta n_{par}$ , and the statistical significance for each amplitude are listed in Tab.3.

The detail magnitudes, phases and fit fractions for the six amplitudes are listed in Tab. 4.

The Dalitz plot projections are shown in Fig.16.

The goodness of the nominal fit is  $\chi^2 = 290.1/280 = 1.04$ .

Table 3: The  $\Delta \ln L$ ,  $\Delta n_{par}$ , and the statistical significance for each amplitude

Amplitude	$\Delta \ln L$	$\Delta n_{par}$	Stat. significance
$D_s^+ \rightarrow \bar{K}^*(892)^0 K^+$	1959.3	2	>20
$D_s^+ \rightarrow \phi(1020)\pi^+$	2303.3	2	>20
$D_s^+ \rightarrow f_0(980)\pi^+ / a_0(980)\pi^+$	270.5	2	>20
$D_s^+ \rightarrow \bar{K}_0^*(1430)^0 K^+$	39.4	2	8.6
$D_s^+ \rightarrow f_0(1370)\pi^+$	22.5	2	6.4
$D_s^+ \rightarrow f_0(1710)\pi^+$	44.7	2	9.2

Table 4: The magnitudes, phases and fit fractions for the six amplitudes

Amplitude	Magnitude	Phase	Fit fractions(%)
$D_s^+ \rightarrow \bar{K}^*(892)^0 K^+$	1.0(fixed)	0.0(fixed)	48.3 $\pm$ 0.9
$D_s^+ \rightarrow \phi(1020)\pi^+$	1.09 $\pm$ 0.02	6.22 $\pm$ 0.07	40.5 $\pm$ 0.7
$D_s^+ \rightarrow f_0(980)\pi^+ / a_0(980)\pi^+$	2.88 $\pm$ 0.14	4.77 $\pm$ 0.07	19.3 $\pm$ 1.7
$D_s^+ \rightarrow \bar{K}_0^*(1430)^0 K^+$	1.26 $\pm$ 0.14	2.91 $\pm$ 0.20	3.0 $\pm$ 0.6
$D_s^+ \rightarrow f_0(1710)\pi^+$	0.79 $\pm$ 0.08	1.02 $\pm$ 0.12	1.9 $\pm$ 0.4
$D_s^+ \rightarrow f_0(1370)\pi^+$	0.58 $\pm$ 0.08	0.59 $\pm$ 0.17	1.2 $\pm$ 0.4

## 5.6 Systematic Uncertainties

Systematic uncertainties taken into account:

- I Variation of resonance masses and widths within  $1\sigma$  error.
  - For  $f_0(980)/a_0(980)$ , the mass and width are shifted within errors from Eq.8 in model independent PWA part.
  - For  $f_0(1370)$ , the mass and width are shifted within errors from [9].
  - For other states, errors used here are taken from PDG [8].
- II Variation of the effective radius of Blatt-Weisskopf Barrier within the range  $[1.0, 5.0] \text{ GeV}^{-1}$  for intermediate resonances and  $[3.0, 7.0] \text{ GeV}^{-1}$  for  $D_s$  meson.
- III Fit bias. The possible bias is given by the result from pull distribution check. With the results obtained from the fit, the signal MC samples are generated with the same size of the data. In this analysis, 300 MC samples with the size equaling to data are used to perform the pull distribution check. The results are listed in Tab.5. The corresponding plots are shown in Fig.17, Fig.18 and Fig.19.




Figure 16:  $D_s^+ \rightarrow K^+ K^- \pi^+$ : Dalitz plot projections from the nominal fit. The data are represented by points with error bars, the fit results by the histograms.

- 1 Thus, the detail results of the systematic uncertainties are summarized in Tab.6. The final results of
- 2 the amplitude analysis are then listed in Tabl.8.
- 3 Below is a comparison between BABAR, CLEO-c and this analysis. Our results are roughly consis-
- 4 tent with those of BABAR and CLEO-c. And for the fit fraction of  $D_s^+ \rightarrow f_0(980)\pi^+/a_0(980)\pi^+$ , we tend
- 5 to agree the result of BABAR.

Table 5: The results of pull distribution checks for the magnitudes, phases and fit fractions for the amplitudes

Amplitude	Phase		Magnitude		Fit fraction	
	mean	width	mean	width	mean	width
$D_s^+ \rightarrow \bar{K}^*(892)^0 K^+$					$-0.13 \pm 0.04$	$0.98 \pm 0.03$
$D_s^+ \rightarrow \phi(1020)\pi^+$	$-0.04 \pm 0.05$	$1.00 \pm 0.03$	$0.07 \pm 0.04$	$0.95 \pm 0.03$	$0.01 \pm 0.04$	$0.95 \pm 0.03$
$D_s^+ \rightarrow f_0(980)\pi^+ / a_0(980)\pi^+$	$-0.07 \pm 0.05$	$1.01 \pm 0.03$	$0.07 \pm 0.05$	$1.10 \pm 0.04$	$0.02 \pm 0.05$	$1.14 \pm 0.04$
$D_s^+ \rightarrow \bar{K}_0^*(1430)^0 K^+$	$0.00 \pm 0.05$	$1.11 \pm 0.04$	$0.14 \pm 0.04$	$0.95 \pm 0.03$	$0.10 \pm 0.04$	$0.99 \pm 0.03$
$D_s^+ \rightarrow f_0(1710)\pi^+$	$0.00 \pm 0.04$	$0.98 \pm 0.03$	$0.08 \pm 0.04$	$0.97 \pm 0.03$	$0.01 \pm 0.04$	$0.99 \pm 0.03$
$D_s^+ \rightarrow f_0(1370)\pi^+$	$-0.11 \pm 0.05$	$1.10 \pm 0.04$	$0.21 \pm 0.04$	$0.99 \pm 0.03$	$0.15 \pm 0.04$	$0.98 \pm 0.03$

Table 6: Systematic uncertainties on the  $\phi$  and FFs for different amplitudes in units of the corresponding statistical uncertainties.

Amplitude		Source			
		I	II	III	Total
$D_s^+ \rightarrow \bar{K}^*(892)^0 K^+$	FF	0.32	0.29	0.13	0.46
$D_s^+ \rightarrow \phi(1020)\pi^+$	$\phi$	0.49	0.10	0.06	0.51
	$\rho$	0.49	0.14	0.18	0.54
	FF	0.44	1.13	0.05	1.21
$D_s^+ \rightarrow f_0(980)\pi^+ / a_0(980)\pi^+$	$\phi$	0.98	0.25	0.06	1.01
	$\rho$	1.11	0.17	0.10	1.13
	FF	1.16	0.15	0.06	1.17
$D_s^+ \rightarrow \bar{K}_0^*(1430)^0 K^+$	$\phi$	1.02	0.48	0.09	1.13
	$\rho$	1.00	0.36	0.13	1.07
	FF	0.76	0.35	0.11	0.85
$D_s^+ \rightarrow f_0(1710)\pi^+$	$\phi$	0.31	0.25	0.09	0.41
	$\rho$	1.17	1.23	0.07	1.70
	FF	0.71	1.21	0.05	1.41
$D_s^+ \rightarrow f_0(1370)\pi^+$	$\phi$	2.66	0.27	0.11	2.67
	$\rho$	1.01	0.32	0.06	1.06
	FF	0.42	0.30	0.15	0.54

Table 7: The final results of the magnitudes, phases and fit fractions for the six amplitudes. The first and second uncertainties are the statistical and systematic uncertainties, respectively.

Amplitude	Magnitude	Phase	Fit fractions(%)
$D_s^+ \rightarrow \bar{K}^*(892)^0 K^+$	1.0(fixed)	0.0(fixed)	$48.3 \pm 0.9 \pm 0.4$
$D_s^+ \rightarrow \phi(1020)\pi^+$	$1.09 \pm 0.02 \pm 0.01$	$6.22 \pm 0.07 \pm 0.04$	$40.5 \pm 0.7 \pm 0.9$
$D_s^+ \rightarrow f_0(980)\pi^+ / a_0(980)\pi^+$	$2.88 \pm 0.14 \pm 0.16$	$4.77 \pm 0.07 \pm 0.07$	$19.3 \pm 1.7 \pm 2.0$
$D_s^+ \rightarrow \bar{K}_0^*(1430)^0 K^+$	$1.26 \pm 0.14 \pm 0.15$	$2.91 \pm 0.20 \pm 0.23$	$3.0 \pm 0.6 \pm 0.5$
$D_s^+ \rightarrow f_0(1710)\pi^+$	$0.79 \pm 0.08 \pm 0.14$	$1.02 \pm 0.12 \pm 0.05$	$1.9 \pm 0.4 \pm 0.6$
$D_s^+ \rightarrow f_0(1370)\pi^+$	$0.58 \pm 0.08 \pm 0.08$	$0.59 \pm 0.17 \pm 0.45$	$1.2 \pm 0.4 \pm 0.2$




Figure 17: The pull distribution check results for phases of the amplitudes in the nominal fit model.

Table 8: Comparison between BABAR, CLEO-c and this analysis.

Amplitude	BABAR	CLEO-c	This Analysis
$D_s^+ \rightarrow \bar{K}^*(892)^0 K^+$	$47.9 \pm 0.5 \pm 0.5$	$47.4 \pm 1.5 \pm 0.4$	$48.3 \pm 0.9 \pm 0.4$
$D_s^+ \rightarrow \phi(1020)\pi^+$	$41.4 \pm 0.8 \pm 0.5$	$42.2 \pm 1.6 \pm 0.3$	$40.5 \pm 0.7 \pm 0.9$
$D_s^+ \rightarrow f_0(980)\pi^+ / a_0(980)\pi^+$	$16.4 \pm 0.7 \pm 2.0$	$28.2 \pm 1.9 \pm 1.8$	$19.3 \pm 1.7 \pm 2.0$
$D_s^+ \rightarrow \bar{K}_0^*(1430)^0 K^+$	$2.4 \pm 0.3 \pm 1.0$	$3.9 \pm 0.5 \pm 0.5$	$3.0 \pm 0.6 \pm 0.5$
$D_s^+ \rightarrow f_0(1710)\pi^+$	$1.1 \pm 0.1 \pm 0.1$	$3.4 \pm 0.5 \pm 0.3$	$1.9 \pm 0.4 \pm 0.6$
$D_s^+ \rightarrow f_0(1370)\pi^+$	$1.1 \pm 0.1 \pm 0.2$	$4.3 \pm 0.6 \pm 0.5$	$1.2 \pm 0.4 \pm 0.2$
$\sum FF(\%)$	$110.2 \pm 0.6 \pm 2.0$	$129.5 \pm 4.4 \pm 2.0$	$114.2 \pm 1.7 \pm 2.3$
$\chi^2/NDF$	$\frac{2843}{2305-14} = 1.2$	$\frac{178}{117} = 1.5$	$\frac{290}{291-10-1} = 1.04$
Events	$96307 \pm 369$	$14400(\text{purity}85\%)$	$4485(\text{purity}99.7\%)$




Figure 18: The pull distribution check results for magnitudes of the amplitudes in the nominal fit model.




Figure 19: The pull distribution check results for fit fractions of the amplitudes in the nominal fit model.



Table 9: The ST yields( $Y_{ST}$ ) and ST efficiencies( $\epsilon_{ST}$ ). The mass windows use the results in Ref. [5]

Tag mode	Mass window(GeV)	$Y_{ST}$	$\epsilon_{ST}(\%)$
$D_s^- \rightarrow K_S^0 K^-$	[1.948, 1.991]	$958648 \pm 1283$	$48.81 \pm 0.07$
$D_s^- \rightarrow K^+ K^- \pi^-$	[1.900, 2.030]	$4064239 \pm 2905$	$40.20 \pm 0.03$
$D_s^- \rightarrow K^+ K^- \pi^- \pi^0$	[1.947, 1.982]	$1164312 \pm 3430$	$10.56 \pm 0.03$
$D_s^- \rightarrow K_S^0 K^- \pi^+ \pi^-$	[1.958, 1.980]	$233843 \pm 1468$	$19.19 \pm 0.12$
$D_s^- \rightarrow K_S^0 K^+ \pi^- \pi^-$	[1.953, 1.983]	$485855 \pm 1313$	$22.59 \pm 0.06$
$D_s^- \rightarrow \pi^- \pi^- \pi^+$	[1.952, 1.984]	$1159359 \pm 3356$	$56.78 \pm 0.16$
$D_s^- \rightarrow \pi^- \eta'_{\pi^+ \pi^- \pi^0}$	[1.940, 1.996]	$251617 \pm 737$	$20.26 \pm 0.06$
$D_s^- \rightarrow K^- \pi^+ \pi^-$	[1.953, 1.983]	$500079 \pm 2870$	$46.83 \pm 0.27$

## 6 Branching Fraction Measurements

### 6.1 Event Selection for Branching Fraction Measurements

After the selection described in Set.3, we continue to select signals used in this part. We use the same 8 tag modes as in Sec.5.1. In the selection of tag  $D_s$ , for multiple candidates, the best candidate is chosen with  $M_{rec}$  closest to mass of  $D_s^*$  in PDF [8]. To further remove the background associated with the larger number of soft  $\pi^\pm(\pi^0)$  from  $D_s^*$  decays, candidates are voted if the momentum of  $\pi^\pm(\pi^0)$  is less than  $0.1 GeV$ .

The single tag(ST) yields are extracted from the fits to the  $D_s$  invariant mass distributions, as shown in Fig.20. In the fit, the mass windows of the tag modes are set to be the same as the Ref. [5]. The signal shape is modeled as MC shape convoluted with a Gaussian function, while background is parameterized as the 2nd-order Chebychev polynomial. The corresponding ST efficiencies are estimated from generic MC. The ST yields( $Y_{ST}$ ) and ST efficiencies( $\epsilon_{ST}$ ) are listed in Tab.9

After selected the candidates of tag  $D_s$ , for each tag mode, we can get double tag candidates.

With the amplitude analysis results obtained from the fit to data, the double efficiencies without any kinematic fit requirement applied are determined and listed in Tab.10.

### 6.2 Analytic Strategy of Branching Fraction Measurement

The data sample for this analysis is collected at  $E_{cm} = 4.178 GeV$ , about  $100 MeV$  higher than  $D_s^* D_s$  threshold. Around this energy region, based on the cross section measurement by CLEO [2], we know that most  $D_s$  production in  $e^+ e^-$  collision comes from  $D_s^{*\pm} D_s^\mp$  events with its cross section approximatedly  $0.9 nb$ , while the cross section for  $D_s^+ D_s^-$  is about a factor of 20 smaller. The  $D_s^*$  decays to either  $\gamma D_s$  or  $\pi^0 D_s$  with branching fractions of  $(93.5 \pm 0.7)\%$  and  $(5.8 \pm 0.7)\%$  [8], respectively. The other charm




Figure 20:  $D_s$  Mass fits from data. The points with error bars are data, and the blue line is the fit. Red short-dashed lines are signal, violet long-dashed lines are background. The red arrows denote the signal region.

Table 10: The DT efficiencies( $\epsilon_{DT}$ ).

Tag mode	$\epsilon_{DT}(\%)$
$D_s^- \rightarrow K_S^0 K^-$	$19.61 \pm 0.14$
$D_s^- \rightarrow K^+ K^- \pi^-$	$18.26 \pm 0.06$
$D_s^- \rightarrow K^+ K^- \pi^- \pi^0$	$4.68 \pm 0.03$
$D_s^- \rightarrow K_S^0 K^- \pi^+ \pi^-$	$8.28 \pm 0.11$
$D_s^- \rightarrow K_S^0 K^+ \pi^- \pi^-$	$9.52 \pm 0.09$
$D_s^- \rightarrow \pi^- \pi^- \pi^+$	$23.55 \pm 0.15$
$D_s^- \rightarrow \pi^- \eta'_{\pi^+ \pi^- \eta_{\gamma\gamma}}$	$8.28 \pm 0.11$
$D_s^- \rightarrow K^- \pi^+ \pi^-$	$20.00 \pm 0.19$

productions total  $7pb$  mainly including  $D^* \bar{D}^*$  with a cross section of  $5nb$ ,  $D^* \bar{D} + D \bar{D}^*$  with a cross section of  $2nb$ , and  $D \bar{D}$  with a cross section of relatively small  $0.2nb$ . There also appears to be  $D \bar{D}^* \pi$  production. The underlying light quark "continuum" background is about  $12nb$ . The relatively large cross sections, relatively large branching fractions, and sufficient luminosities allow us to employ double tag (DT) technique to study this study.

As  $D_s^- \rightarrow K^+ K^- \pi^-$  is not only our signal mode but also one of our tag modes, we divide the events into two categories:

- Cat. A: Tag  $D_s$  decays to tag modes except  $D_s^- \rightarrow K^+ K^- \pi^-$ . The generic MC sample with the signal removed shows no peaking background around the fit range of  $1.90 < M_{sig} < 2.03 GeV$ . Thus, the double tag yield is determined by the fit to  $M_{sig}$ , which has shown in Fig.21(a). And the background is described with  $2^{nd}$ -order Chebychev polynomial. The double tag yield is  $3503 \pm 64$ .
- Cat. B: Tag  $D_s$  decays  $K^+ K^- \pi^+$ . As both of the two  $D_s$  mesons decay to our signal modes, we fit  $aM$  (the average mass of  $D_s$  at signal side and tag side), which is shown in Fig.21(b). Here, the background is described with  $2^{nd}$ -order Chebychev polynomial. The double tag yield is  $1629 \pm 43$ .

To measure the branching fraction of this decay, we start from the following equations with one tag mode:

$$N_{tag}^{obs} = 2N_{D_s^+ D_s^-} \mathcal{B}_{tag} \epsilon_{tag}, \quad (19)$$

$$\begin{aligned} N_{sig}^{obsA} &= 2N_{D_s^+ D_s^-} \mathcal{B}_{tag} \mathcal{B}_{sig} \epsilon_{tag, sig}, & \text{for Cat. A} \\ N_{sig}^{obsB} &= N_{D_s^+ D_s^-} \mathcal{B}_{tag} \mathcal{B}_{sig} \epsilon_{tag, sig}, & \text{for Cat. B} \end{aligned} \quad (20)$$

where  $N_{D_s^+ D_s^-}$  is the total number of  $D_s^{*\pm} D_s^\mp$  produced from  $e^+ e^-$  collision;  $N_{tag}^{obs}$  is the number of observed tag modes;  $N_{sig}^{obsA}$  and  $N_{sig}^{obsB}$  are the number of observed signals for Cat. A and Cat. B, respectively;




Figure 21: Fit of (a)Cat. A and (b)Cat. B. We fit  $M_{sig}$  and  $aM$  for Cat. A and Cat. B, respectively. The signal shapes are the corresponding simulated shapes convoluted with a Gaussian function and the background shapes are described with  $2^{nd}$ -order Chebychev polynomial.

$\mathcal{B}_{tag}$  and  $\mathcal{B}_{sig}$  are the branching fractions of specific tag mode and signal mode, respectively;  $\epsilon_{tag}$  is the reconstruction efficiency of the tag mode;  $\epsilon_{tag,sig}$  is the reconstruction efficiency of both the tag and signal decay modes.

Using the above equations, it's easy to obtain:

$$\mathcal{B}_{sig} = \frac{N_{sig}^{obsA} + 2N_{sig}^{obsB}}{\sum_{\alpha} N_{tag}^{\alpha} \epsilon_{tag,sig}^{\alpha} / \epsilon_{tag}^{\alpha}}, \quad (21)$$

where the yields  $N_{tag}^{obsA}$ ,  $N_{tag}^{obsB}$  and  $N_{tag}^{\alpha}$  can be obtained from data, while  $\epsilon_{tag}$  and  $\epsilon_{tag,sig}$  can be obtained from the appropriate Monte-Carlo(MC) samples,  $\alpha = 1, 2, 3 \dots$  represents the tag mode.

### 6.3 Results of Branching Fraction

Then we can get the branching fraction  $\mathcal{B}(D_s^+ \rightarrow K^+ K^- \pi^+) = (5.38 \pm 0.07)\%$  (only statistical uncertainty) according to Eq.21.

### 6.4 Systematic Uncertainties

The following sources are taken in account to calculate systematic uncertainties.

- Signal shape. The systematic uncertainty due to the signal shape is studied with the fit without the Gaussian function convoluted, the double tag yield shift is taken as the related effect.
- Background shape and fit range. For background shape and the fit range, the MC shape is used to instead the  $1^{st}$ -order Chebychev polynomial and the fit range of  $[1.90, 2.03] GeV$  is changed to  $[1.89, 2.04] GeV$ . The largest branching fraction shift is taken as the related effect.

Table 11: Systematic uncertainties of branching fraction.

Source	Sys. Uncertainty
Signal shape	0.3
Background shape and fit range	1.1
Fit bias	0.2
$K^\pm$ PID efficiency	1.1
$\pi^\pm$ PID efficiency	0.4
$K^\pm$ Tracking efficiency	1.1
$\pi^\pm$ Tracking efficiency	0.2
MC statistics	0.2
total	2.0

- Fit bias. The possible bias is estimated by the IO check with using the round 30-40 of generic MC.
- $K^\pm$  and  $\pi^\pm$  Tracking/PID efficiency. Based on the works [12] and [13] by Xingyu Shan and Sanqiang Qu, etc, we find that it's enough to assign 1.1% , 0.4%, 1.1% and 0.2% as the systematic uncertainty for  $K^\pm$  PID,  $\pi^\pm$  PID,  $K^\pm$  tracking,  $\pi^\pm$  tracking efficiencies, respectively.
- MC statistics. The uncertainty of MC statistics is obtained by  $\sqrt{\sum_i f_i \frac{\delta \epsilon_i}{\epsilon_i}}$ , where  $f_i$  is the tag yield fraction and  $\epsilon_i$  is the signal efficiency of tag mode  $i$ .

All of the systematic uncertainties mentioned above are summarized in Tab.11.

The branching fraction with systematic uncertainties is  $\mathcal{B}(D_s^+ \rightarrow K^+ K^- \pi^+) = (5.38 \pm 0.07_{stat.} \pm 0.11_{sys.})\%$ .

Table 12: The result of this analysis.

Amplitude	Magnitude	Phase	Fit fractions(%)
$D_s^+ \rightarrow \bar{K}^*(892)^0 K^+$	1.0(fixed)	0.0(fixed)	$48.3 \pm 0.9 \pm 0.4$
$D_s^+ \rightarrow \phi(1020)\pi^+$	$1.09 \pm 0.02 \pm 0.01$	$6.22 \pm 0.07 \pm 0.04$	$40.5 \pm 0.7 \pm 0.9$
$D_s^+ \rightarrow f_0(980)\pi^+ / a_0(980)\pi^+$	$2.88 \pm 0.14 \pm 0.16$	$4.77 \pm 0.07 \pm 0.07$	$19.3 \pm 1.7 \pm 2.0$
$D_s^+ \rightarrow \bar{K}_0^*(1430)^0 K^+$	$1.26 \pm 0.14 \pm 0.15$	$2.91 \pm 0.20 \pm 0.23$	$3.0 \pm 0.6 \pm 0.5$
$D_s^+ \rightarrow f_0(1710)\pi^+$	$0.79 \pm 0.08 \pm 0.14$	$1.02 \pm 0.12 \pm 0.05$	$1.9 \pm 0.4 \pm 0.6$
$D_s^+ \rightarrow f_0(1370)\pi^+$	$0.58 \pm 0.08 \pm 0.08$	$0.59 \pm 0.17 \pm 0.45$	$1.2 \pm 0.4 \pm 0.2$

## 7 Summary

This analysis presents the amplitude analysis for the decay  $D_s^+ \rightarrow K^+ K^- \pi^+$ . The results are listed in Tab12. And for the fit fraction of  $D_s^+ \rightarrow f_0(980)\pi^+ / a_0(980)\pi^+$ , we tend to agree the result of BABAR.

We also measured the branching fraction and the value is  $\mathcal{B}(D_s^+ \rightarrow K^+ K^- \pi^+) = (5.38 \pm 0.07_{stat.} \pm 0.11_{sys.})\%$ .

## References

- [1] D.Cronin-Hennessy *et al.* (CLEO Collaboration), Phys. Rev. D**80**, 072001 (2009).
- [2] P.del Amo Sanchez *et al.* (BARBAR Collaboration), Phys. Rev. D**83**, 052001 (2011).
- [3] Andy Julin, Hajime Muramatsu and Ron Poling, BESIII DocDB 580-v1.
- [4] C. Patrignani *et al.* (Particle Data Group), Chin. Phys. C **40**,100001 (2016)
- [5] Sifan Zhang and Hailong Ma, BESIII DocDB 630-v35.
- [6] Yu Lu and Liaoyuan Dong, BESIII DocDB 416-v30.
- [7] B. S. Zou and D. V. Bugg, Eur. Phys. J. A**16**, 537(2003).
- [8] K. A. Olive *et al.* (Particle Data Group), Chin. Phys. C **40**, 100001 (2016)
- [9] M. Alblikim *et al.* (BESIII Collaboration), Phys. Lett. B**607** 243 (2005).
- [10] G. Bonvicini *et al.* (CLEO Collaboration), Phys. Rev. D**78**, 052001 (2001).
- [11] Yu Lu and Liaoyuan Dong, BESIII DocDB 613-v16.
- [12] <https://indico.ihep.ac.cn/event/8006/contribution/1/material/slides/0.pdf>
- [13] <https://indico.ihep.ac.cn/event/8023/contribution/1/material/slides/0.pdf>

# **1 Appendices**



<sup>1</sup> **A To be supplemented**