

Study of sensitivities of the P2O project

M. M. Kirsanov¹**e-mail:** mikhail.m.kirsanov@gmail.com, A. Sokolov²

¹ Institute for Nuclear Research of the Russian Academy of Sciences,
117312 Moscow, Russia

² Institute for High Energy Physics, Protvino, Russia
(Dated: September 9, 2024)

The sensitivities of the project to some oscillation parameters as a function of power and systematic errors are studied.

I. INTRODUCTION

P2O is the project of neutrino beam from IHEP Protvino to ORCA [1]

II. TECHNICAL NOTES ON THE REPOSITORY

The files for this project, including Papers and Notes, are kept in <https://github.com/kirsanov-protvino/P2O>. In order to work on them one should create account on github. Next step is to create an SSH key. Add this key to your account on the page <https://github.com/settings/keys>. After that you will be able to clone the repository using the command `git clone git@github.com:kirsanov-protvino/P2O.git` and push your changes in it. The command above is shown when you click the green button "Code" on the repository page and switch to SSH instead of HTTPS. To check that your repository clone is "pushable" type inside it `git remote -v`.

III. TECHNICAL NOTES ON THE GLOBES PROGRAM RUNNING

In order to check that the statistics corresponds to the Proposal, the following call was used: `glbGetChannelRatePtr(0, ichannel, GLB_PRE)`. It returns the numbers of events in energy bins, to be summed over the them. It is to be called AFTER the calculation of sensitivities, otherwise some variables are not initialized. It is found that with the normalization factor 40 in the glb file the total number of events corresponds to the one in Figure 7 of the Proposal [1] (20000 $\nu_\mu CC$ events by eye from the figure, after scaling from 90 MW to 450 MW).

In the Globes running the file `p2o.glb` was used. Some parameters there, such as energy resolution of 30%, are taken from the Proposal [1], some from the proposal of KM3NET <https://iopscience.iop.org/article/10.1088/0954-3899/43/8/084001>.

IV. STUDY OF THE EXPERIMENT RESOLUTION

The value of $\sqrt{\chi^2}$ as a function of hypothetical δ for the true values $\delta = \pi/2, \pi, 3\pi/2$ is shown in Fig. 1. The energy resolution here is nominal, 30%, as specified in the Proposal.




FIG. 1. Value of $\sqrt{\chi^2}$ as a function of hypothetical δ for $\delta = \pi/2$ (blue), π (green), $3\pi/2$ (red). Energy resolution is 30%.

The resolution of δ measurement for the true values $\delta = 0, \pi/2, \pi, 3\pi/2$ is shown in Fig. 2. It is determined as the deviation of δ from the minimum of χ^2 at which the latter increases by 1, averaged over the left and right sides.




FIG. 2. Resolution of the δ CP measurement as a function of energy resolution for the true values $\delta = 0$ (black), $\pi/2$ (blue), π (green), $3\pi/2$ (red).

V. EFFECT OF ν_e FRACTION IN THE BEAM

This fraction is specified in the spectra file. To avoid rewriting the spectra files, different fractions were simulated by changing the efficiency with which the channel "nu-e-beam" is added to the backgrounds. This is possible because the program does not have a protection against efficiencies greater than 1. For example the nominal efficiency for the ν_e appearance is 0.85. The fraction of ν_e increased by factor of 10 was simulated by increasing this efficiency to 8.5. This changes the delta CP measurement accuracy from 61.65 to 62.55 degrees (delta CP = $3/2 \pi$). Decreasing the ν_e fraction is not visible if we print out only 3 digits of delta CP measurement.

I understand this small effect like this. All events without identified muon are summed up ("shower type" events) and used in the analysis. In the model described in our glb file we have one error specified for this: 5%. So the effect of increased ν_e fraction remains small if the fraction of ν_e in the total number of "shower type" events does not dominate.

The effect of ν_e fraction can be more significant if we assume that we don't know it with sufficient accuracy and specify this separate accuracy in the glb file. "Sufficient" here means a value of the order of 5% mentioned above.

VI. EFFECT OF THE TYPE OF SYSTEMATIC ERROR

No significant effect was observed when it was changed from "chiSpectrumCalib" to "chiSpectrumTilt". Artefacts in the dependencies also did not change.

-
- [1] A. V. Akindinov *et al.*, Eur. Phys. J. C **79**, 758 (2019), arXiv:1902.06083 [physics.ins-det].