

a
Departures from RNA by protein class
≥0.5 log2
≥1
≥0.5
≥1
Plasma-derived (12)
67%
42%
Secreted (374)
17%
5%
Matrix (154)
17%
3%
Transmembrane (2,332)
4%
0.6%
Intracellular (6,708)
2%
0.5%
20
0
20
40
% of class: protein below ← RNA prediction → above
b
Terminal complement
C5
C6
C8A
RNA low/ND
C8B
C8G
C9
RNA low/ND
Protein (95% CI)
RNA, other rats
0
2
log2 change
(protein 25 vs 9; RNA 27 vs 6 months)
c
Excess over plasma proteins
Plasma proteins (16)
2.59×
Albumin
1.83×
C3
2.15×
0.0
0.5
1.0
1.5
2.0
Terminal-complement excess
over comparator, 25 vs 9 months (log2)
d
UMOD peptide map
Upstream (18)
C-terminal (3)
Boundary (1)
22 sequences
R591
0
200
400
600
Rat UMOD residue (precursor, 644 aa)
e
Upstream region accumulates
Upstream
C-terminal
Region, log2 vs 9 months
2
1
0
9
18
21
25
Age (months)
f
Regions diverge within one precursor
Regions
Upstream
C-terminal
TAL markers (6)
Contrasts
Upstream − C-terminal
q = 0.004
Upstream − TAL
0
1
2
log2 change, 25 vs 9 months
g
Basement-membrane components change selectively
25 / 9
Collagen IV
COL4A1
1.74×
COL4A2
1.93×
COL4A3
2.55×
COL4A4
1.90×
COL4A5
2.60×
Laminins
LAMA5
1.28×
LAMB1
1.26×
LAMB2
1.28×
LAMC1
1.32×
Nidogens
NID1
1.43×
NID2
0.97×
Fibrillar collagens (recovered)
COL1A1
0.95×
COL1A2
0.98×
COL3A1
1.17×
−0.5
0.0
0.5
1.0
1.5
2.0
log2 change, 25 vs 9 months (95% CI)
h
Collagen IV rises without RNA
COL4A3
COL4A5
COL4A2
COL4A1
COL4A4
Laminins
NID1
COL3A1
NID2
COL1A1
Protein log2 change, 25 vs 9 months
1.5
1.0
0.5
0.0
−1
0
1
RNA log2 change, 27 vs 6 months
(54 other rats)