
a
Differentiated tubular programs
Mean log2 vs 9 months
0.0
−0.1
−0.2
−0.3
PT identity
OXPHOS
FAO
Transport
Glycolysis
Ribosome
Proteasome
9
18
21
25
Age (months)
b
Mature segment markers
PT-S1 (13)
PT-S2 (12)
PT-S3 (14)
−0.6
−0.5
−0.4
−0.3
−0.2
−0.1
0.0
0.1
log2 change, 25 vs 9 months
c
Coordination among age-matched kidneys
r = 0.909
OXPHOS residual (log2)
0.4
0.2
0.0
−0.2
−0.4
−0.3
−0.2
−0.1
0.0
0.1
0.2
0.3
PT identity residual (log2)
d
Contralateral NoRI geometry
Nuclear density
Lumen fraction
Profile area
Epithelial protein
PT-profile fraction
−20
0
20
40
60
80
Change as % of observed 9-month mean
e
Original and PT-conditioned differences
OXPHOS
FAO
Transport
Original
With PT score
−0.5
−0.4
−0.3
−0.2
−0.1
0.0
Change, 25 vs 9 months (log2)
f
Specificity control for OXPHOS conditioning
3,626 matched declining sets
Median 1.2%
PT: 25.8%
Accepted matched sets
300
200
100
0
−30
−20
−10
0
10
20
30
Age difference remaining (% of original)