

# Secant Method of Numerical analysis

Last Updated : 19 Aug, 2020

**Secant method** is also a recursive method for finding the root for the polynomials by successive approximation. It's similar to the **Regular-falsi** method but here we don't need to check **f(x<sub>1</sub>)f(x<sub>2</sub>) < 0** again and again after every approximation. In this method, the neighbourhoods roots are approximated by secant line or chord to the function **f(x)**. It's also advantageous of this method that we don't need to differentiate the given function **f(x)**, as we do in **Newton-raphson** method.




Figure – Secant Method

Now, we'll derive the formula for secant method. The equation of Secant line passing through two points is :

$$Y - Y_1 = m(X - X_1)$$

Here, m=slope

So, apply for **(x<sub>1</sub>, f(x<sub>1</sub>))** and **(x<sub>0</sub>, f(x<sub>0</sub>))**

$$Y - f(x_1) = \frac{f(x_0) - f(x_1)}{(x_0 - x_1)} (x - x_1) \quad \text{Equation (1)}$$

As we're finding root of function f(x) so, Y=f(x)=0 in Equation (1) and the point where the secant line cut the x-axis is,

$$x = x_1 - \frac{(x_0 - x_1)}{(f(x_0) - f(x_1))} f(x_1)$$

We use the above result for successive approximation for the root of function f(x). Let's say the first approximation is **x=x<sub>2</sub>**:

$$x_2 = x_1 - \frac{(x_0 - x_1)}{(f(x_0) - f(x_1))} f(x_1)$$

Similarly, the second approximation would be **x=x<sub>3</sub>**:

$$x_3 = x_2 - \frac{(x_1 - x_2)}{(f(x_1) - f(x_2))} f(x_2)$$

And so on, **till k<sup>th</sup> iteration**,

$$x_{k+1} = x_k - \frac{(x_{k-1} - x_k)}{(f(x_{k-1}) - f(x_k))} f(x_k)$$

**Note:** To start the solution of the function f(x) two initial guesses are required such that **f(x<sub>0</sub>) < 0** and **f(x<sub>1</sub>) > 0**. Usually it hasn't been asked to find, that root of the polynomial f(x) at which **f(x) = 0**. Mostly You would only be asked by the problem to find the root of the **f(x)** till two decimal places or three decimal places or four etc.

## Example-1 :

Compute the root of the equation  $x^2 e^{-x/2} = 1$  in the interval [0, 2] using the secant method. The root should be correct to three decimal places.

### Solution -

$x_0 = 1.42$ ,  $x_1 = 1.43$ ,  $f(x_0) = -0.0086$ ,  $f(x_1) = 0.00034$ .

Apply, **secant method**, The first approximation is,

$$x_2 = x_1 - \frac{(x_0 - x_1)}{(f(x_0) - f(x_1))} f(x_1)$$
$$= 1.43 - \frac{(1.42 - 1.43)}{(0.00034 - (-0.0086))} (0.00034)$$
$$= 1.4296$$
$$f(x_2) = -0.000011 \text{ (-ve)}$$

The second approximation is,

$$x_3 = x_2 - \frac{(x_1 - x_2)}{(f(x_1) - f(x_2))} f(x_2)$$
$$= 1.4296 - \frac{(1.42 - 1.4296)}{(0.00034 - (-0.000011))} (-0.000011)$$
$$= 1.4292$$

Since, **x<sub>2</sub>** and **x<sub>3</sub>** matching up to **three decimal places**, the required root is **1.429**.

## Example-2 :

A real root of the equation  $f(x) = x^3 - 5x + 1 = 0$  lies in the interval (0, 1). Perform four iterations of the secant method.

### Solution -

We have,  $x_0 = 0$ ,  $x_1 = 1$ ,  $f(x_0) = 1$ ,  $f(x_1) = -3$

$$x_2 = x_1 - \frac{(x_0 - x_1)}{(f(x_0) - f(x_1))} f(x_1)$$
$$= 1 - \frac{(0 - 1)}{(1 - (-3))} (-3)$$
$$= 0.25$$

$$f(x_2) = -0.234375$$

The second approximation is,

$$x_3 = x_2 - \frac{(x_1 - x_2)}{(f(x_1) - f(x_2))} f(x_2)$$
$$= (-0.234375) - \frac{(1 - 0.25)}{(-3 - (-0.234375))} (-0.234375)$$
$$= 0.186441$$
$$f(x_3) = x_3 - \frac{(x_2 - x_3)}{(f(x_2) - f(x_3))} f(x_3)$$
$$= 0.186441 - \frac{(0.25 - 0.186441)}{(-0.234375 - (-0.074276))} (-0.074276)$$
$$= \mathbf{0.201736}$$

$$f(x_4) = -0.000470$$

The fourth approximation is,

$$x_5 = x_4 - \frac{(x_3 - x_4)}{(f(x_3) - f(x_4))} f(x_4)$$
$$= 0.201736 - \frac{(0.186441 - 0.201736)}{(0.074276 - (-0.000470))} (-0.000470)$$
$$= 0.201640$$

Attention reader! Don't stop learning now. Get hold of all the important CS Theory concepts for SDE interviews with the [CS Theory Course](#) at a student-friendly price and become industry ready.

Like 0

Previous

Global Variables in MS SQL Server

Next

PPP Automaton States

## RECOMMENDED ARTICLES

Page : 1 2

01 Absolute, Relative and Percentage errors in Numerical Analysis  
30, Dec 20

05 Runge-Kutta 2nd order method to solve Differential equations  
03, Apr 20

02 Univariate, Bivariate and Multivariate data and its analysis  
14, Aug 18

06 Multiplicative Congruence method for generating Pseudo Random Numbers  
16, Jul 20

03 Program for Gauss Siedel Method (Computational Mathematics)  
17, Jun 19

07 Additive Congruence method for generating Pseudo Random Numbers  
16, Jul 20

04 Program for Picard's iterative method | Computational Mathematics  
27, Jun 19

08 Linear Congruence method for generating Pseudo Random Numbers  
16, Jul 20

## Article Contributed By :

 **madhav\_mohan**  
@madhav\_mohan

## Vote for difficulty

Easy Normal Medium Hard Expert

Article Tags : Engineering Mathematics


Improve Article

Report Issue

Writing code in comment? Please use [ide.geeksforgeeks.org](https://ide.geeksforgeeks.org), generate link and share the link here.


Load Comments

## WHAT'S NEW




Ad free experience with GeeksforGeeks Premium

[View Details](#)



DSA Self Paced Course

[View Details](#)



DSA Live Classes for Working Professionals

[View Details](#)

## MOST POPULAR IN ENGINEERING MATHEMATICS

Arrow Symbols in LaTeX

Univariate, Bivariate and Multivariate data and its analysis

Difference between Propositional Logic and Predicate Logic

Proof of De-Morgan's laws in boolean algebra

Mathematics | Eigen Values and Eigen Vectors

## MORE RELATED ARTICLES IN ENGINEERING MATHEMATICS

Mathematics | Mean, Variance and Standard Deviation

Mathematics | Set Operations (Set theory)

Inclusion-Exclusion and its various Applications

Mathematics | Planar Graphs and Graph Coloring

Mathematics | Limits, Continuity and Differentiability