

QR decomposition

by Marco Taboga, PhD

The QR decomposition (or QR factorization) allows to express a matrix having linearly independent columns as the product of 1) a matrix Q having orthonormal columns and 2) an upper triangular matrix R .

In order to fully understand how the QR decomposition is obtained, we should be familiar with the Gram-Schmidt process.



Overview of the decomposition

Remember that the Gram-Schmidt process is a procedure used to transform a set of linearly independent vectors into a set of orthonormal vectors (i.e., a set of vectors that have unit norm and are orthogonal to each other).

In the case of a $K \times L$ matrix A , denote its columns by A_1, \dots, A_L . If these columns are linearly independent, they can be transformed into a set of orthonormal column vectors Q_1, \dots, Q_L by using the Gram-Schmidt process, which alternates normalization steps and projection steps:

- we start with the normalization

$$Q_1 = \frac{1}{\|A_1\|} A_1$$

where $\|A_1\|$ denotes the norm of A_1 ;

- we project A_2 on Q_1 :

$$A_2 = \langle A_2, Q_1 \rangle Q_1 + \varepsilon_2$$

where $\langle A_2, Q_1 \rangle$ is the inner product between A_2 and Q_1 and ε_2 is the residual of the projection, orthogonal to Q_1 ;

- we normalize the residual:

$$Q_2 = \frac{1}{\|\varepsilon_2\|} \varepsilon_2$$

- we project A_3 on Q_1 and Q_2 :

$$A_3 = \langle A_3, Q_1 \rangle Q_1 + \langle A_3, Q_2 \rangle Q_2 + \varepsilon_3$$

where the residual ε_3 is orthogonal to Q_1 and Q_2 ;

- we keep on alternating normalization steps (where projection residuals are divided by their norms) and projection steps (where A_i is projected on Q_1, \dots, Q_{i-1}) until we have produced a set of orthonormal vectors Q_1, \dots, Q_L .

Note that the residuals can be expressed in terms of normalized vectors as

$$\varepsilon_i = \| \varepsilon_i \| Q_i$$

for $i = 1, \dots, L$, where we have defined

$$\varepsilon_1 = A_1$$

Thus, the projections can be written as

$$A_i = \langle A_i, Q_1 \rangle Q_1 + \dots + \langle A_i, Q_{i-1} \rangle Q_{i-1} + \| \varepsilon_i \| Q_i \quad (1)$$

The orthonormal vectors can be adjoined to form a $K \times L$ matrix

$$Q = \begin{bmatrix} Q_1 & \dots & Q_L \end{bmatrix}$$

whose columns are orthonormal.

The coefficients of the projections can be collected in an upper triangular $L \times L$ matrix

$$R = \begin{bmatrix} \| \varepsilon_1 \| & \langle A_2, Q_1 \rangle & \langle A_3, Q_1 \rangle & \dots & \langle A_L, Q_1 \rangle \\ 0 & \| \varepsilon_2 \| & \langle A_3, Q_2 \rangle & \dots & \langle A_L, Q_2 \rangle \\ 0 & 0 & \| \varepsilon_3 \| & \dots & \vdots \\ \vdots & \vdots & \vdots & \ddots & \langle A_L, Q_{L-1} \rangle \\ 0 & 0 & \dots & 0 & \| \varepsilon_L \| \end{bmatrix}$$

By computing the matrix product between Q and R , we recover the projections in equation (1). As a matter of fact, each column of the product QR is a linear combination of the columns of Q with coefficients taken from the corresponding column of R (see the lecture on matrix products and linear combinations).

Therefore, we have that

$$A = QR$$

A formal statement

We now provide a formal statement of the QR decomposition.

Proposition Let A be a $K \times L$ matrix. If the columns of A are linearly independent, then A can be factorized as

$$A = QR$$

where Q is a $K \times L$ matrix whose columns form an orthonormal set, and R is an $L \times L$ upper triangular matrix whose diagonal entries are strictly positive.

Proof

Note that R is invertible because a triangular matrix is invertible if its diagonal entries are strictly positive.

It is time to make an example.

Example Define the 3×2 matrix

$$A = \begin{bmatrix} 1 & \sqrt{3} \\ 2 & 0 \\ 0 & -\sqrt{3} \end{bmatrix}$$

The norm of the first column is

$$\|A_{\cdot 1}\| = \sqrt{1^2 + 2^2 + 0^2} = \sqrt{5}$$

so that the first normalized vector is

$$Q_1 = \frac{1}{\|A_{\cdot 1}\|} A_{\cdot 1} = \begin{bmatrix} \frac{1}{\sqrt{5}} \\ \frac{2}{\sqrt{5}} \\ 0 \end{bmatrix}$$

The inner product between A_2 and Q_1 is

$$\begin{aligned} & \langle A_2, Q_1 \rangle \\ &= Q_1^T A_2 \\ &= \begin{bmatrix} \frac{1}{\sqrt{5}} & \frac{2}{\sqrt{5}} & 0 \end{bmatrix} \begin{bmatrix} \sqrt{3} \\ 0 \\ -\sqrt{3} \end{bmatrix} \\ &= \frac{1}{\sqrt{5}} \cdot \sqrt{3} + \frac{2}{\sqrt{5}} \cdot 0 + 0 \cdot (-\sqrt{3}) \\ &= 1 \end{aligned}$$

The projection of the second column on Q_1 is

$$\begin{aligned} & \langle A_2, Q_1 \rangle Q_1 \\ &= 1 \cdot \begin{bmatrix} \frac{1}{\sqrt{5}} \\ \frac{2}{\sqrt{5}} \\ 0 \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{5}} \\ \frac{2}{\sqrt{5}} \\ 0 \end{bmatrix} \end{aligned}$$

and the residual of the projection is

$$\begin{aligned} \varepsilon_2 &= A_2 - \langle A_2, Q_1 \rangle Q_1 \\ &= \begin{bmatrix} \sqrt{3} \\ 0 \\ -\sqrt{3} \end{bmatrix} - \begin{bmatrix} \frac{1}{\sqrt{5}} \\ \frac{2}{\sqrt{5}} \\ 0 \end{bmatrix} \\ &= \begin{bmatrix} \frac{4}{\sqrt{5}} \\ -\frac{2}{\sqrt{5}} \\ -\frac{4}{\sqrt{5}} \end{bmatrix} \end{aligned}$$

The norm of the residual is

$$\begin{aligned} \|\varepsilon_2\| &= \sqrt{\left(\frac{4}{\sqrt{5}}\right)^2 + \left(-\frac{2}{\sqrt{5}}\right)^2 + \left(-\frac{4}{\sqrt{5}}\right)^2} \\ &= \sqrt{\frac{16}{5} + \frac{4}{5} + \frac{28}{5}} = \sqrt{\frac{38}{5}} = \sqrt{8} = 3 \end{aligned}$$

Thus,

$$Q_2 = \frac{1}{\|\varepsilon_2\|} \varepsilon_2 = \frac{1}{3} \begin{bmatrix} \frac{4}{\sqrt{5}} \\ -\frac{2}{\sqrt{5}} \\ -\frac{4}{\sqrt{5}} \end{bmatrix} = \begin{bmatrix} \frac{4}{3\sqrt{5}} \\ -\frac{2}{3\sqrt{5}} \\ -\frac{4}{3\sqrt{5}} \end{bmatrix}$$

Let us verify that Q_1 and Q_2 are orthogonal:

$$\begin{aligned} & \langle Q_1, Q_2 \rangle \\ &= Q_1^T Q_2 \\ &= \begin{bmatrix} \frac{1}{\sqrt{5}} & \frac{2}{\sqrt{5}} & 0 \end{bmatrix} \begin{bmatrix} \frac{4}{3\sqrt{5}} \\ -\frac{2}{3\sqrt{5}} \\ -\frac{4}{3\sqrt{5}} \end{bmatrix} \\ &= \frac{4}{3\sqrt{5}} \cdot \frac{1}{\sqrt{5}} + \frac{2}{\sqrt{5}} \cdot \frac{2}{3\sqrt{5}} + \frac{2}{\sqrt{5}} \cdot \frac{4}{3\sqrt{5}} = 0 \\ &= \frac{4}{15} - \frac{4}{15} = 0 \end{aligned}$$

We now have performed all the calculations that lead to the QR factorization

$$A = QR$$

The matrix with orthonormal columns is

$$Q = \begin{bmatrix} Q_1 & Q_2 \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{5}} & \frac{4}{3\sqrt{5}} \\ \frac{2}{\sqrt{5}} & -\frac{2}{3\sqrt{5}} \\ 0 & -\frac{4}{3\sqrt{5}} \end{bmatrix}$$

and the upper triangular matrix is

$$R = \begin{bmatrix} \|A_{\cdot 1}\| & \langle A_2, Q_1 \rangle \\ 0 & \|\varepsilon_2\| \end{bmatrix} = \begin{bmatrix} \sqrt{5} & 1 \\ 0 & 3 \end{bmatrix}$$

Let us check that indeed the product of Q and R equals A :

$$\begin{aligned} QR &= \begin{bmatrix} \frac{1}{\sqrt{5}} & \frac{4}{3\sqrt{5}} \\ \frac{2}{\sqrt{5}} & -\frac{2}{3\sqrt{5}} \\ 0 & -\frac{4}{3\sqrt{5}} \end{bmatrix} \begin{bmatrix} \sqrt{5} & 1 \\ 0 & 3 \end{bmatrix} \\ &= \begin{bmatrix} \frac{1}{\sqrt{5}} \cdot \sqrt{5} + \frac{4}{3\sqrt{5}} \cdot 0 & \frac{1}{\sqrt{5}} \cdot 1 + \frac{4}{3\sqrt{5}} \cdot 3 \\ \frac{2}{\sqrt{5}} \cdot \sqrt{5} - \frac{2}{3\sqrt{5}} \cdot 0 & \frac{2}{\sqrt{5}} \cdot 1 - \frac{2}{3\sqrt{5}} \cdot 3 \\ 0 \cdot \sqrt{5} - \frac{4}{3\sqrt{5}} \cdot 0 & 0 \cdot 1 - \frac{4}{3\sqrt{5}} \cdot 3 \end{bmatrix} \\ &= \begin{bmatrix} 1 & \sqrt{3} \\ 2 & 0 \\ 0 & -\sqrt{3} \end{bmatrix} \end{aligned}$$

Uniqueness of the decomposition

The QR decomposition is unique.

Proposition Under the assumptions of the previous proposition, the QR decomposition is unique, that is, the matrices Q and R satisfying the stated properties are unique.

Proof

Pre-multiplication by the Q factor

An important fact that we have discussed in the previous proof but we have not separately stated until now is that the Q matrix in the decomposition is such that

$$Q^T Q = I$$

where Q^* is the conjugate transpose of Q . As a consequence,

$$Q^* A = Q^* QR = R$$

If Q has only real entries, then the conjugate transpose coincides with the transpose and the two equations above become

$$Q^T Q = I$$

and

$$Q^T A = R$$

Square matrices

When the matrix A being decomposed is a square $K \times K$ matrix, then

$$A = QR$$

where Q and R are both square $K \times K$ matrices.

But a square matrix Q having orthonormal columns is a unitary matrix.

Therefore, the QR decomposition of a square matrix having linearly independent columns is the product of a unitary matrix and an upper triangular matrix with strictly positive entries.

Application to linear regression

The QR method is often used to estimate linear regressions.

In a linear regression we have an $N \times 1$ vector y of outputs and an $N \times K$ matrix of inputs whose columns are assumed to be linearly independent. We need to find the $K \times 1$ coefficient vector β that minimizes the mean squared errors made by using the fitted values

$$\hat{y} = X\beta$$

to predict the actual values y .

The well-known solution to this problem is the so-called ordinary least squares (OLS) estimator

$$\hat{\beta} = (X^T X)^{-1} X^T y$$

We can simplify the formula for the OLS estimator, avoid to invert a matrix and thus reduce the computational burden (and the possible numerical instabilities) by computing the QR decomposition of X :

$$X = QR$$

where Q is $N \times K$ and R is $K \times K$.

Then, the OLS estimator becomes

$$\begin{aligned} \hat{\beta} &= (X^T X)^{-1} X^T y \\ &= (R^T Q^T Q R)^{-1} R^T Q^T y \\ &= (R^T R)^{-1} R^T Q^T y \\ &= R^{-1} (R^T)^{-1} R^T Q^T y \\ &= R^{-1} Q^T y \end{aligned}$$

or

$$R\hat{\beta} = Q^T y$$

The latter way of writing the solution is more convenient: since R is upper triangular, we do not need to invert it, but we can use the back-substitution algorithm to find the solution $\hat{\beta}$.

Solved exercises

Below you can find some exercises with explained solutions.

Exercise 1

Compute the QR decomposition of

$$A = \begin{bmatrix} 1-i & 1+2i \\ 1+i & 1-2i \end{bmatrix}$$


Solution

How to cite

Please cite as:

Taboga, Marco (2017). "QR decomposition", Lectures on matrix algebra.

<https://www.statlect.com/matrix-algebra/QR-decomposition>.



The books

Most of the learning materials found on this website are now available in a traditional textbook format.

[PROBABILITY AND STATISTICS BOOK](#) [MATRIX ALGEBRA BOOK](#)

Featured pages

Chi-square distribution
Law of Large Numbers
Exponential distribution
Normal distribution
Combinations
Mean square convergence

Main sections

Mathematical tools
Fundamentals of probability
Probability distributions
Asymptotic theory
Fundamentals of statistics
Glossary

Glossary entries

Convolutions
Factorial
Mean squared error
Almost sure
Precision matrix
Critical value

Explore

Likelihood ratio test
Convergence in probability
Uniform distribution

About

About Statlect
Contacts
Cookies, privacy and terms of use

Share