




Contents lists available at SciVerse ScienceDirect

## International Journal of Cardiology

journal homepage: [www.elsevier.com/locate/ijcard](http://www.elsevier.com/locate/ijcard)

Letter to the Editor

## Insight into relation between autonomic function and hypertension

Kirthana Kunikullaya U<sup>\*</sup>, Jaisri Goturu

Department of Physiology, M S Ramaiah Medical College, MSRT Post, MSR Nagar, Bangalore, Karnataka, India – 560054

## ARTICLE INFO

## Article history:

Received 7 August 2011

Accepted 13 August 2011

Available online xxxx

## Keywords:

Cardiac autonomic function

Pre hypertension

Hypertension

## To the Editor,

I was very glad to read Erdogan et al. article [1]. Prehypertension confers higher cardiovascular risk [2] and was included in JNC VII to encourage patients and clinicians to intervene and prevent/delay development of hypertension [3]. Studies have shown beneficial effect of diet and lifestyle modification among prehypertensives [4,5].

This study [1] has used heart rate recovery (HRR) as a simple tool to test autonomic nervous system (ANS) function. Thus, HRR can serve as a good screening test among normotensives and prehypertensives. I commend the authors for this simple but innovative work. Attenuated HRR can predict endothelial dysfunction in coronary artery disease patients [6].

This study [1] also shows that autonomic dysfunction precedes the development of hypertension. There are hardly few studies which have assessed the autonomic function of prehypertensives. Normotensives with reduced heart rate variability (HRV) are at greater risk of developing hypertension [7]. Borderline hypertensives have increased sympathetic and decreased parasympathetic activity that correlates significantly with renin release [8].

A small study population and the lack of a larger control group are the two major limitations of this study as acknowledged by the author [1]. The sample size of each of the groups was not calculated based on any prior published data.

Another limitation of this study [1] is the lack of data on the family history of hypertension among the subjects tested. A landmark study demonstrated decreased parasympathetic activity in normotensive subjects with a positive family history of hypertension [9]. They are

also characterized by altered cardiovascular morphology [10] along with significantly different autonomic function [11].

The study [1] has not looked into the interaction of the autonomic function with the renin–angiotensin system, and other biomarkers. Angiotensin II may contribute to activation of sympathetic activity and suppression of parasympathetic activity [12]. High Plasma Renin Activity (PRA) is an independent determinant of diminished vagal activity [13]. Review on interaction between endothelium and ANS has shown that decreased HRV is associated with endothelial dysfunction [12]. Nitric oxide acts as a sympatholytic agent [14]. Reduced HRV triggers subclinical inflammation [15,16]. Inflammatory markers increase the incidence of prehypertension suggesting that prehypertension might be a pro-inflammatory condition [17]. Oxidative stress impairs endothelial function [18]. Vitamin E, antioxidant, reduces catecholamine levels and improves HRV [19].

Further studies are warranted to unravel the effect of biomarkers on autonomic function and a specific targeted therapy towards them to prevent development of hypertension. Probably integration of various risk factors rather than reliance on a single risk factor will be an efficient approach to implement when screening large populations as a basis for preventive interventions.

The author of this manuscript has certified that she complies with the principles of ethical publishing in the International Journal of Cardiology (Shewan and Coats 2010;144:1–2).

## References

- [1] Erdogan D, Gonul E, Icli A, et al. Effects of normal blood pressure, prehypertension, and hypertension on autonomic nervous system function. *Int Jour Cardiol* 2011;151:50–3.
- [2] Grotto I, Grossman E, Huerta M, et al. Prevalence of prehypertension and associated cardiovascular risk profiles among young Israeli adults. *Hypertension* 2006;48:254–9.
- [3] Chobanian AV, Bakris GL, Black HR, et al. The seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Pressure. The JNC 7 Report. *JAMA* 2003;289:2560–71.
- [4] Márquez-Celedonio FG, Téxon-Fernández O, Chávez-Negrete A, et al. Clinical effect of lifestyle modification on cardiovascular risk in prehypertensives: PREHIPER I study. *Rev Esp Cardiol* 2009;62(1):86–90.
- [5] Wister A, Loewen N, Kennedy-Symonds H, et al. One-year follow-up of a therapeutic lifestyle intervention targeting cardiovascular disease risk. *CMAJ* 2007;177:859–65.
- [6] Huang PH, Leu HB, Chen JW, et al. Usefulness of attenuated heart rate recovery immediately after exercise to predict endothelial dysfunction in patients with suspected coronary artery disease. *Am J Cardiol* 2004;93(1):10–3.
- [7] Schroeder EB, Liao D, Chambless LE, et al. Hypertension, blood pressure, and heart rate variability: The Atherosclerosis Risk in Communities (ARIC) Study. *Hypertension* 2003;42:1106–11.
- [8] Duprez DA, De Sutter JH, De Buyzere ML, et al. Renin–angiotensin–aldosterone system, RR interval, and blood pressure variability during postural changes in borderline arterial hypertension. *Am J Hypertens* 1995;8:683–8.

<sup>\*</sup> Corresponding author. Tel.: +91 9742334950.

E-mail address: [kirthanaku@gmail.com](mailto:kirthanaku@gmail.com) (K. Kunikullaya U).

- [9] Wu JS, Lu FH, Yang YC, et al. Epidemiological study on the effect of prehypertension and family history of hypertension on cardiac autonomic function. *J Am Coll Cardiol* 2008;51:1896–901.
- [10] Muldoon F, Terrel DF, Bunker CH, et al. Family history studies in hypertension research. *Am J Hypertens* 1993;6:76–88.
- [11] Pitzalis MV, Iacoviello M, Massari F, et al. Influence of gender and family history of hypertension on autonomic control of heart rate, diastolic function and brain natriuretic peptide. *J Hypertens* 2001;19:143–8.
- [12] Harris KF, Matthews KA. Interactions between autonomic nervous system activity and endothelial function: a model for the development of cardiovascular disease. *Psychosom Med* 2004;66:153–64.
- [13] Virtanen R, Jula A, Kuusela T, et al. Reduced heart rate variability in hypertension: associations with lifestyle factors and plasma renin activity. *J Hum Hypertens* 2003;17:171–9.
- [14] Chowdhary S, Vaile JC, Fletcher J, et al. Nitric oxide and cardiac autonomic control in humans. *Hypertension* 2000;36(2):264–9.
- [15] Tracey KJ. The inflammatory reflex. *Nature* 2002;420:853–9.
- [16] Carney RM, Freedland KE, Stein PK, et al. Heart rate variability and markers of inflammation and coagulation in depressed patients with coronary heart disease. *J Psychosom Res* 2007;62(4):463–7.
- [17] Chrysoshoou C, Pitsavos C, Panagiotakos DB, et al. Association between prehypertension status and inflammatory markers related to atherosclerotic disease: the ATTICA Study. *Am J Hypertens* 2004;17(7):568–73.
- [18] Vaziri ND. Causal link between oxidative stress, inflammation, and hypertension. *Iran J Kidney Dis* 2008;2:1–10.
- [19] Manzella D, Barbieri M, Ragno E, et al. Chronic administration of pharmacologic doses of vitamin E improves the cardiac autonomic nervous system in patients with type 2 diabetes. *Am J Clin Nutr* 2001;73:1052–7.