

**Home assignment № 1****Task 1.**

$U$  is a finite set,  $f : U \rightarrow U$  and  $f$  is a surjective function. It means that

$$\forall y \in U \exists x \in U : f(x) = y$$

Let  $f$  is **not** an injective function. Then  $\exists x_1 \neq x_2 : f(x_1) = f(x_2)$ . And also  $\exists y' \nexists x : f(x) = y'$ . But  $U$  is finite set and  $\forall y \in U \exists x \in U : f(x) = y$ . Therefore,

$$\forall x_1, x_2 : f(x_1) = f(x_2) \Rightarrow x_1 = x_2$$

It means that  $f$  is also an injective function.

**Task 2.**

a) Asymmetric and transitive: 4231

Compute this number by a Haskell script *TransitiveAsymmetricRelations.hs*.

b) Antisymmetric and antireflexive:  $3^{10}$

Consider the relation matrix  $P$ . Diagonal elements  $p_{ii}$  must be always 0. For non-diagonal elements:

$$\forall i, j : (p_{ij} = 1 \wedge p_{ji} = 0) \vee (p_{ij} = 1 \wedge p_{ji} = 0) \vee (p_{ij} = 0 \wedge p_{ji} = 0)$$

There are 10 pairs of different set elements and all of them have 3 of 4 variants of values. Therefore, we have a result:  $3^{10}$ .

**Task 3.**

In Task 2 I wrote a Haskell script (*TransitiveAsymmetricRelations.hs*) including a function *isTransitive*. This function get a relation matrix  $P$  ( $n \times n$ ), square it ( $O(n^3)$ ) and check that result of multiplication is including on original relation matrix  $P^2 \subseteq P$  ( $O(n^2)$ ). As result we have a complexity:  $O(n^3 + n^2) = O(n^3)$ .

**Task 4.****Task 5.**

$$R : (x_1, y_1)R(x_2, y_2) \leftrightarrow x_1 \leq x_2, y_1 \leq y_2.$$

$R$  is reflexive.

$$\forall (x, y) \in Z^2 : x \leq x, y \leq y, \text{ because } \leq \text{ is reflexive } \Rightarrow (x, y)R(x, y).$$

$R$  is transitive.

$$\begin{aligned} \forall (x_1, y_1), (x_2, y_2), (x_3, y_3) : (x_1, y_1)R(x_2, y_2), (x_2, y_2)R(x_3, y_3) &\Rightarrow x_1 \leq x_2, y_1 \leq y_2, x_2 \leq x_3, \\ y_2 \leq y_3. \leq \text{ is transitive } &\Rightarrow x_1 \leq x_3, y_1 \leq y_3 \Rightarrow (x_1, y_1)R(x_3, y_3). \end{aligned}$$

$R$  is antisymmetric.

$$\forall (x_1, y_1), (x_2, y_2) : (x_1, y_1)R(x_2, y_2), (x_2, y_2)R(x_1, y_1) \Rightarrow x_1 \leq x_2, y_1 \leq y_2, x_2 \leq x_1, y_2 \leq y_1.$$

$\leq$  is antisymmetric  $\Rightarrow x_1 = x_2, y_1 = y_2 \Rightarrow (x_1, y_1) = (x_2, y_2)$ .

It means that  $R$  is a partial order.

1)  $A_1 = (x, y) \mid x \leq 3, y \leq 4$   
 $\nexists \min, \max = (3, 4)$ .

2)  $A_2 = (x, y) \mid x^2 + y^2 \leq 4$   
 $\min = (-2, -2), \max = (2, 2)$ .