



张宇考研数学高等数学基础阶段模考试卷

一、选择题：1 ~ 10 小题，每小题 5 分，共 50 分。下列每题给出的四个选项中，只有一个选项是最符合题目要求的。

1. 设 $f(x) = \begin{cases} e^x, & x \leq 0, \\ x^2, & x > 0, \end{cases}$ 则 $I = \lim_{x \rightarrow 0^+} \left[\int_{-\infty}^x f(t) dt \right]^{\frac{1}{x(1-\cos x)}} = (\quad)$.

- (A) $e^{\frac{2}{3}}$ (B) $e^{\frac{3}{2}}$ (C) $\frac{2}{3}$ (D) $\frac{3}{2}$

2. 设 $g(x) = \begin{cases} 2-x, & x \leq 0, \\ 2+x, & x > 0, \end{cases}$ $f(x) = \begin{cases} x^2, & x < 0, \\ -x-1, & x \geq 0, \end{cases}$ 则 $x=0$ 是 $g[f(x)]$ 的().

- (A) 可去间断点 (B) 跳跃间断点 (C) 连续点 (D) 第二类间断点

3. (数学一、数学三) 设级数 $\sum_{n=1}^{\infty} a_n$ 与 $\sum_{n=1}^{\infty} b_n$ 都发散, 则().

- (A) $\sum_{n=1}^{\infty} (a_n + b_n)$ 发散 (B) $\sum_{n=1}^{\infty} a_n b_n$ 发散
(C) $\sum_{n=1}^{\infty} (|a_n| + |b_n|)$ 发散 (D) $\sum_{n=1}^{\infty} (a_n^2 + b_n^2)$ 发散

(数学二) 等边三角形当高为 8 时, 其面积对高的变化率为().

- (A) $\frac{4}{\sqrt{3}}$ (B) $\frac{8}{\sqrt{3}}$ (C) $\frac{16}{\sqrt{3}}$ (D) $\frac{32}{\sqrt{3}}$

4. 设 $f(x) = \ln(2+x)$, 则 $f'''(0) = (\quad)$.

- (A) $\frac{1}{2}$ (B) $\frac{1}{3}$ (C) $\frac{1}{4}$ (D) 1

5. 曲线 $f(x) = (x-1)^2(x-3)^2$ 的拐点个数为().

- (A) 0 (B) 1 (C) 2 (D) 3

6. 设 $y(x)$ 满足 $\Delta y = \frac{1-x}{\sqrt{2x-x^2}} \Delta x + o(\Delta x)$, 且 $y(0) = 0$, 则 $\int_0^2 y(x) dx = (\quad)$.

- (A) π (B) $\frac{\pi}{2}$ (C) $\frac{\pi}{3}$ (D) $\frac{\pi}{4}$

7. 设 n 为正整数, S_n 是第一象限内曲线 $y = n \cos nx$ 与该曲线在点 $(\frac{\pi}{2n}, 0)$ 处的切线所围成的平面图形的面积, 则().

- (A) S_n 与 n 有关 (B) S_n 与 n 无关
(C) 无法判断 S_n 与 n 的关系 (D) 以上都不正确

8. 设函数 $f(x), g(x)$ 均有二阶连续导数, 满足 $f(0) > 0, g(0) < 0$, 且 $f'(0) = g'(0) = 0$, 则函数 $z = f(x)g(y)$ 在点 $(0, 0)$ 处取得极小值的一个充分条件是().

- (A) $f''(0) < 0, g''(0) > 0$ (B) $f''(0) < 0, g''(0) < 0$
(C) $f''(0) > 0, g''(0) > 0$ (D) $f''(0) > 0, g''(0) < 0$

9. 设 $D: x^2 + y^2 \leq y, x \geq 0$, $f(x, y)$ 在 D 上连续, 且

$$f(x, y) = \sqrt{1 - x^2 - y^2} - \frac{8}{\pi} \iint_D f(x, y) dx dy,$$

则 $\iint_D f(x, y) dx dy = ()$.

- (A) $\frac{1}{6} \left(\frac{\pi}{2} - \frac{3}{2} \right)$ (B) $\frac{1}{6} \left(\frac{\pi}{2} - \frac{2}{3} \right)$ (C) $\frac{1}{6} \left(\frac{3}{2} - \frac{\pi}{2} \right)$ (D) $\frac{1}{6} \left(\frac{2}{3} - \frac{\pi}{2} \right)$

10. 设 $y = x$ 和 $y = e^{2x}$ 都是某 n 阶常系数齐次线性微分方程的解, 则 n 最低是().

- (A) 1 (B) 2 (C) 3 (D) 4

二、填空题: 11 ~ 16 小题, 每小题 5 分, 共 30 分.

11. 曲线 $y = \frac{(1+x)^{\frac{3}{2}}}{\sqrt{x}}$ 的斜渐近线方程为_____.

12. (数学一) 函数 $z = e^{3(x-1)+4y} \cos(xy)$ 在点 $(1, 0)$ 处的最大变化率为_____.

(数学二) 曲线 $\begin{cases} x = t - \ln(1+t^2), \\ y = \arctan t \end{cases}$ 在 $t = 2$ 处的曲率为_____.

(数学三) 设函数 $y = f(x)$ 由方程 $e^{2x+y} - \cos(xy) = e - 1$ 确定, 则曲线 $y = f(x)$ 在点 $(0, 1)$ 处的法线方程为_____.

13. (数学一) 设抛物面壳 $\Sigma: z = \frac{1}{2}(x^2 + y^2) (0 \leq z \leq 1)$ 的面密度为 z , 则该抛物面壳的质量为_____.

(数学二) 设不定积分 $\int \frac{x^2 + ax + 2}{(x+1)(x^2+1)} dx$ 的结果中不含反正切函数, 则 $a =$ _____.

(数学三) 设 E 表示 $[0, 4\pi]$ 这一闭区间, 则定积分 $\int_E |\cos x| \sqrt{\sin x} dx =$ _____.

14. (数学一) 若常数 λ 可使向量 $2xy(x^4 + y^2)^\lambda \mathbf{i} - x^2(x^4 + y^2)^\lambda \mathbf{j}$ 在右半平面 $x > 0$ 上成为某二元函数 $u(x, y)$ 的梯度, 则 $\lambda =$ _____.

(数学二、数学三) 设 $f(x+y, x-y) = 2(x^2 + y^2)e^{x^2 - y^2}$, 则 $f'_x(x, y) - f'_y(x, y) =$ _____.

15. 设 $\begin{cases} y' - \left(2x - \frac{1}{x}\right)y = x^2, x \geq 1, \\ y(1) = y_1, \end{cases}$ 若 $\lim_{x \rightarrow +\infty} \frac{y(x)}{x}$ 存在, 则 $y_1 =$ _____.

16. (数学一、数学三) 若幂级数 $\sum_{n=0}^{\infty} a^n x^n (a > 0)$ 的收敛域为 $(-\infty, +\infty)$, 则 a 应满足_____.

(数学二) 累次积分 $\int_0^{\frac{\pi}{2}} d\theta \int_0^{\cos \theta} f(r \cos \theta, r \sin \theta) r^2 dr$ 在直角坐标系先 x 后 y 的积分次序为_____.

三、解答题: 17 ~ 22 小题, 共 70 分. 解答应写出文字说明、证明过程或演算步骤.

17. (本题满分 10 分)

求 a, b 的值, 使得 $f(x) = \ln(1 - ax) + \frac{x}{1 + bx}$ 在 $x \rightarrow 0$ 时是 x 的 3 阶无穷小.

18. (本题满分 12 分)

设 $f(x)$ 可导, 满足 $xf'(x) = f'(-x) + 1, f(0) = 0$.

(1) 求 $f'(x)$ 的表达式;

(2) 求 $f(x)$ 的极值.

微信公众号【神灯考研】
考研人的精神家园

19. (本题满分 12 分)

求微分方程 $xdy + (x - 2y)dx = 0$ 的一个解 $y = y(x)$, 使得由曲线 $y = y(x)$ 与直线 $x = 1$, $x = 2$ 及 x 轴所围成的平面图形绕 x 轴旋转一周所得旋转体体积最小.

20. (本题满分 12 分)

现有函数 $u(x, t)$, 试利用变量代换 $\begin{cases} \xi = x - 2t, \\ \eta = x + 3t \end{cases}$ 将 u 关于变量 x, t 的方程 $6 \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial x \partial t} - \frac{\partial^2 u}{\partial t^2} = 0$ 化为 u 关于变量 ξ, η 的方程, 其中 u 具有二阶连续偏导数.

微信公众号【神灯考研】
考研人的精神家园

21. (本题满分 12 分)

(数学一) 计算 $I = \oint_{\Sigma} \frac{x dydz + y dzdx + z dxdy}{(x^2 + y^2 + 4z^2)^{\frac{3}{2}}}$, 其中 $\Sigma: (x-1)^2 + y^2 + z^2 = a^2 (a > 0 \text{ 且 } a \neq 1)$, 取外侧.

(数学二、数学三) 设 D 是由封闭曲线 $x^2 + y^2 = a(x + \sqrt{x^2 + y^2})$ 所围成的有界闭区域, 其中常数 $a > 0$. 求二重积分 $I = \iint_D [x^2 \ln(y + \sqrt{1 + y^2}) + x \sqrt{x^2 + y^2}] dx dy$.

微信公众号【神灯考研】
考研人的精神家园

22. (本题满分 12 分)

(数学一、数学三) 求幂级数 $\sum_{n=2}^{\infty} \frac{n}{n^2-1} x^n$ 的收敛域与和函数.

(数学二) 设 $f(x)$ 在 $[0,1]$ 上非负且连续.

(1) 证明: 存在 $\xi \in (0,1)$, 使得 $\xi f(\xi) = \int_{\xi}^1 f(t) dt$;

(2) 又设 $f(x)$ 在 $(0,1)$ 内可导, 且 $f'(x) > -\frac{2f(x)}{x}$, 证明(1) 中的 ξ 是唯一的.

微信公众号【神灯考研】
考研人的精神家园

张宇考研数学高等数学 基础阶段模考参考答案及评分细则

一、选择题

1. 答 应选(A).

解 当 $x > 0$ 时, $\int_{-\infty}^x f(t) dt = \int_{-\infty}^0 e^t dt + \int_0^x t^2 dt = 1 + \frac{1}{3}x^3$, 于是

$$I = \lim_{x \rightarrow 0^+} \left(1 + \frac{1}{3}x^3\right)^{\frac{1}{x(1-\cos x)}} \stackrel{1^\infty}{=} e^A,$$

其中
$$A = \lim_{x \rightarrow 0^+} \frac{1}{x(1-\cos x)} \cdot \left(1 + \frac{1}{3}x^3 - 1\right) = \lim_{x \rightarrow 0^+} \frac{1}{\frac{1}{2}x^3} \cdot \frac{1}{3}x^3 = \frac{2}{3},$$

故 $I = e^{\frac{2}{3}}$.

2. 答 应选(B).

解 方法一 先求出 $g[f(x)]$.

$$\begin{aligned} g[f(x)] &= \begin{cases} 2+f(x), & f(x) > 0, \\ 2-f(x), & f(x) \leq 0 \end{cases} = \begin{cases} 2+x^2, & x < 0, \\ 2-(-x-1), & x \geq 0 \end{cases} \\ &= \begin{cases} 2+x^2, & x < 0, \\ 3+x, & x \geq 0. \end{cases} \end{aligned}$$

$$\lim_{x \rightarrow 0^+} g[f(x)] = \lim_{x \rightarrow 0^+} (3+x) = 3, \quad \lim_{x \rightarrow 0^-} g[f(x)] = \lim_{x \rightarrow 0^-} (2+x^2) = 2.$$

因此, 应选(B).

方法二 不必先求 $g[f(x)]$ 的表达式, 直接计算 $\lim_{x \rightarrow 0^+} g[f(x)]$ 和 $\lim_{x \rightarrow 0^-} g[f(x)]$.

$$\lim_{x \rightarrow 0^+} g[f(x)] = \lim_{x \rightarrow 0^+} g(-x-1) = \lim_{x \rightarrow 0^+} [2 - (-x-1)] = 3,$$

$$\lim_{x \rightarrow 0^-} g[f(x)] = \lim_{x \rightarrow 0^-} g(x^2) = \lim_{x \rightarrow 0^-} (2+x^2) = 2.$$

因此, 应选(B).

3. (数学一、数学三) 答 应选(C).

解 用反证法证明(C)是正确的, 假设 $\sum_{n=1}^{\infty} (|a_n| + |b_n|)$ 收敛, 由于 $0 \leq |a_n| \leq |a_n| + |b_n|$,

于是 $\sum_{n=1}^{\infty} |a_n|$ 必收敛, 进而 $\sum_{n=1}^{\infty} a_n$ 也就收敛了, 这与 $\sum_{n=1}^{\infty} a_n$ 发散矛盾, 于是假设不成立, 所以

$\sum_{n=1}^{\infty} (|a_n| + |b_n|)$ 发散. 另外, 其余选项不正确, 可自行举反例排除. 微信公众号【神灯考研】

(数学二) 答 应选(C).

解 设 h 表示等边三角形的高, 于是面积 $S = \frac{h^2}{\sqrt{3}}$, 于是 $S'_h = \frac{2h}{\sqrt{3}}$, 当 $h = 8$ 时, $S'_h = \frac{16}{\sqrt{3}}$, 选(C). 考研人的精神家园

4. 答 应选(C).

解 $f(x) = \ln\left[2\left(1 + \frac{x}{2}\right)\right] = \ln 2 + \ln\left(1 + \frac{x}{2}\right) = \ln 2 + \frac{x}{2} - \frac{1}{2}\left(\frac{x}{2}\right)^2 + \frac{1}{3}\left(\frac{x}{2}\right)^3 - \dots$, 由 $\frac{f'''(0)}{3!} = \frac{1}{24}$, 得 $f'''(0) = \frac{1}{4}$.

5. 答 应选(C).

解 $f'(x) = 2(x-1)(x-3)^2 + 2(x-1)^2(x-3) = 2(x-1)(x-3)(2x-4)$, 显然 $f'(1) = f'(2) = f'(3) = 0$, 于是必存在 $\xi_1 \in (1, 2), \xi_2 \in (2, 3)$, 分别使得 $f''(\xi_1) = 0$ 和 $f''(\xi_2) = 0$, 而注意到 $f(x)$ 是四次多项式, 于是 $f''(x)$ 就是二次多项式, 综上 $f''(x) = 0$ 只有上述两个根 ξ_1 和 ξ_2 , 可以设 $f''(x) = 12(x-\xi_1)(x-\xi_2)$, 其中 $\xi_1 \in (1, 2), \xi_2 \in (2, 3)$, 显然 $f''(x)$ 在 ξ_1 和 ξ_2 两侧都变号, 于是 $f(x)$ 的拐点共有 2 个.

6. 答 应选(B).

解 由 $\Delta y = \frac{1-x}{\sqrt{2x-x^2}}\Delta x + o(\Delta x)$, 知 $y' = \frac{1-x}{\sqrt{2x-x^2}}$, 于是 $y = \int \frac{1-x}{\sqrt{2x-x^2}} dx = \frac{1}{2} \int \frac{1}{\sqrt{2x-x^2}} d(2x-x^2) = \sqrt{2x-x^2} + C$,

由 $y(0) = 0$, 知 $C = 0$, 所以 $y = \sqrt{2x-x^2}$, 进而

$$\int_0^2 y(x) dx = \int_0^2 \sqrt{2x-x^2} dx = \frac{1}{2} \pi \times 1^2 = \frac{1}{2} \pi.$$

7. 答 应选(B).

解 $y = n \cos nx$ 在点 $\left(\frac{\pi}{2n}, 0\right)$ 处的切线为 $y - 0 = y'\left(\frac{\pi}{2n}\right)\left(x - \frac{\pi}{2n}\right)$, 即 $y = -n^2 x + \frac{\pi}{2} n$, 注意到曲线 $y = n \cos nx$ 在区间 $\left(0, \frac{\pi}{2n}\right)$ 内是凸的, 于是切线在曲线上方, 从而所围图形面积 $S_n = \int_0^{\frac{\pi}{2n}} \left[\left(-n^2 x + \frac{\pi}{2} n\right) - n \cos nx \right] dx = \left(-\frac{n^2 x^2}{2} + \frac{\pi}{2} nx - \sin nx \right) \Big|_0^{\frac{\pi}{2n}} = \frac{\pi^2}{8} - 1$, 故 S_n 与 n 无关, 选(B).

8. 答 应选(A).

解 由 $z = f(x)g(y)$, 得

$$\frac{\partial z}{\partial x} = f'(x)g(y), \frac{\partial z}{\partial y} = f(x)g'(y),$$

$$\frac{\partial^2 z}{\partial x^2} = f''(x)g(y), \frac{\partial^2 z}{\partial x \partial y} = f'(x)g'(y), \frac{\partial^2 z}{\partial y^2} = f(x)g''(y),$$

$$A = \frac{\partial^2 z}{\partial x^2} \Big|_{(0,0)} = f''(0)g(0), B = \frac{\partial^2 z}{\partial x \partial y} \Big|_{(0,0)} = f'(0)g'(0) = 0,$$

$$C = \frac{\partial^2 z}{\partial y^2} \Big|_{(0,0)} = f(0)g''(0).$$

由于


$$\left. \frac{\partial z}{\partial x} \right|_{(0,0)} = f'(0)g(0) = 0, \left. \frac{\partial z}{\partial y} \right|_{(0,0)} = f(0)g'(0) = 0, f(0) > 0, g(0) < 0,$$

显然只有当 $f''(0) < 0, g''(0) > 0$ 时, $B^2 - AC < 0$, 且 $A > 0$, 即此时 $z = f(x)g(y)$ 在点 $(0,0)$ 处取得极小值. 因此选项(A) 是正确的.

9. 答 应选(B).

解 令 $\iint_D f(x,y) dx dy = A$, 则原等式成为 $f(x,y) = \sqrt{1-x^2-y^2} - \frac{8}{\pi}A$, 两端在 D 上同时作积分, 得

$$\begin{aligned} A &= \iint_D \sqrt{1-x^2-y^2} dx dy - \iint_D \frac{8}{\pi}A dx dy \\ &= \iint_D \sqrt{1-x^2-y^2} dx dy - \frac{8}{\pi}A \cdot \frac{1}{2}\pi \left(\frac{1}{2}\right)^2, \end{aligned}$$



从而 $A = \iint_D \sqrt{1-x^2-y^2} dx dy - A$, 所以

$$2A = \int_0^{\frac{\pi}{2}} d\theta \int_0^{\sin \theta} \sqrt{1-r^2} \cdot r dr = \frac{1}{3} \int_0^{\frac{\pi}{2}} (1 - \cos^3 \theta) d\theta = \frac{1}{3} \left(\frac{\pi}{2} - \frac{2}{3} \right),$$

$$\text{故 } A = \frac{1}{6} \left(\frac{\pi}{2} - \frac{2}{3} \right).$$

10. 答 应选(C).

分析 对 n 阶常系数齐次线性微分方程, 其特征根与解的关系如下.

若 λ 是单根, 必有解 $e^{\lambda x}$;

若 λ 是二重根, 必有解 $e^{\lambda x}, xe^{\lambda x}$;

若 λ 是三重根, 必有解 $e^{\lambda x}, xe^{\lambda x}, x^2 e^{\lambda x}$.

依次类推更高重实根即可.

若 $\alpha \pm \beta i$ 是一重虚根, 必有解 $e^{\alpha x} \cos \beta x, e^{\alpha x} \sin \beta x$;

若 $\alpha \pm \beta i$ 是二重虚根, 必有解 $e^{\alpha x} \cos \beta x, xe^{\alpha x} \cos \beta x, e^{\alpha x} \sin \beta x, xe^{\alpha x} \sin \beta x$.

依次类推更高重虚根即可.

解 $y = x = xe^{0x}$ 是解 $\Rightarrow 0$ 至少是二重根, 且 e^{0x} (也就是 1) 也是解, $y = e^{2x}$ 是解 $\Rightarrow 2$ 至少是单根, 综上 n 最低是 3, 选(C).

二、填空题

11. 答 应填 $y = x + \frac{3}{2}$.

$$\text{解 } k = \lim_{x \rightarrow +\infty} \frac{y}{x} = \lim_{x \rightarrow +\infty} \frac{\frac{(1+x)^{\frac{3}{2}}}{\sqrt{x}}}{x} = \lim_{x \rightarrow +\infty} \frac{(1+x)^{\frac{3}{2}}}{x\sqrt{x}} = 1,$$

微信公众号【神灯考研】

考研人的精神家园

$$\begin{aligned} b &= \lim_{x \rightarrow +\infty} (y - kx) = \lim_{x \rightarrow +\infty} \left[\frac{(1+x)^{\frac{3}{2}}}{\sqrt{x}} - x \right] = \lim_{x \rightarrow +\infty} \frac{(1+x)^{\frac{3}{2}} - x^{\frac{3}{2}}}{\sqrt{x}} \\ &= \lim_{x \rightarrow +\infty} x^{\frac{3}{2}} \cdot \frac{\left(1 + \frac{1}{x}\right)^{\frac{3}{2}} - 1}{\sqrt{x}} = \lim_{x \rightarrow +\infty} x^{\frac{3}{2}} \cdot \frac{\frac{3}{2} \cdot \frac{1}{x}}{\sqrt{x}} = \frac{3}{2}, \end{aligned}$$

于是斜渐近线方程为 $y = x + \frac{3}{2}$.

12. (数学一) 答 应填 5.

解 函数 $z = e^{3(x-1)+4y} \cos(xy)$ 在点 $(1,0)$ 处的梯度 $\text{grad } z|_{(1,0)} = (z'_x, z'_y)|_{(1,0)} = (3,4)$,

于是函数 $z = e^{3(x-1)+4y} \cos(xy)$ 在点 $(1,0)$ 处的最大变化率为 $\sqrt{3^2 + 4^2} = 5$.

(数学二) 答 应填 $\frac{5}{\sqrt{2}}$.

$$\text{解 } \frac{dy}{dx} = \frac{y'_t}{x'_t} = \frac{1}{(t-1)^2}, \frac{d^2y}{dx^2} = \frac{d}{dt} \left[\frac{1}{(t-1)^2} \right] \cdot \frac{dt}{dx} = \frac{-2}{(t-1)^3} \cdot \frac{1}{x'_t} = -\frac{2(1+t^2)}{(t-1)^5},$$

$$t=2 \text{ 处的曲率为 } \frac{|y''|}{(1+y'^2)^{\frac{3}{2}}} \Big|_{t=2} = \frac{5}{\sqrt{2}}.$$

(数学三) 答 应填 $y = \frac{1}{2}x + 1$.

解 方程 $e^{2x+y} - \cos(xy) = e - 1$ 两端同时对 x 求导, 得

$$e^{2x+y}(2+y') + \sin(xy) \cdot (y + xy') = 0,$$

将 $x=0, y=1$ 代入, 得 $y' = -2$, 于是点 $(0,1)$ 处的法线方程为 $y-1 = \frac{1}{2}(x-0)$, 即 $y = \frac{1}{2}x + 1$.

13. (数学一) 答 应填 $\frac{2\pi}{15}(6\sqrt{3}+1)$.

$$\begin{aligned} \text{解 质量 } m &= \iint_{\Sigma} z dS = \iint_{x^2+y^2 \leq 2} \frac{1}{2}(x^2+y^2) \cdot \sqrt{1+z_x'^2+z_y'^2} dx dy \\ &= \iint_{x^2+y^2 \leq 2} \frac{1}{2}(x^2+y^2) \cdot \sqrt{1+x^2+y^2} dx dy \\ &= \int_0^{2\pi} d\theta \int_0^{\sqrt{2}} \frac{1}{2}r^2 \cdot \sqrt{1+r^2} \cdot r dr \\ &= \frac{2\pi}{15}(6\sqrt{3}+1). \end{aligned}$$

(数学二) 答 应填 -1 .

解 $\frac{x^2+ax+2}{(x+1)(x^2+1)} = \frac{A}{x+1} + \frac{Bx+C}{x^2+1} = \frac{A}{x+1} + \frac{Bx}{x^2+1} + \frac{C}{x^2+1}$, 要使积分结果不含反正切函数, 必有 $C=0$, 此时 $\frac{x^2+ax+2}{(x+1)(x^2+1)} = \frac{A}{x+1} + \frac{Bx}{x^2+1}$, 通分, 对比两端分子, 有 $x^2 +$

$$ax + 2 = A(x^2 + 1) + Bx(x + 1) = (A + B)x^2 + Bx + A, \text{ 于是 } \begin{cases} 1 = A + B, \\ a = B, \\ 2 = A, \end{cases} \text{ 故 } a = -1.$$

(数学三) 答 应填 $\frac{8}{3}$.

分析 这里的 $\sqrt{\sin x}$ 是关键, $\sin x$ 只能在 $[0, \pi]$ 及 $[2\pi, 3\pi]$ 取非负值而使得 $\sqrt{\sin x}$ 有意义.

$$\begin{aligned} \text{解 } I &= \int_0^{\pi} |\cos x| \sqrt{\sin x} dx + \int_{2\pi}^{3\pi} |\cos x| \sqrt{\sin x} dx \\ &= \int_0^{\frac{\pi}{2}} \cos x \sqrt{\sin x} dx - \int_{\frac{\pi}{2}}^{\pi} \cos x \sqrt{\sin x} dx + \int_{2\pi}^{\frac{5\pi}{2}} \cos x \sqrt{\sin x} dx - \int_{\frac{5\pi}{2}}^{3\pi} \cos x \sqrt{\sin x} dx \\ &= \frac{8}{3}. \end{aligned}$$

14. (数学一) 答 应填 -1 .

解 令 $P(x, y) = 2xy(x^4 + y^2)^\lambda$, $Q(x, y) = -x^2(x^4 + y^2)^\lambda$. 由题意知 $\frac{\partial Q}{\partial x} = \frac{\partial P}{\partial y}$, 则

$$4x(x^4 + y^2)^\lambda(\lambda + 1) = 0.$$

于是, 当且仅当 $\lambda = -1$ 时, 所给向量场在右半平面 $x > 0$ 上是梯度场.

【注】 如何进一步求解 $u(x, y)$, 解答如下: 在 $x > 0$ 的半平面内, 以任意一点, 如 $(1, 0)$, 作为积分路径的起点, 则得

$$\begin{aligned} u(x, y) &= \int_{(1,0)}^{(x,y)} \frac{2xydx - x^2dy}{x^4 + y^2} + C \\ &= \int_1^x \frac{2x \cdot 0}{x^4 + 0^2} dx - \int_0^y \frac{x^2}{x^4 + y^2} dy + C \\ &= -\arctan \frac{y}{x^2} + C, \end{aligned}$$

式中, C 为任意常数.

(数学二、数学三) 答 应填 $(x - y)e^{xy}(2 - x^2 - y^2)$.

$$\text{解 由 } \begin{cases} x + y = u, \\ x - y = v \end{cases} \text{ 解得 } \begin{cases} x = \frac{u+v}{2}, \\ y = \frac{u-v}{2}, \end{cases} \text{ 于是 } f(u, v) = (u^2 + v^2)e^{uv}, \text{ 即 } f(x, y) = (x^2 +$$

$y^2)e^{xy}$, 所以 $f'_x(x, y) - f'_y(x, y) = (x - y)e^{xy}(2 - x^2 - y^2)$.

15. 答 应填 -1 .

解

$$y' - \left(2x - \frac{1}{x}\right)y = x^2,$$

$$y = e^{\int (2x - \frac{1}{x}) dx} \left[\int x^2 e^{-\int (2x - \frac{1}{x}) dx} dx + C \right] = \frac{e^{x^2}}{x} \left(\int x^3 e^{-x^2} dx + C \right) \\ = \frac{e^{x^2}}{x} \left(-\frac{1}{2} x^2 e^{-x^2} - \frac{1}{2} e^{-x^2} + C \right) = -\frac{x}{2} - \frac{1}{2x} + C \frac{e^{x^2}}{x}.$$

由 $y(1) = y_1$, 得 $C = (y_1 + 1)e^{-1}$, 得初值问题的解为

$$y = -\frac{x}{2} - \frac{1}{2x} + (y_1 + 1)e^{-1} \frac{e^{x^2}}{x}.$$

$$\lim_{x \rightarrow +\infty} \frac{y}{x} = \lim_{x \rightarrow +\infty} \left[-\frac{1}{2} - \frac{1}{2x^2} + (y_1 + 1)e^{-1} \frac{e^{x^2}}{x^2} \right].$$

若 $y_1 + 1 \neq 0$, 则上式最后一项趋于无穷, 故应取 $y_1 = -1$.

当且仅当 $y_1 = -1$ 时上述极限存在, 为 $-\frac{1}{2}$.

16. (数学一、数学三) 答 应填 $0 < a < 1$.


解 要想使得收敛域为 $(-\infty, +\infty)$, 只需 $\rho = \lim_{n \rightarrow \infty} \frac{a^{(n+1)^2}}{a^{n^2}} = \lim_{n \rightarrow \infty} a^{2n+1} = 0 (a > 0) \Rightarrow 0 < a < 1$.

(数学二) 答 应填 $\int_0^{\frac{1}{2}} dy \int_{\frac{1}{2}-\sqrt{\frac{1}{4}-y^2}}^{\frac{1}{2}+\sqrt{\frac{1}{4}-y^2}} f(x, y) \sqrt{x^2+y^2} dx$.

解 $r = \cos \theta \Rightarrow r^2 = r \cos \theta \Rightarrow x^2 + y^2 = x$ (见图), 配方得

$$\left(x - \frac{1}{2}\right)^2 + y^2 = \frac{1}{4}, \text{ 于是}$$

$$x = \frac{1}{2} \pm \sqrt{\frac{1}{4} - y^2}, 0 \leq y \leq \frac{1}{2},$$



所以

$$\int_0^{\frac{\pi}{2}} d\theta \int_0^{\cos \theta} f(r \cos \theta, r \sin \theta) r^2 dr = \int_0^{\frac{1}{2}} dy \int_{\frac{1}{2}-\sqrt{\frac{1}{4}-y^2}}^{\frac{1}{2}+\sqrt{\frac{1}{4}-y^2}} f(x, y) \sqrt{x^2+y^2} dx.$$

注意, 看清题目是 $r^2 dr$ 不是 $r dr$.

三、解答题

17. 解 $\ln(1-ax) = -ax - \frac{1}{2}(-ax)^2 + \frac{1}{3}(-ax)^3 + o(x^3), \dots\dots 3 \text{ 分}$

$$x \cdot \frac{1}{1+bx} = x \cdot [1 - bx + (-bx)^2 + o(x^2)] = x - bx^2 + b^2 x^3 + o(x^3),$$

$\dots\dots 6 \text{ 分}$

$$f(x) = \ln(1-ax) + \frac{x}{1+bx} = (-a+1)x - \left(\frac{1}{2}a^2+b\right)x^2 + \left(-\frac{1}{3}a^3+b^2\right)x^3 + o(x^3),$$

$\dots\dots 8 \text{ 分}$

若想使 $f(x)$ 在 $x \rightarrow 0$ 时是关于 x 的 3 阶无穷小, 则 $-a+1=0, -\left(\frac{1}{2}a^2+b\right)=0$, 即 $a=$

1, $b = -\frac{1}{2}$, 此时 $f(x) = -\frac{1}{12}x^3 + \cdots$ 刚好是 x 的 3 阶无穷小.10 分

18. 解 (1) 将 $xf'(x) = f'(-x) + 1$ 中的 x 替换为 $-x$, 得 $-xf'(-x) = f'(x) + 1$,3 分

结合两式, 解得 $f'(x) = \frac{x-1}{x^2+1}$5 分

(2) 由 $f'(x) = \frac{x-1}{x^2+1} = 0$, 得唯一驻点 $x = 1$6 分

当 $x < 1$ 时, $f'(x) < 0$, 当 $x > 1$ 时, $f'(x) > 0$, 于是 $f(1)$ 是 $f(x)$ 的唯一极小值.8 分

$$\begin{aligned} f(1) &= \int_0^1 f'(x) dx + f(0) = \int_0^1 \frac{x-1}{x^2+1} dx + 0 \\ &= \int_0^1 \frac{x}{x^2+1} dx - \int_0^1 \frac{1}{x^2+1} dx = \frac{1}{2} \ln 2 - \frac{\pi}{4}. \end{aligned} \quad \text{.....12 分}$$

19. 解 原方程可化为

$$\frac{dy}{dx} - \frac{2}{x}y = -1,$$

这是一阶非齐次线性微分方程, 故直接套用公式得

$$y = e^{\int \frac{2}{x} dx} \left(-\int e^{-\int \frac{2}{x} dx} dx + C \right) = x^2 \left(\frac{1}{x} + C \right) = x + Cx^2. \quad \text{.....4 分}$$

由曲线 $y = x + Cx^2$, 直线 $x = 1, x = 2$ 以及 x 轴所围成的平面图形绕 x 轴旋转一周所得旋转体体积为

$$V(C) = \int_1^2 \pi(x + Cx^2)^2 dx = \pi \left(\frac{x^3}{3} + \frac{C}{2}x^4 + \frac{C^2}{5}x^5 \right) \Big|_1^2 = \pi \left(\frac{31}{5}C^2 + \frac{15}{2}C + \frac{7}{3} \right), \quad \text{.....8 分}$$

令 $V'(C) = 0$, 得

$$\pi \left(\frac{62}{5}C + \frac{15}{2} \right) = 0,$$

解出 $C = -\frac{75}{124}$10 分

又 $V''(C) = \frac{62}{5}\pi > 0$, 故 $C = -\frac{75}{124}$ 为唯一极小值点, 也就是最小值点. 因此

$$y = x - \frac{75}{124}x^2$$

为所求解.12 分

20. 解 由于 $\xi = x - 2t, \eta = x + 3t$, 有

$$\frac{\partial u}{\partial x} = \frac{\partial u}{\partial \xi} \cdot 1 + \frac{\partial u}{\partial \eta} \cdot 1, \frac{\partial u}{\partial t} = \frac{\partial u}{\partial \xi} \cdot (-2) + \frac{\partial u}{\partial \eta} \cdot 3. \quad \text{.....2 分}$$

进一步,

$$\frac{\partial^2 u}{\partial x^2} = \frac{\partial^2 u}{\partial \xi^2} \cdot 1 + \frac{\partial^2 u}{\partial \xi \partial \eta} \cdot 1 + \frac{\partial^2 u}{\partial \eta \partial \xi} \cdot 1 + \frac{\partial^2 u}{\partial \eta^2} \cdot 1 = \frac{\partial^2 u}{\partial \xi^2} + 2 \frac{\partial^2 u}{\partial \xi \partial \eta} + \frac{\partial^2 u}{\partial \eta^2}, \quad \cdots \cdots 4 \text{ 分}$$

$$\begin{aligned} \frac{\partial^2 u}{\partial t^2} &= -2 \left[\frac{\partial^2 u}{\partial \xi^2} \cdot (-2) + \frac{\partial^2 u}{\partial \xi \partial \eta} \cdot 3 \right] + 3 \left[\frac{\partial^2 u}{\partial \eta \partial \xi} \cdot (-2) + \frac{\partial^2 u}{\partial \eta^2} \cdot 3 \right] \\ &= 4 \frac{\partial^2 u}{\partial \xi^2} - 12 \frac{\partial^2 u}{\partial \xi \partial \eta} + 9 \frac{\partial^2 u}{\partial \eta^2}, \quad \cdots \cdots 6 \text{ 分} \end{aligned}$$

$$\begin{aligned} \frac{\partial^2 u}{\partial x \partial t} &= \frac{\partial^2 u}{\partial \xi^2} \cdot (-2) + \frac{\partial^2 u}{\partial \xi \partial \eta} \cdot 3 + \frac{\partial^2 u}{\partial \eta \partial \xi} \cdot (-2) + \frac{\partial^2 u}{\partial \eta^2} \cdot 3 \\ &= -2 \frac{\partial^2 u}{\partial \xi^2} + \frac{\partial^2 u}{\partial \xi \partial \eta} + 3 \frac{\partial^2 u}{\partial \eta^2}, \quad \cdots \cdots 8 \text{ 分} \end{aligned}$$

此时, 方程 $6 \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial x \partial t} - \frac{\partial^2 u}{\partial t^2} = 0$ 化为

$$6 \left(\frac{\partial^2 u}{\partial \xi^2} + 2 \frac{\partial^2 u}{\partial \xi \partial \eta} + \frac{\partial^2 u}{\partial \eta^2} \right) + \left(-2 \frac{\partial^2 u}{\partial \xi^2} + \frac{\partial^2 u}{\partial \xi \partial \eta} + 3 \frac{\partial^2 u}{\partial \eta^2} \right) - \left(4 \frac{\partial^2 u}{\partial \xi^2} - 12 \frac{\partial^2 u}{\partial \xi \partial \eta} + 9 \frac{\partial^2 u}{\partial \eta^2} \right) = 0, \quad \cdots \cdots 10 \text{ 分}$$

从而得 $25 \frac{\partial^2 u}{\partial \xi \partial \eta} = 0$, 即 $\frac{\partial^2 u}{\partial \xi \partial \eta} = 0$12 分

21. (数学一) 分析 若 a 比较小, 当 $0 < a < 1$ 时, Σ 没有包含原点 $(0, 0, 0)$, 此时可以直接使用高斯公式; 若 a 比较大, 当 $a > 1$ 时, Σ 包含原点 $(0, 0, 0)$, 则可在挖去原点 $(0, 0, 0)$ 后使用高斯公式.

解 容易验证 $\frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z} = 0, (x, y, z) \neq (0, 0, 0)$2 分

若 $0 < a < 1$, 则 $I = \iiint_{\Omega} \left(\frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z} \right) dv = 0$;4 分

若 $a > 1$, 构造 $\Sigma_0: x^2 + y^2 + 4z^2 = \epsilon^2$, 取外侧, 此时根据高斯公式, 有

$$\oiint_{\Sigma + \Sigma_0} \frac{xdydz + ydzdx + zdx dy}{(x^2 + y^2 + 4z^2)^{\frac{3}{2}}} = 0,$$

故

$$I = \oiint_{\Sigma + \Sigma_0} - \iint_{\Sigma_0} = 0 - \iint_{\Sigma_0} = \iint_{\Sigma_0} \frac{xdydz + ydzdx + zdx dy}{(x^2 + y^2 + 4z^2)^{\frac{3}{2}}} \quad \cdots \cdots 6 \text{ 分}$$

$$= \frac{1}{\epsilon^3} \iint_{\Sigma_0} xdydz + ydzdx + zdx dy \quad \cdots \cdots 8 \text{ 分}$$

$$= \frac{1}{\epsilon^3} \iiint_{\Omega: x^2 + y^2 + 4z^2 \leq \epsilon^2} (1 + 1 + 1) dv \quad \cdots \cdots 10 \text{ 分}$$

$$= \frac{3}{\epsilon^3} \cdot \frac{4}{3} \pi \cdot \epsilon \cdot \epsilon \cdot \frac{\epsilon}{2} = 2\pi. \quad \cdots \cdots 12 \text{ 分}$$

(数学二、数学三) 解 曲线 $x^2 + y^2 = a(x + \sqrt{x^2 + y^2})$ 在极坐标系下是 $r = a(1 + \cos \theta), a > 0$,4 分

这是心形线, 如下图所示. 由于 $x^2 \ln(y + \sqrt{1 + y^2})$ 对 y 是奇函数, 且积分区域 D 关于 x 轴对

称,则


$$I = 0 + 2 \iint_{D_t} x \sqrt{x^2 + y^2} dx dy \quad \dots\dots 6 \text{ 分}$$

$$= 2 \int_0^\pi d\theta \int_0^{a(1+\cos\theta)} r^3 \cos\theta dr \quad \dots\dots 8 \text{ 分}$$

$$= \frac{a^4}{2} \int_0^\pi (1 + \cos\theta)^4 \cos\theta d\theta$$

$$= \frac{a^4}{2} \int_0^\pi (1 + 4\cos\theta + 6\cos^2\theta + 4\cos^3\theta + \cos^4\theta) \cos\theta d\theta \quad \dots\dots 10 \text{ 分}$$

$$= a^4 \int_0^{\frac{\pi}{2}} (4\cos^2\theta + 4\cos^4\theta) d\theta = \frac{7\pi a^4}{4}. \quad \dots\dots 12 \text{ 分}$$



22. (数学一、数学三) 解 (1) 由

$$\lim_{n \rightarrow \infty} \frac{\frac{n+1}{(n+1)^2 - 1}}{\frac{n}{n^2 - 1}} = \lim_{n \rightarrow \infty} \frac{n+1}{(n+1)^2 - 1} \cdot \frac{n^2 - 1}{n} = 1,$$

知收敛半径 $R = 1$. \dots\dots 2 \text{ 分}

收敛区间为 $(-1, 1)$, 对 $x = -1$, 原级数成为 $\sum_{n=2}^{\infty} (-1)^n \frac{n}{n^2 - 1}$, 是交错级数, 满足莱布尼茨定理, 故收敛;

对 $x = 1$, 原级数成为 $\sum_{n=2}^{\infty} \frac{n}{n^2 - 1}$, 是正项级数, 与 $\sum_{n=2}^{\infty} \frac{1}{n}$ 同敛散, 故发散.

综上, 收敛域为 $[-1, 1)$. \dots\dots 4 \text{ 分}

$$\begin{aligned} (2) \sum_{n=2}^{\infty} \frac{n}{n^2 - 1} x^n &= \sum_{n=2}^{\infty} \frac{n}{(n-1)(n+1)} x^n = \frac{1}{2} \sum_{n=2}^{\infty} \left(\frac{1}{n-1} + \frac{1}{n+1} \right) x^n \\ &= \frac{1}{2} \sum_{n=2}^{\infty} \frac{1}{n-1} x^n + \frac{1}{2} \sum_{n=2}^{\infty} \frac{1}{n+1} x^n \\ &= \frac{x}{2} \sum_{n=2}^{\infty} \frac{1}{n-1} x^{n-1} + \frac{1}{2x} \sum_{n=2}^{\infty} \frac{1}{n+1} x^{n+1}, \quad -1 < x < 1, \text{ 且 } x \neq 0. \quad \dots\dots 6 \text{ 分} \end{aligned}$$

记 $S_1(x) = \sum_{n=2}^{\infty} \frac{1}{n-1} x^{n-1}$, $S_1(0) = 0$, 则 $S_1'(x) = \sum_{n=2}^{\infty} x^{n-2}$ (等比级数) $= \frac{1}{1-x}$, 于是

$$S_1(x) = \int_0^x S_1'(t) dt + S_1(0) = \int_0^x \frac{1}{1-t} dt + 0 = -\ln(1-x); \quad \dots\dots 8 \text{ 分}$$

记 $S_2(x) = \sum_{n=2}^{\infty} \frac{1}{n+1} x^{n+1}$, $S_2(0) = 0$, 则 $S_2'(x) = \sum_{n=2}^{\infty} x^n$ (等比级数) $= \frac{x^2}{1-x}$, 于是

$$S_2(x) = \int_0^x S_2'(t) dt + S_2(0) = \int_0^x \frac{t^2}{1-t} dt + 0 = \int_0^x \frac{t^2 - 1 + 1}{1-t} dt = -\frac{x^2}{2} - x - \ln(1-x);$$

.....10 分

所以 $\sum_{n=2}^{\infty} \frac{n}{n^2-1} x^n = -\frac{x}{2} \ln(1-x) - \frac{1}{2x} \left[\frac{x^2}{2} + x + \ln(1-x) \right]$, $-1 \leq x < 1$, 且 $x \neq 0$;

而 $\sum_{n=2}^{\infty} \frac{n}{n^2-1} x^n \Big|_{x=0} = \left(\frac{2}{3} x^2 + \dots \right) \Big|_{x=0} = 0$.

综上,

$$\sum_{n=2}^{\infty} \frac{n}{n^2-1} x^n = \begin{cases} -\frac{x}{2} \ln(1-x) - \frac{1}{2x} \left[\frac{x^2}{2} + x + \ln(1-x) \right], & -1 \leq x < 1, \text{ 且 } x \neq 0, \\ 0, & x = 0. \end{cases}$$

.....12 分

(数学二) 证明 (1) 令

$$F(x) = -x \int_x^1 f(t) dt,$$

.....4 分

则

$$F(0) = 0, F(1) = 0.$$

.....5 分

由罗尔定理知, 存在 $\xi \in (0, 1)$, 使得 $F'(\xi) = 0$, 即 $-\int_{\xi}^1 f(t) dt - \xi[-f(\xi)] = 0$, 亦即

$$\xi f(\xi) = \int_{\xi}^1 f(t) dt.$$

.....6 分

(2) 对于方程 $xf(x) = \int_x^1 f(t) dt$, 在(1)中已证明了其在 $(0, 1)$ 内至少有一个根 ξ , 再考虑

$$\left[xf(x) - \int_x^1 f(t) dt \right]' = f(x) + xf'(x) + f(x) = 2f(x) + xf'(x), \quad \text{.....8 分}$$

由于 $f'(x) > -\frac{2f(x)}{x}$, $0 < x < 1$, 于是

$$2f(x) + xf'(x) > 0, 0 < x < 1,$$

所以

$$\left[xf(x) - \int_x^1 f(t) dt \right]' > 0,$$

.....10 分

也就是说 $xf(x) - \int_x^1 f(t) dt$ 在 $(0, 1)$ 内单调, 故上述 ξ 唯一.

.....12 分

微信公众号【神灯考研】
考研人的精神家园