

**Theorem**

Let the homogeneous linear equation with constant coefficients  $a_0T(n) + a_1T(n-1) + \dots + a_kT(n-k) = 0$  be given. If its characteristic equation  $a_0r^k + a_1r^{k-1} + \dots + a_kr^0 = 0$  has  $k$  distinct solutions  $r_1, r_2, \dots, r_k$ , then the solution is given by  $T(n) = c_1r_1^n + c_2r_2^n + \dots + c_kr_k^n$  where  $c_i$  are constants.

**Example1:**

Solve  $T(0) = 0, T(1) = 1, T(n) - 3T(n-1) - 4T(n-2) = 0$  if  $n > 1$

**Solution**

**Step1: Find the characteristic equation:**  $r^2 - 3r - 4 = 0, (r-4)(r+1) = 0$ , implies  $r = 4$  or  $r = -1$

**Step2: Apply the theorem:**  $T(n) = c_14^n + c_2(-1)^n$

**Step 3: Find  $c_1$  and  $c_2$** 

$$T(0) = c_14^0 + c_2(-1)^0 = 0 \text{ implies } c_1 + c_2 = 0 \dots (1)$$

$$T(1) = c_14^1 + c_2(-1)^1 = 1 \text{ implies } 4c_1 - c_2 = 1 \dots (2)$$

Solve equations (1) & (2) to get  $c_1 = \frac{1}{5}, c_2 = -\frac{1}{5}$ . Hence, the solution is:  $T(n) = \frac{1}{5}4^n - \frac{1}{5}(-1)^n$

**Example2:**

Solve  $T(0) = 0, T(1) = 1, T(n) - T(n-1) - T(n-2) = 0$  if  $n > 1$

**Solution**

**Step1: Find the characteristic equation:**  $r^2 - r - 1 = 0, r = \frac{1 \pm \sqrt{5}}{2}$

**Step2: Apply the theorem:**  $T(n) = c_1(\frac{1+\sqrt{5}}{2})^n + c_2(\frac{1-\sqrt{5}}{2})^n$

**Step 3: Find  $c_1$  and  $c_2$** 

$$T(0) = c_1(\frac{1+\sqrt{5}}{2})^0 + c_2(\frac{1-\sqrt{5}}{2})^0 = 0 \text{ implies } c_1 + c_2 = 0 \dots (1)$$

$$T(1) = c_1(\frac{1+\sqrt{5}}{2})^1 + c_2(\frac{1-\sqrt{5}}{2})^1 = 1 \text{ implies } c_1(\frac{1+\sqrt{5}}{2}) + c_2(\frac{1-\sqrt{5}}{2}) = 1 \dots (2)$$

Solve equations (1) & (2) to get  $c_1 = \frac{1}{\sqrt{5}}, c_2 = -\frac{1}{\sqrt{5}}$ . Hence, the solution is:  $T(n) = ((\frac{1+\sqrt{5}}{2})^n - (\frac{1-\sqrt{5}}{2})^n) / \sqrt{5}$

**Theorem**

Let  $r$  be a root of multiplicity  $m$ , then  $T(n) = r^n, T(n) = nr^n, T(n) = n^2r^n, \dots, T(n) = n^{m-1}r^n$  are all solutions.

**Example:**

Solve  $T(0) = 0, T(1) = 1, T(2) = 2, T(n) - 7T(n-1) + 15T(n-2) - 9T(n-3) = 0$  if  $n > 2$

**Solution**

**Step1: Find the characteristic equation:**  $r^3 - 7r^2 + 15r - 9 = 0$  implies  $(r-1)(r-3)^2 = 0$ , implies  $r = 1$  or  $r = 3$  of multiplicity 2

**Step2: Apply the theorem:**  $T(n) = c_1 1^n + c_2 3^n + c_3 n 3^n$

**Step 3: Find  $c_1, c_2$ , and  $c_3$**

$$c_1 = -1, c_2 = 1, c_3 = -\frac{1}{3}$$

Hence the solution is:  $T(n) = (-1)1^n + (1)3^n + \left(-\frac{1}{3}\right)n3^n = -1 + 3^n - n3^{n-1}$