

# Lecture 33: Residue theorem

Math 660—Jim Fowler

Thursday, August 5, 2010



## Theorem (Cauchy)

*If  $f(z)$  is analytic in a region  $\Omega$ , then*

$$n(\gamma, a) f(a) = \frac{1}{2\pi i} \int_{\gamma} \frac{f(z) dz}{z - a}$$

*if  $\gamma$  is homologous to zero in  $\Omega$ .*

The **residue** of  $f(z)$  at an isolated singularity  $a$  is the complex number  $R = \operatorname{Res}_{z=a} f(z)$  which makes

$$f(z) - \frac{R}{z - a} = F'(z)$$

for some single-valued analytic function  $F$  defined in a small annulus around  $a$ .

Suppose  $\Omega$  is simply connected, and  
Suppose  $\Omega' = \Omega - \{a_1, \dots, a_n\}$ .

Suppose  $\Omega$  is simply connected, and

Suppose  $\Omega' = \Omega - \{a_1, \dots, a_n\}$ .

Then for any  $\gamma \in \Omega'$ ,

$$[\gamma] = [c_1\gamma_1 + \dots + c_n\gamma_n] \in H_1(\Omega')$$

for curves  $\gamma_i$  around  $a_i$ , with  $c_i = n(\gamma, a_i)$ .

$$\frac{1}{2\pi i} \int_{\gamma} f \, dz$$

$$\frac{1}{2\pi i} \int_{\gamma} f \, dz = \sum_j \frac{c_j}{2\pi i} \int_{\gamma_j} f \, dz$$



$$\begin{aligned}\frac{1}{2\pi i} \int_{\gamma} f \, dz &= \sum_j \frac{c_j}{2\pi i} \int_{\gamma_j} f \, dz \\ &= \sum_j \frac{c_j}{2\pi i} \int_{\gamma_j} \frac{\operatorname{Res}_{z=a_i} f(z)}{z-a} \, dz\end{aligned}$$

$$\begin{aligned}
\frac{1}{2\pi i} \int_{\gamma} f \, dz &= \sum_j \frac{c_j}{2\pi i} \int_{\gamma_j} f \, dz \\
&= \sum_j \frac{c_j}{2\pi i} \int_{\gamma_j} \frac{\operatorname{Res}_{z=a_i} f(z)}{z-a} \, dz \\
&= \sum_j c_j \operatorname{Res}_{z=a_i} f(z)
\end{aligned}$$

$$\begin{aligned}
\frac{1}{2\pi i} \int_{\gamma} f \, dz &= \sum_j \frac{c_j}{2\pi i} \int_{\gamma_j} f \, dz \\
&= \sum_j \frac{c_j}{2\pi i} \int_{\gamma_j} \frac{\operatorname{Res}_{z=a_i} f(z)}{z - a} \, dz \\
&= \sum_j c_j \operatorname{Res}_{z=a_i} f(z) \\
&= \sum_j n(\gamma, a_i) \cdot \operatorname{Res}_{z=a_i} f(z)
\end{aligned}$$

$$\begin{aligned}
\frac{1}{2\pi i} \int_{\gamma} f \, dz &= \sum_j \frac{c_j}{2\pi i} \int_{\gamma_j} f \, dz \\
&= \sum_j \frac{c_j}{2\pi i} \int_{\gamma_j} \frac{\operatorname{Res}_{z=a_i} f(z)}{z - a} \, dz \\
&= \sum_j c_j \operatorname{Res}_{z=a_i} f(z) \\
&= \sum_j n(\gamma, a_i) \cdot \operatorname{Res}_{z=a_i} f(z)
\end{aligned}$$

This is Cauchy's **residue theorem** for when there are finitely many singularities.

If there are infinitely many singularities, still only finitely many have nonzero winding number with  $\gamma$ ,

If there are infinitely many singularities, still only finitely many have nonzero winding number with  $\gamma$ , and therefore

## **Theorem (Residue theorem)**

*Let  $f(z)$  be analytic in  $\Omega$ , except for isolated singularities  $a_j$ . Then,*

$$\frac{1}{2\pi i} \int_{\gamma} f \, dz = \sum_j n(\gamma, a_j) \cdot \operatorname{Res}_{z=a_j} f(z)$$

*for any cycle which is homologous to zero in  $\Omega$  and does not pass through any of the  $a_j$ .*

If only we had a procedure for computing  
 $\operatorname{Res}_{z=a} f(z) \dots$

If only we had a procedure for computing  
 $\operatorname{Res}_{z=a} f(z) \dots$

Wait, we do!



Consider  $f(z) = b_k(z - a)^{-k} + \cdots b_1(z - a)^{-1} + \varphi(z)$ ,  
for  $\varphi$  analytic.

Consider  $f(z) = b_k(z - a)^{-k} + \cdots + b_1(z - a)^{-1} + \varphi(z)$ ,  
for  $\varphi$  analytic.

Then  $\operatorname{Res}_{z=a} f(z) = b_1$ .

Consider  $f(z) = b_k(z - a)^{-k} + \cdots + b_1(z - a)^{-1} + \varphi(z)$ ,  
for  $\varphi$  analytic.

Then  $\operatorname{Res}_{z=a} f(z) = b_1$ .

For *simple* poles, it is even easier: then

$$\operatorname{Res}_{z=a} f(z) = \lim_{z \rightarrow a} (z - a) \cdot f(z).$$

# Example 1

$$f(z) = \frac{e^z}{(z-a)(z-b)} \text{ with } a \neq b.$$

Compute  $\int_{\gamma} f \, dz$  for a curve around  $a$  and  $b$ .

## Example 2

$$f(z) = \frac{1}{z^2 + z - 1}.$$

Compute  $\int_{\gamma} f \, dz$  for  $\gamma = \partial B_3(0)$ .

## Example 3

$$f(z) = \frac{1}{\sin z}$$

Compute  $\int_{\gamma} f \, dz$  for  $\gamma$  a circle of radius one around the origin.

## Example 3

$$f(z) = \frac{1}{z^4 + 4}$$

Compute  $\int_{\gamma} f \, dz$  for  $\gamma$  a circle of radius two and center 1.

# Example 4

Evaluate

$$\int_{-\infty}^{\infty} e^{-z^2}, dz$$