

Lecture 41: Taylor series

Math 660—Jim Fowler

Tuesday, August 17, 2010

Taylor series

If $f(z)$ is analytic in the region $\Omega \ni z_0$, then

$$f(z) = \sum_{n=0}^{\infty} \frac{f^{(n)}(z_0)}{n!} (z - z_0)^n$$

on the largest open disk of center z_0 contained in Ω .

Taylor series for log

Taylor series for log

$$\log(1 + z) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1} z^n}{n}$$

Taylor series for log

$$\log(1 + z) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1} z^n}{n}$$

Sums and products

Find the Taylor series for a product $f(z)g(z)$,
given the Taylor series for $f(z)$ and $g(z)$.

Sums and products

Find the Taylor series for a product $f(z)g(z)$, given the Taylor series for $f(z)$ and $g(z)$.

What is the radius of convergence?

Composition

Find the Taylor series for a composition $f(g(z))$, given the Taylor series for $f(z)$ and $g(z)$.

Inverse function

Find the Taylor series for $f^{-1}(z)$,
given the Taylor series for $f(z)$.

Laurent series

A **Laurent series** is one of the form

$$\sum_{n=-\infty}^{\infty} a_n z^n.$$

The series is **convergent** if the parts consisting of nonnegative powers and negative powers are separately convergent.

Laurent series

A **Laurent series** is one of the form

$$\sum_{n=-\infty}^{\infty} a_n z^n.$$

The series is **convergent** if the parts consisting of nonnegative powers and negative powers are separately convergent.

Convergence occurs in an annulus.