

Lecture 1: Introduction to complex numbers

Math 660—Jim Fowler

Monday, June 20, 2011

Syllabus

Syllabus

Summer 2011

Math 660

Jim Fowler

This is a beginning graduate course in complex analysis; as such, we study *analysis* (e.g., the rigorous foundations of calculus) as it applies to *functions of a complex variable*. The resulting theory is strikingly beautiful.

Resources

We present 5 resources to help you to learn complex analysis.

Professor's office hours

If you have questions, want to work through problems, or just talk about mathematics, please attend office hours.

Name: Jim Fowler

Office: MW658 Mathematics Tower

Phone: (773) 809-5659

Email: fowler@math.osu.edu


Website: <http://www.math.osu.edu/~fowler/>

Office Hours: Monday 3:30–5:00P.M.

Wednesday 3:30–5:00P.M.

and by appointment

Please email me with any concerns you have; the success of this course depends on open communication.



Today's Goal

Chapter 1 of *Complex Analysis*

We're going to move fast—some details won't be justified entirely for a while.

$$i = \sqrt{-1}$$

$$i = \sqrt{-1}$$

$$i^2 =$$

$$i = \sqrt{-1}$$

$$i^2 = -1$$

$$i = \sqrt{-1}$$

$$i^2 = -1 \quad (-i)^2 =$$

$$i = \sqrt{-1}$$

$$i^2 = -1 \quad (-i)^2 = -1$$

$$a + bi \in \mathbb{C}$$

$$a, b \in \mathbb{R}$$

The Complex Field

$$(a + bi) + (c + di) = (a + c) + (b + d)i$$

The Complex Field

$$(a + bi) + (c + di) = (a + c) + (b + d)i$$

$$(a + bi) - (c + di) = (a - c) + (b - d)i$$

The Complex Field

$$(a + bi) + (c + di) = (a + c) + (b + d)i$$

$$(a + bi) - (c + di) = (a - c) + (b - d)i$$

$$(a + bi) \cdot (c + di) = (ac - bd) + (ad + bc)i$$

The Complex Field

$$(a + bi) + (c + di) = (a + c) + (b + d)i$$

$$(a + bi) - (c + di) = (a - c) + (b - d)i$$

$$(a + bi) \cdot (c + di) = (ac - bd) + (ad + bc)i$$

$$\frac{a + bi}{c + di} = \left(\frac{ac + bd}{c^2 + d^2} \right) + \left(\frac{bc - ad}{c^2 + d^2} \right) i$$

Dividing Complex Numbers

$$\frac{a + bi}{c + di} =$$

Putting complex numbers in order

If $i > 0$, then $i^2 > 0$, meaning $-1 > 0$.

Putting complex numbers in order

If $i > 0$, then $i^2 > 0$, meaning $-1 > 0$.

If $i < 0$, then $i^2 > 0$, meaning $-1 > 0$.

Putting complex numbers in order

If $i > 0$, then $i^2 > 0$, meaning $-1 > 0$.

If $i < 0$, then $i^2 > 0$, meaning $-1 > 0$.

Something is wrong...

Putting complex numbers in order

If $i > 0$, then $i^2 > 0$, meaning $-1 > 0$.

If $i < 0$, then $i^2 > 0$, meaning $-1 > 0$.

Something is wrong...

The complex numbers aren't ordered.

Why write i ? Why not $\sqrt{-1}$?

$$\sqrt{a} \sqrt{b} = \sqrt{ab}$$

Why write i ? Why not $\sqrt{-1}$?

$$\sqrt{a} \sqrt{b} = \sqrt{ab}$$

But what if $a = -1$ and $b = -1$...?

Why write i ? Why not $\sqrt{-1}$?

$$\sqrt{a} \sqrt{b} = \sqrt{ab}$$

But what if $a = -1$ and $b = -1$...

$$\sqrt{-1} \sqrt{-1} = -\sqrt{(-1) \cdot (-1)}$$

There are two square roots. This is confusing.
Choose one of them, and call it i .

Complex Conjugation

Definition

Define $\overline{a + bi} = a - bi$.

$$\overline{\overline{z}} = z. \qquad \overline{a + b} = \overline{a} + \overline{b}. \qquad \overline{ab} = \overline{a} \cdot \overline{b}.$$

Complex Conjugation

Definition

Define $\overline{a + bi} = a - bi$.

$$\overline{\overline{z}} = z. \qquad \overline{a + b} = \overline{a} + \overline{b}. \qquad \overline{ab} = \overline{a} \cdot \overline{b}.$$

Could you tell the difference between a world in which i was replaced by $-i$?

Absolute Value

Definition

$$|z|^2 = z \cdot \bar{z}.$$

Absolute Value

Definition

$$|z|^2 = z \cdot \bar{z}.$$

- ▶ $|z| \geq 0$, where $|z| = 0$ iff $z = 0$.

Absolute Value

Definition

$$|z|^2 = z \cdot \bar{z}.$$

- ▶ $|z| \geq 0$, where $|z| = 0$ iff $z = 0$.
- ▶ $|z + w| \leq |z| + |w|$.

Absolute Value

Definition

$$|z|^2 = z \cdot \bar{z}.$$

- ▶ $|z| \geq 0$, where $|z| = 0$ iff $z = 0$.
- ▶ $|z + w| \leq |z| + |w|$.
- ▶ $|z \cdot w| = |z| \cdot |w|$.

Absolute Value


Definition

$$|z|^2 = z \cdot \bar{z}.$$

- ▶ $|z| \geq 0$, where $|z| = 0$ iff $z = 0$.
- ▶ $|z + w| \leq |z| + |w|$.
- ▶ $|z \cdot w| = |z| \cdot |w|$.

By Pythagoras' theorem, $|a + bi| = \sqrt{a^2 + b^2}$

Argument



Theorem

$$\arg(z_1 z_2) \equiv \arg z_1 + \arg z_2 \pmod{2\pi}.$$

Absolute value and argument

$$z = |z| \cdot (\cos \arg z + i \sin \arg z) .$$


Absolute value and argument

$$z = |z| \cdot (\cos \arg z + i \sin \arg z) .$$


$$zw = |z| \cdot |w| \cdot ((\cos \arg z \cos \arg w - \sin \arg z \sin \arg w) \\ + (\cos \arg z \sin \arg w + \sin \arg z \cos \arg w)i) .$$

$$zw = |z| \cdot |w| \cdot (\cos(\arg z + \arg w) + i \sin(\arg z + \arg w))$$


Complex numbers as vectors




Complex numbers as vectors




Complex numbers as vectors




Complex numbers as vectors




Complex numbers as vectors




Complex numbers as vectors




Complex numbers as vectors




Complex numbers as vectors




Complex numbers as vectors




Complex numbers as vectors




Complex numbers as vectors




Complex numbers as vectors




Complex numbers as vectors



Complex numbers as vectors



Complex numbers as vectors



Binomial theorem

$$z = r(\cos \theta + i \sin \theta).$$

Binomial theorem

$$z = r(\cos \theta + i \sin \theta).$$

$$z^n = r^n (\cos(n\theta) + i \sin(n\theta))$$

Binomial theorem

$$z = r(\cos \theta + i \sin \theta).$$

$$z^n = r^n (\cos(n\theta) + i \sin(n\theta))$$

$$\sqrt[n]{z} = \sqrt[n]{r} \left(\cos \frac{\theta}{n} + i \sin \frac{\theta}{n} \right)$$

Binomial theorem

$$z = r(\cos \theta + i \sin \theta).$$

$$z^n = r^n (\cos(n\theta) + i \sin(n\theta))$$

$$\sqrt[n]{z} = \sqrt[n]{r} \left(\cos \frac{\theta}{n} + i \sin \frac{\theta}{n} \right)$$

$$\sqrt[n]{z} = \sqrt[n]{r} \left(\cos \frac{\theta + k \cdot 2\pi}{n} + i \sin \frac{\theta + k \cdot 2\pi}{n} \right),$$

$$\text{for } k \in \{0, \dots, n-1\}.$$

Binomial theorem

$$z = r(\cos \theta + i \sin \theta).$$

$$z^n = r^n (\cos(n\theta) + i \sin(n\theta))$$


$$\sqrt[n]{z} = \sqrt[n]{r} \left(\cos \frac{\theta}{n} + i \sin \frac{\theta}{n} \right)$$

$$\sqrt[n]{z} = \sqrt[n]{r} \left(\cos \frac{\theta + k \cdot 2\pi}{n} + i \sin \frac{\theta + k \cdot 2\pi}{n} \right),$$


$$\text{for } k \in \{0, \dots, n-1\}.$$

So n^{th} roots exist; they form vertices of a regular polygon.


Roots of unity




Roots of unity




Roots of unity




Roots of unity



Roots of unity



Roots of unity



Powers of a complex number

We can now compute $z^{p/q}$.

By continuity, we could define z^x for $x \in \mathbb{R}$.

What about, say, z^i ? This will have to wait.

Stereographic projection

Extend \mathbb{C} to $\mathbb{C} \cup \{\infty\}$.

Consider $S^2 = \{(x_1, x_2, x_3) \in \mathbb{R}^3 : \sum_i x_i^2 = 1\}$,

Define $f : S^2 \rightarrow \mathbb{C} \cup \{\infty\}$ by

$$z = x + iy = f(x_1, x_2, x_3) = \frac{x_1 + ix_2}{1 - x_3}$$

The points $(0, 0, 1)$, $(x, y, 0)$, and (x_1, x_2, x_3) are collinear.

Powers on the Riemann sphere