

Topology of Piecewise-Linear Manifolds

Jim Fowler

Lecture 6
Summer 2010

Where are we now?



Where are we now?

Objects



Where are we now?

Objects
polyhedra



Where are we now?

Objects

polyhedra

simplicial complexes



Where are we now?

Objects

polyhedra

simplicial complexes

Maps



Where are we now?

Objects

polyhedra

simplicial complexes

Maps

PL maps



Where are we now?

Objects

polyhedra

simplicial complexes

Maps

PL maps

simplicial maps

Where are we now?

Objects

polyhedra

simplicial complexes

Maps

PL maps

simplicial maps

Invariants

Where are we now?

Objects

polyhedra

simplicial complexes

Maps

PL maps

simplicial maps

Invariants

χ

Where are we now?

Objects

polyhedra

simplicial complexes

Maps

PL maps

simplicial maps

Invariants

χ

b_0

Where are we now?

Objects

polyhedra

simplicial complexes

Maps

PL maps

simplicial maps

Invariants

χ

b_0

w_1

Piecewise linear map

Let P, Q be polyhedra.

$f : P \rightarrow Q$ is a PL map

if each point $p \in P$

has a closed star $N = p * L$

so that $f(\lambda p + (1 - \lambda)x) = \lambda f(p) + (1 - \lambda)f(x)$

for $x \in L$ and $\lambda \in [0, 1]$.

It locally maps conical rays to conical rays.

Convexity

Definition

A subset $C \in \mathbb{R}^n$ is **convex**
if for any $p, q \in C$,
the segment $\{p\} * \{q\}$
is contained in C .

Cells

A compact convex subset $C \in \mathbb{R}^n$
is a k -dimensional **cell**
if it spans a k -dimensional subspace.

Cells

A compact convex subset $C \in \mathbb{R}^n$
is a k -dimensional **cell**
if it spans a k -dimensional subspace.

For $x \in C$, define $\langle x, C \rangle$ to be
the union of $\{x\}$ and
all lines L such that $L \cap C$ contains x in its interior.

Cells

A compact convex subset $C \in \mathbb{R}^n$
is a k -dimensional **cell**
if it spans a k -dimensional subspace.

For $x \in C$, define $\langle x, C \rangle$ to be
the union of $\{x\}$ and
all lines L such that $L \cap C$ contains x in its interior.

The subset $C_x = C \cap \langle x, C \rangle$ is
the **face** of C containing x .

Cells

A compact convex subset $C \in \mathbb{R}^n$
is a k -dimensional **cell**
if it spans a k -dimensional subspace.

For $x \in C$, define $\langle x, C \rangle$ to be
the union of $\{x\}$ and
all lines L such that $L \cap C$ contains x in its interior.

The subset $C_x = C \cap \langle x, C \rangle$ is
the **face** of C containing x .

Write $D < C$ if D is a face of C .

Cell complex

A **cell complex** is

a finite collection K of cells such that

- ▶ If $C \in K$, and $D < C$, then $D \in K$.
- ▶ If $C, D \in K$, then $C \cap D$ is a face of C and D .

Cell complex

A **cell complex** is

a finite collection K of cells such that

- ▶ If $C \in K$, and $D < C$, then $D \in K$.
- ▶ If $C, D \in K$, then $C \cap D$ is a face of C and D .

The **underlying polyhedron**, $|K|$ is
the union of all cells in K .

Cell complex

A **cell complex** is

a finite collection K of cells such that

- ▶ If $C \in K$, and $D < C$, then $D \in K$.
- ▶ If $C, D \in K$, then $C \cap D$ is a face of C and D .

The **underlying polyhedron**, $|K|$ is
the union of all cells in K .

A **cellular map** $f : K \rightarrow L$ is
a PL map $|f| : |K| \rightarrow |L|$
which is linear on cells of K ,
and sends cells to cells.

Subdivision

A cell complex L is a **subdivision** of K
if $|L| = |K|$
and each cell of L is contained in a cell K .

Subdivision

A cell complex L is a **subdivision** of K
if $|L| = |K|$
and each cell of L is contained in a cell K .

Write $L \triangleleft K$ if L is a subdivision of K .

Simplicial complexes, not abstract

A cell complex is a **simplicial complex** if each $C \in K$ is a **simplex** (i.e., an n -cell which is the join of $n + 1$ independent points).

A **triangulation** of a polyhedron P is a simplicial complex K with a PL homeomorphism $f : |K| \rightarrow P$.

From now on,
complex
means
simplicial complex,
(not abstract)



Simplicial Collapse

Principal Simplexes

Definition

Let K be a complex, and
 $\sigma \in K$ a simplex.

Call σ a **principal simplex**

if the only simplex containing σ
is σ itself

(i.e., it isn't contained in a larger simplex).

Principal Simplexes

Definition

Let K be a complex, and $\sigma \in K$ a simplex.

Call σ a **principal simplex** if the only simplex containing σ is σ itself

(i.e., it isn't contained in a larger simplex).

Problem

Does every complex have a principal simplex?

Free faces

Definition

Let K be a complex,
and $\sigma \in K$ a simplex,
and $\tau < \sigma$ a face.

Call τ a **free face** of σ
if the only simplexes containing τ
are τ and σ .

Free faces

Definition

Let K be a complex,
and $\sigma \in K$ a simplex,
and $\tau < \sigma$ a face.

Call τ a **free face** of σ
if the only simplexes containing τ
are τ and σ .

Problem

Does every complex have a simplex with a free face?

Free faces

Definition

Let K be a complex,
and $\sigma \in K$ a simplex,
and $\tau < \sigma$ a face.

Call τ a **free face** of σ
if the only simplexes containing τ
are τ and σ .

Problem

Does every complex have a simplex with a free face?

Does any complex have a simplex with a free face?

Elementary simplicial collapse

Definition

Let L and $K = L \cup \text{cl}\{\sigma, \tau\}$ be complexes

If σ is a principal simplex of K , and

τ is a free face of σ , then

L is an **elementary simplicial collapse** of K .

Simplicial collapse

Definition

Let K_1, K_2, \dots, K_n be complexes, with K_{i+1} an elementary simplicial collapse of K_i .

Call K_n a **simplicial collapse** of K_1 , and write $K_1 \searrow K_n$.

Call K_1 a **simplicial expansion** of K_n , and write $K_n \nearrow K_1$.

Simple homotopy equivalence

Definition

K is **simple homotopy equivalent** to L
(sometimes abbreviated s.h.e.)

if you can reach transform K into L
via a sequence of

- ▶ PL homeomorphisms,
- ▶ simplicial collapses,
- ▶ simplicial expansions.

In this case, we write $K \searrow \nearrow L$.

Full subcomplex

$L \subset K$ are complexes.

Define $f_L : K \rightarrow [0, 1]$ on vertices by

$$f_L(v) = \begin{cases} 0 & \text{if } v \in L \\ 1 & \text{if } v \notin L \end{cases}$$

and extending linearly to simplexes.

If $L = f_L^{-1}(0)$

we say L is a **full subcomplex** of K ,
and write $L \subseteq K$.

Derived subdivision

$L \subset K$ are complexes.

Choose $a_i \in \text{int } A_i$ for each $A_i \in K$ with $A_i \notin L$.

$K' \triangleleft K$ is “ K derived away from L ”
is defined inductively over skeleta by

$$A_i' = \begin{cases} \{a_i\} * \partial A_i' & \text{if } A_i \notin L, \\ A_i & \text{if } A_i \in L. \end{cases}$$

$L \subset K$ are complexes.

Define the simplicial neighborhood of L in K as

$$N(L, K) = \{A \in K : A < B, B \cap |L| \neq \emptyset\}.$$

The **simplicial complement** of L in K is

$$C(L, K) = \{A \in K : A \cap |L| = \emptyset\}.$$

$L \subset K$ are complexes.

Define the simplicial neighborhood of L in K as

$$N(L, K) = \{A \in K : A < B, B \cap |L| \neq \emptyset\}.$$

The **simplicial complement** of L in K is

$$C(L, K) = \{A \in K : A \cap |L| = \emptyset\}.$$

$$K = N(L, K) \cup C(L, K)$$

$L \subset K$ are complexes.

Define the simplicial neighborhood of L in K as

$$N(L, K) = \{A \in K : A < B, B \cap |L| \neq \emptyset\}.$$

The **simplicial complement** of L in K is

$$C(L, K) = \{A \in K : A \cap |L| = \emptyset\}.$$

$$K = N(L, K) \cup C(L, K)$$

$$\partial N(L, K) := N(L, K) \cap C(L, K)$$

A derived of K near L
is obtained by
deriving K away from $L \cup C(L, K)$,
i.e., deriving simplexes meeting $|L|$ but not in L .

A derived of K near L
is obtained by
deriving K away from $L \cup C(L, K)$,
i.e., deriving simplexes meeting $|L|$ but not in L .

$L \in K'$ if K' is K derived near L .

Why?

If K' is K derived near L ,
 $N(L, K')$ is a
derived neighborhood of L in K .

Theorem

*If K_1 and K_2 are deriveds of K near L ,
then $s : K_1 \rightarrow K_2$
throws $N(L, K_1)$ onto $N(L, K_2)$
and is the identity on $L \cup C(L, K)$.*

Regular neighborhoods

Suppose $X \subset Y$ are polyhedra.

$|K|$ a neighborhood of X in Y ,

$|L| = X$

$L \subseteq K$

K' derived of K near L .

$|N(L, K')|$ is called a **regular neighborhood**

Theorem

*If N_1 and N_2 are regular neighborhoods of X in Y ,
then there is a homeomorphism $h : Y \rightarrow Y$
which throws N_1 onto N_2 , and
which is the identity on X
and the identity outside a compact subset of Y .*

Regular neighborhoods in manifolds

Theorem

A regular neighborhood N of a polyhedron X in a manifold M is a manifold with boundary.

Theorem (Simplicial neighborhood theorem)

X a compact polyhedron

M a manifold

$X \subset \text{int } M$

N a neighborhood of X in $\text{int } M$.

Then N is a regular neighborhood iff

- ▶ *N is a compact manifold with boundary*
- ▶ *there are triangulations (K, L, J) of $(N, X, \partial N)$ with $L \subseteq K$, $K = N(L, K)$ and $J = \partial N(L, K)$.*

Theorem (SNT version 2)

X a compact polyhedron

M a manifold

$X \subset \text{int } M$

N a polyhedral neighborhood of X in $\text{int } M$.

Then N is a regular neighborhood iff

- ▶ N is a compact manifold with boundary
- ▶ $N \searrow X$.

Theorem (SNT version 2)

X a compact polyhedron

M a manifold

$X \subset \text{int } M$

N a polyhedral neighborhood of X in $\text{int } M$.

Then N is a regular neighborhood iff

- ▶ N is a compact manifold with boundary
- ▶ $N \searrow X$.

e.g., if $X \searrow$ point, N is a ball.

Theorem (SNT version 2)

X a compact polyhedron

M a manifold

$X \subset \text{int } M$

N a polyhedral neighborhood of X in $\text{int } M$.

Then N is a regular neighborhood iff

- ▶ N is a compact manifold with boundary
- ▶ $N \searrow X$.

e.g., if $X \searrow$ point, N is a ball.

e.g., a collapsible manifold is a ball.

Corollary

*If $B^n \subset S^n$, then
 $\text{cl}(S^n - B^n)$ is a ball.*

Corollary

*If $B^n \subset S^n$, then
 $\text{cl}(S^n - B^n)$ is a ball.*

Proof.

$$S^n = \partial\Delta^{n+1}.$$

A point $p \in B^n$ is contained in some $\Delta^n \subset \Delta^{n+1}$,
and both Δ^n and B are regular neighborhoods of P .
Regular neighborhoods are unique. □

next time
we will introduce the work of

E. C. Zeeman