

# Lecture 24: Removable singularities

Math 660—Jim Fowler

Monday, July 25, 2011

# Removable singularities

$f : \Omega - \{a\} \rightarrow \mathbb{C}$  analytic  
can be extended to  $\Omega \rightarrow \mathbb{C}$   
iff  $\lim_{z \rightarrow a} f(z)(z - a) = 0$ .

# Removable singularities

$f : \Omega - \{a\} \rightarrow \mathbb{C}$  analytic  
can be extended to  $\Omega \rightarrow \mathbb{C}$   
iff  $\lim_{z \rightarrow a} f(z)(z - a) = 0$ .

Why? Cauchy's formula is valid.

Apply this trick to

$$F(z) = \frac{f(z) - f(a)}{z - a}$$

and get  $f(z) = f(a) + (z - a)F(z)$ .

Apply this trick to

$$F(z) = \frac{f(z) - f(a)}{z - a}$$

and get  $f(z) = f(a) + (z - a)F(z)$ .  
Rinse, repeat.

## Consequently . . .

If  $f : \Omega \rightarrow \mathbb{C}$  is analytic,  
for any  $a \in \Omega$ , we can write

$$f(z) = f(a) + \sum_{n=1}^k \frac{f^{(n)}(a)}{n!} (z - a)^n + F(z)(z - a)^{k+1}$$

for some analytic  $F : \Omega \rightarrow \mathbb{C}$ .

## Say more about $F$

If  $f : \Omega \rightarrow \mathbb{C}$  is analytic,  
for any  $a \in \Omega$ , we can write

$$f(z) = f(a) + \sum_{n=1}^k \frac{f^{(n)}(a)}{n!} (z - a)^n + F(z)(z - a)^{k+1}$$

for the analytic  $F : \Omega \rightarrow \mathbb{C}$ .

$$F(z) = \frac{1}{2\pi i \eta(z, \gamma)} \int_{\gamma} \frac{f(w)}{(w - a)^n (w - z)} dw.$$

# Zeroes and poles

If all derivatives of  $f$  vanish at  $a$ ,  
then  $f(z) = F(z)(z - a)^n$

# Zeroes and poles

If all derivatives of  $f$  vanish at  $a$ ,  
then  $f(z) = F(z)(z - a)^n$   
Estimate the size of  $F(z)$ .

# Zeroes and poles

If all derivatives of  $f$  vanish at  $a$ ,

$$\text{then } f(z) = F(z)(z - a)^n$$

Estimate the size of  $F(z)$ .

$f(z)$  vanishes on  $\Omega$ .

## Zero of order $k$

If  $f^{(n)}(a) = 0$  for  $n \in \{0, \dots, k - 1\}$ ,  
but  $f^{(k)}(a) \neq 0$ ,  
we say  $f$  has a **zero of order  $k$** .

## Zero of order $k$

If  $f^{(n)}(a) = 0$  for  $n \in \{0, \dots, k - 1\}$ ,  
but  $f^{(k)}(a) \neq 0$ ,  
we say  $f$  has a **zero of order  $k$** .  
It's like a polynomial!

## Zeroes are isolated

Zeroes of analytic function are isolated.

## Zeroes are isolated

Zeroes of analytic function are isolated.

So if  $f(z) = g(z)$  for  $z \in S$ ,  
(what sort of set is  $S$ ?)

then  $f \equiv g$  on  $\Omega$ .

# Classifying singularities

$\Delta$  a disk, with center  $a$ .

Suppose  $f : \Delta - \{a\} \rightarrow \mathbb{C}$  is analytic.

# Classifying singularities

$\Delta$  a disk, with center  $a$ .

Suppose  $f : \Delta - \{a\} \rightarrow \mathbb{C}$  is analytic.

If  $\lim_{z \rightarrow a} f(z) = \infty$ ,

we call  $a$  a **pole**.

Consider  $1/f(z)$ ,

which has a zero at  $a$  of order  $k$ ,

now called the order of the pole.

# Classifying singularities

$\Delta$  a disk, with center  $a$ .

Suppose  $f : \Delta - \{a\} \rightarrow \mathbb{C}$  is analytic.

If  $\lim_{z \rightarrow a} f(z) = \infty$ ,

we call  $a$  a **pole**.

Consider  $1/f(z)$ ,

which has a zero at  $a$  of order  $k$ ,

now called the order of the pole.

**Meromorphic** means holomorphic except for poles.

# Meromorphic functions

# Meromorphic functions

Quotients of analytic functions

# Meromorphic functions

Quotients of analytic functions

Poles of meromorphic functions are isolated

# Meromorphic functions

Quotients of analytic functions

Poles of meromorphic functions are isolated

Think of the relationship between  $\mathbb{Z}$  and  $\mathbb{Q}$ .

$$Z = \{\alpha \in \mathbb{R} : \lim_{z \rightarrow a} |(z-a)^\alpha f(z)| = 0\}$$

$$P = \{\alpha \in \mathbb{R} : \lim_{z \rightarrow a} |(z-a)^\alpha f(z)| = \infty\}$$

$$Z = \{\alpha \in \mathbb{R} : \lim_{z \rightarrow a} |(z - a)^\alpha f(z)| = 0\}$$

$$P = \{\alpha \in \mathbb{R} : \lim_{z \rightarrow a} |(z - a)^\alpha f(z)| = \infty\}$$

If  $\alpha \in Z$  and  $\beta > \alpha$ , then  $\beta \in Z$ .

If  $\alpha \in P$  and  $\beta < \alpha$ , then  $\beta \in P$ .

$$Z = \{\alpha \in \mathbb{R} : \lim_{z \rightarrow a} |(z - a)^\alpha f(z)| = 0\}$$

$$P = \{\alpha \in \mathbb{R} : \lim_{z \rightarrow a} |(z - a)^\alpha f(z)| = \infty\}$$

If  $\alpha \in Z$  and  $\beta > \alpha$ , then  $\beta \in Z$ .

If  $\alpha \in P$  and  $\beta < \alpha$ , then  $\beta \in P$ .

- ▶  $Z = \mathbb{R}$  and  $P = \emptyset$ ,
- ▶  $Z = (n, \infty)$  and  $P = (-\infty, n)$ ,
- ▶  $Z = \emptyset$  and  $P = \emptyset$ .

## Partial fractions redux

If  $f(z)$  has a pole of order  $k$ , consider

$$(z-a)^k f(z) = a_0 + a_1(z-a) + \cdots + a_{k-1}(z-a)^{k-1} + F(z)(z-a)^k$$

and divide by  $(z - a)^k$  to get the **singular part**.

$Z = \emptyset$  and  $P = \emptyset$

These are **essential singularities**.

Theorem (Weierstrass)

$f : \Omega - \{a\} \rightarrow \mathbb{C}$  analytic  
with essential singularity at  $a$ .

For every  $w \in \mathbb{C}$  and  $\epsilon > 0$ ,  
there is  $z \in B_\epsilon(a) \cap \Omega$   
so that  $f(z) = w$ .

# Singularities at infinity

To determine the behavior of  $f(z)$  at infinity,  
consider instead  $f(1/z)$ .

# Singularities at infinity

To determine the behavior of  $f(z)$  at infinity,  
consider instead  $f(1/z)$ .

What sort of singularity does  $\sin z$  have near infinity?



