

# Lecture 39: Schwarz' theorem

Math 660—Jim Fowler

Friday, August 13, 2010

# Poisson integral

Given a continuous function  $U : [0, 2\pi] \rightarrow \mathbb{R}$ , consider

$$P_*(z) = \frac{1}{2\pi} \int_0^{2\pi} \operatorname{re} \frac{e^{i\theta} + z}{e^{i\theta} - z} U(\theta) d\theta.$$

# Poisson integral

Given a continuous function  $U : [0, 2\pi] \rightarrow \mathbb{R}$ , consider

$$P_*(z) = \frac{1}{2\pi} \int_0^{2\pi} \operatorname{re} \frac{e^{i\theta} + z}{e^{i\theta} - z} U(\theta) d\theta.$$

►  $P_{U+V} = P_U + P_V$

# Poisson integral

Given a continuous function  $U : [0, 2\pi] \rightarrow \mathbb{R}$ , consider

$$P_*(z) = \frac{1}{2\pi} \int_0^{2\pi} \operatorname{re} \frac{e^{i\theta} + z}{e^{i\theta} - z} U(\theta) d\theta.$$

- ▶  $P_{U+V} = P_U + P_V$
- ▶  $P_{kU} = kP_U$

# Poisson integral

Given a continuous function  $U : [0, 2\pi] \rightarrow \mathbb{R}$ , consider

$$P_*(z) = \frac{1}{2\pi} \int_0^{2\pi} \operatorname{re} \frac{e^{i\theta} + z}{e^{i\theta} - z} U(\theta) d\theta.$$

- ▶  $P_{U+V} = P_U + P_V$
- ▶  $P_{kU} = kP_U$
- ▶  $P_c = c$

# Poisson integral

Given a continuous function  $U : [0, 2\pi] \rightarrow \mathbb{R}$ , consider

$$P_*(z) = \frac{1}{2\pi} \int_0^{2\pi} \operatorname{re} \frac{e^{i\theta} + z}{e^{i\theta} - z} U(\theta) d\theta.$$

- ▶  $P_{U+V} = P_U + P_V$
- ▶  $P_{kU} = kP_U$
- ▶  $P_c = c$
- ▶  $P_U \geq 0$  if  $U \geq 0$ .

# Schwarz' theorem

The function  $P_U(z)$  is harmonic in the interior of the disk  $B_1(0)$ ,  
and continuous on the closed disk (provided  $U$  is continuous).

# Reflection principle



# Reflection principle

If  $u(z)$  is harmonic, then  $u(\bar{z})$  is harmonic.

# Reflection principle

If  $u(z)$  is harmonic, then  $u(\bar{z})$  is harmonic.

If  $f(z)$  is analytic, then  $\overline{f(\bar{z})}$  is analytic.

# Reflection principle

If  $u(z)$  is harmonic, then  $u(\bar{z})$  is harmonic.

If  $f(z)$  is analytic, then  $\overline{f(\bar{z})}$  is analytic.

Consider  $f(z) - \overline{f(\bar{z})}$ .

Let  $\Omega^+$  be the part in the upper half plane of a symmetric region  $\Omega$ ,  
let  $\sigma$  be the part of the real axis in  $\Omega$ .

Let  $\Omega^+$  be the part in the upper half plane of a symmetric region  $\Omega$ ,  
let  $\sigma$  be the part of the real axis in  $\Omega$ .

Suppose  $v(x)$  is continuous in  $\Omega^+ \cup \sigma$ , harmonic in  $\Omega^+$ , and zero on  $\sigma$ .

Let  $\Omega^+$  be the part in the upper half plane of a symmetric region  $\Omega$ ,  
let  $\sigma$  be the part of the real axis in  $\Omega$ .

Suppose  $v(x)$  is continuous in  $\Omega^+ \cup \sigma$ , harmonic in  $\Omega^+$ , and zero on  $\sigma$ .

Then  $v$  has a harmonic extension to  $\Omega$  which satisfies the symmetry relation  $v(\bar{z}) = -v(z)$ .

Let  $\Omega^+$  be the part in the upper half plane of a symmetric region  $\Omega$ ,  
let  $\sigma$  be the part of the real axis in  $\Omega$ .

Suppose  $v(x)$  is continuous in  $\Omega^+ \cup \sigma$ , harmonic in  $\Omega^+$ , and zero on  $\sigma$ .

Then  $v$  has a harmonic extension to  $\Omega$  which satisfies the symmetry relation  $v(\bar{z}) = -v(z)$ .

In the same situation, if  $v$  is the imaginary part of an analytic function  $f(z)$  in  $\Omega^+$ , then  $f(z)$  has an analytic extension which satisfies  $f(z) = \overline{f(\bar{z})}$ .