

Lecture 37: More on Harmonic functions

Math 660—Jim Fowler

Wednesday, August 11, 2010

Harmonic functions in polar coordinates

Harmonic functions in polar coordinates

$$r \frac{\partial}{\partial r} \left(r \frac{\partial u}{\partial r} \right) + \frac{\partial^2 u}{\partial \theta^2} = 0$$

Harmonic functions in polar coordinates

$$r \frac{\partial}{\partial r} \left(r \frac{\partial u}{\partial r} \right) + \frac{\partial^2 u}{\partial \theta^2} = 0$$

So if $\frac{\partial u}{\partial \theta} = 0$, then $u = a \log r + b$.

Integrals of harmonic functions

If u is harmonic, with conjugate harmonic function v ,

Integrals of harmonic functions

If u is harmonic, with conjugate harmonic function v , then $f(x, y) = u(x, y) + i v(x, y)$ is analytic,

Integrals of harmonic functions

If u is harmonic, with conjugate harmonic function v , then $f(x, y) = u(x, y) + i v(x, y)$ is analytic, and we can find $\int f dz$.

Integrals of harmonic functions

If u is harmonic, with conjugate harmonic function v , then $f(x, y) = u(x, y) + i v(x, y)$ is analytic, and we can find $f dz$.

$$dv = -\frac{\partial u}{\partial y} dx + \frac{\partial u}{\partial x} dy.$$

Integrals of harmonic functions

If u is harmonic, with conjugate harmonic function v , then $f(x, y) = u(x, y) + i v(x, y)$ is analytic, and we can find $f dz$.

$$dv = -\frac{\partial u}{\partial y} dx + \frac{\partial u}{\partial x} dy.$$

But in practice, there may be no conjugate harmonic function,
so we take as a definition

$$*du = -\frac{\partial u}{\partial y} dx + \frac{\partial u}{\partial x} dy.$$

Hodge theory

For any cycle γ ,

$$\int_{\gamma} du = 0$$

Hodge theory

For any cycle γ ,

$$\int_{\gamma} du = 0$$

And if γ is homologous to zero, then

$$\int_{\gamma} f dz = 0$$

Hodge theory

For any cycle γ ,

$$\int_{\gamma} du = 0$$

And if γ is homologous to zero, then

$$\int_{\gamma} f dz = 0$$

And so $\int_{\gamma} *du = 0$ for cycles homologous to zero.

Simply connected regions

In a simply connected regions, $\int_{\gamma} *du = 0$ for all cycles, so u has a single-valued conjugate function v .

For every cycle γ homologous to zero,

$$\int_{\gamma} u_1 * du_2 - u_2 * du_1 = 0$$

For every cycle γ homologous to zero,

$$\int_{\gamma} u_1 * du_2 - u_2 * du_1 = 0$$

Proof: If conjugate harmonic functions exist (e.g., in a simply connected region like a rectangle), then

$$\begin{aligned} u_1 * du_2 - u_2 * du_1 &= u_1 dv_2 - u_2 dv_1 \\ &= u_1 dv_2 + v_1 du_2 - d(u_2 v_1) \\ &= (\operatorname{im}(u_1 + iv_1)(du_2 + idv_2)) - d(u_2 v_1) \end{aligned}$$

For every cycle γ homologous to zero,

$$\int_{\gamma} u_1 * du_2 - u_2 * du_1 = 0$$

Proof: If conjugate harmonic functions exist (e.g., in a simply connected region like a rectangle), then

$$\begin{aligned} u_1 * du_2 - u_2 * du_1 &= u_1 dv_2 - u_2 dv_1 \\ &= u_1 dv_2 + v_1 du_2 - d(u_2 v_1) \\ &= (\operatorname{im}(u_1 + iv_1)(du_2 + idv_2)) - d(u_2 v_1) \end{aligned}$$

so the integral vanishes.



For every cycle γ homologous to zero,

$$\int_{\gamma} u_1 * du_2 - u_2 * du_1 = 0$$

Apply this to $u_1 = \log r$ and $u_2 = u$.

Mean-value property

Suppose u is harmonic in a punctured disk; then the arithmetic mean of u over concentric circles is a linear function of $\log r$,

$$\frac{1}{2\pi} \int_{|z|=r} u \, d\theta = a \log r + \beta$$

and if u is harmonic in the whole disk, then $a = 0$.

Maximum principle

A nonconstant harmonic function has neither a maximum nor a minimum in the interior of a disk; the maximum and minimum on a closed bounded set occur on the boundary.

Poisson's formula

If $u(z)$ is continuous on a closed bounded set E ,
and harmonic in the interior of E ,
then it is uniquely determined by its values on the
boundary.

Poisson's formula

If $u(z)$ is continuous on a closed bounded set E ,
and harmonic in the interior of E ,
then it is uniquely determined by its values on the
boundary.

Can we get a formula for the function on the interior,
given the values on the boundary?

Recall

$$u(0) = \frac{1}{2\pi} \int_{|z|=r} u(re^{i\theta}) d\theta$$

Recall

$$u(0) = \frac{1}{2\pi} \int_{|z|=r} u(re^{i\theta}) d\theta$$

Consider $S_a(z) = \frac{R(Rz+a)}{R+\bar{a}z}$.

Recall

$$u(0) = \frac{1}{2\pi} \int_{|z|=r} u(re^{i\theta}) d\theta$$

Consider $S_a(z) = \frac{R(Rz+a)}{R+\bar{a}z}$.

$$u(a) = \frac{1}{2\pi} \int_{|z|=1} u(S_a(e^{i\theta})) d\theta.$$