

Topology of Piecewise-Linear Manifolds

Jim Fowler

Lecture 7
Summer 2010

Regular neighborhoods

Suppose $X \subset M$.

Regular neighborhoods

Suppose $X \subset M$. The polyhedron X might have many neighborhoods in M .

Regular neighborhoods

Suppose $X \subset M$. The polyhedron X might have many neighborhoods in M .

But there is an “essentially unique” regular neighborhood.

E. C. Zeeman

E. C. Zeeman

Goal: *“Unknotting combinatorial balls”*

E. C. Zeeman

Goal: *“Unknotting combinatorial balls”*

a paper in the **Annals**

Main result

Theorem

*Any embedding $S^p \subset S^q$
is unknotted, if $q - p \geq 3$.*

Main result

Theorem

*Any embedding $S^p \subset S^q$
is unknotted, if $q - p \geq 3$.*

Corollary

*Any embedding $S^1 \subset S^4$
is unknotted.*

Some background

Can S^1 be knotted in S^3 ?

Some background

Can S^1 be knotted in S^3 ?

What does *knotted* even mean?

One point compactification

$$\mathbb{R}^1 \cong S^1 - \text{point}$$

One point compactification

$$\mathbb{R}^1 \cong S^1 - \text{point}$$

$$\mathbb{R}^2 \cong S^2 - \text{point}$$

One point compactification

$$\mathbb{R}^1 \cong S^1 - \text{point}$$

$$\mathbb{R}^2 \cong S^2 - \text{point}$$

$$\mathbb{R}^3 \cong S^3 - \text{point}$$


One point compactification


$$\mathbb{R}^1 \cong S^1 - \text{point}$$


$$\mathbb{R}^2 \cong S^2 - \text{point}$$

$$\mathbb{R}^3 \cong S^3 - \text{point}$$

So we can draw $S^1 \subset S^3$ as if the circle were in \mathbb{R}^3 .








Unknot




3_1



4_1



5_1



5_2



6_1



6_2



6_3




7_1



7_2



7_3



7_4



7_5




7_6




7_7


Reidemeister Move—Type I




Reidemeister Move—Type II



Reidemeister Move—Type III



Reidemeister Moves



Knots

Theorem

Two knots are the same if a diagram for the one can be transformed into the other via Reidemeister moves.

High dimensional knots?

Are there “Reidemeister moves” for S^2 's in S^4 ?

Pairs

Definition

“(p, q)-sphere pair” means $S^q \subset S^p$.

We sometimes write (S^p, S^q) .

“(p, q)-ball pair” means $B^q \subset B^p$,
with B^q properly embedded in B^p ,
meaning $\partial B^q \subset \partial B^p$ and $\text{int } B^q \subset \text{int } B^p$.

We sometimes write (B^p, B^q) .

Pairs

Definition

“(p, q)-sphere pair” means $S^q \subset S^p$.

We sometimes write (S^p, S^q) .

“(p, q)-ball pair” means $B^q \subset B^p$,
with B^q properly embedded in B^p ,
meaning $\partial B^q \subset \partial B^p$ and $\text{int } B^q \subset \text{int } B^p$.

We sometimes write (B^p, B^q) .

“Pair” means either a ball-pair or sphere-pair.

Pairs

Definition

“(p, q)-sphere pair” means $S^q \subset S^p$.

We sometimes write (S^p, S^q) .

“(p, q)-ball pair” means $B^q \subset B^p$,
with B^q properly embedded in B^p ,
meaning $\partial B^q \subset \partial B^p$ and $\text{int } B^q \subset \text{int } B^p$.


We sometimes write (B^p, B^q) .

“Pair” means either a ball-pair or sphere-pair.

$$\partial(B^p, B^q) = (S^{p-1}, S^{q-1}).$$

Joins of pairs

$(S^p, S^q) * S^k$ is a



Joins of pairs

$(S^p, S^q) * S^k$ is a sphere pair.



Joins of pairs

$(S^p, S^q) * S^k$ is a sphere pair.

$(S^p, S^q) * B^k$ is a

Joins of pairs

$(S^p, S^q) * S^k$ is a sphere pair.

$(S^p, S^q) * B^k$ is a ball pair.

Joins of pairs

$(S^p, S^q) * S^k$ is a sphere pair.

$(S^p, S^q) * B^k$ is a ball pair.

$(B^p, B^q) * S^k$ is a

Joins of pairs

$(S^p, S^q) * S^k$ is a sphere pair.

$(S^p, S^q) * B^k$ is a ball pair.

$(B^p, B^q) * S^k$ is a ball pair.

Joins of pairs

$(S^p, S^q) * S^k$ is a sphere pair.

$(S^p, S^q) * B^k$ is a ball pair.

$(B^p, B^q) * S^k$ is a ball pair.

$(B^p, B^q) * B^k$ is a

Joins of pairs

$(S^p, S^q) * S^k$ is a sphere pair.

$(S^p, S^q) * B^k$ is a ball pair.

$(B^p, B^q) * S^k$ is a ball pair.

$(B^p, B^q) * B^k$ is a ball pair.

Joins of pairs

$(S^p, S^q) * S^k$ is a sphere pair.

$(S^p, S^q) * B^k$ is a ball pair.

$(B^p, B^q) * S^k$ is a ball pair.

$(B^p, B^q) * B^k$ is a ball pair.

Joins of pairs

$(S^p, S^q) * S^k$ is a sphere pair.

$(S^p, S^q) * B^k$ is a ball pair.

$(B^p, B^q) * S^k$ is a ball pair.

$(B^p, B^q) * B^k$ is a ball pair.

We'll call the join of a pair and a point a **cone pair**.

Subpairs

$X = (X^p, X^q)$ and $Y = (Y^r, Y^s)$ are pairs
we say Y is a subpair of X
(written $Y \subset X$ or $X \supset Y$)
if $Y^r \subset X^p$ and $T^s = X^q \cap Y^r$.

Subpairs

$X = (X^p, X^q)$ and $Y = (Y^r, Y^s)$ are pairs

we say Y is a subpair of X

(written $Y \subset X$ or $X \supset Y$)

if $Y^r \subset X^p$ and $T^s = X^q \cap Y^r$.

If $P = (S^p, S^q) \supset Q = (B^p, B^q)$, then

$$P - \text{int } Q = (S^p - \text{int } B^p, S^q - \text{int } B^q)$$

is a ball pair (via regular neighborhood machinery).

Faces of pairs

If $Q' = (B^{p-1}, B^{q-1})$ is contained in the boundary of $Q = (B^p, B^q)$, we call Q' a **face** of Q .

Faces of pairs

If $Q' = (B^{p-1}, B^{q-1})$ is contained in the boundary of $Q = (B^p, B^q)$, we call Q' a **face** of Q .

Theorem

If ball pairs intersect in their common boundary, their union is a sphere pair.

Faces of pairs

If $Q' = (B^{p-1}, B^{q-1})$ is contained in the boundary of $Q = (B^p, B^q)$, we call Q' a **face** of Q .

Theorem

If ball pairs intersect in their common boundary, their union is a sphere pair.

Theorem

If ball pairs intersect in a face, their union is a ball pair.

Standard pairs

$\Gamma^{p,q} = (S^{p-q}\Delta^q, \Delta^q)$ is the standard (p, q) -ball pair.
 $\partial\Gamma^{p+1,q+1}$ is the standard (p, q) -sphere pair.

Standard pairs

$\Gamma^{p,q} = (S^{p-q}\Delta^q, \Delta^q)$ is the standard (p, q) -ball pair.
 $\partial\Gamma^{p+1,q+1}$ is the standard (p, q) -sphere pair.


A pair is **unknotted**
if it is homeomorphic to a standard pair.

Theorem (**BallThm** _{p,q})

If $p - q \geq 3$, then any (p, q) -ball pair is unknotted.

Theorem (**SphereThm** _{p,q})

If $p - q \geq 3$, then any (p, q) -sphere pair is unknotted.



Theorem (**BallThm** _{p,q})

If $p - q \geq 3$, then any (p, q) -ball pair is unknotted.

Theorem (**SphereThm** _{p,q})

If $p - q \geq 3$, then any (p, q) -sphere pair is unknotted.

Proof.

By induction.



Overview of the induction

Prove **BallThm** _{$p,0$} by hand.

Overview of the induction

Prove **BallThm** _{$p,0$} by hand.

BallThm _{p,q} implies **SphereThm** _{p,q} .

Overview of the induction

Prove **BallThm** _{$p,0$} by hand.

BallThm _{p,q} implies **SphereThm** _{p,q} .

BallThm _{$p-1,q-1$} and **SphereThm** _{$p-1,q-1$}
together imply **BallThm** _{p,q} .

Base case

Lemma

BallThm _{$p,0$} *is true.*

Base case

Lemma

BallThm _{$p,0$} *is true.*

Proof.

A ball B^p with a marked point B^0
is homeomorphic
to any other such
(via regular neighborhood theory).



Overview of the induction

Prove **BallThm** _{$p,0$} by hand.

BallThm _{p,q} implies **SphereThm** _{p,q} .

BallThm _{$p-1,q-1$} and **SphereThm** _{$p-1,q-1$}
together imply **BallThm** _{p,q} .

Overview of the induction

✓ Prove **BallThm** _{$p,0$} by hand.

BallThm _{p,q} implies **SphereThm** _{p,q} .

BallThm _{$p-1,q-1$} and **SphereThm** _{$p-1,q-1$}
together imply **BallThm** _{p,q} .

Lemma

$$\mathbf{BallThm}_{p,q} \Rightarrow \mathbf{SphereThm}_{p,q}$$

Lemma

BallThm $_{p,q} \Rightarrow$ **SphereThm** $_{p,q}$

Proof.

$P = (S^p, S^q)$. Choose vertex $x \in S^q$.

Lemma

$$\mathbf{BallThm}_{p,q} \Rightarrow \mathbf{SphereThm}_{p,q}$$

Proof.

$P = (S^p, S^q)$. Choose vertex $x \in S^q$.

$$P = Q \cup \{x\} * \partial Q,$$

Lemma

$$\mathbf{BallThm}_{p,q} \Rightarrow \mathbf{SphereThm}_{p,q}$$

Proof.

$P = (S^p, S^q)$. Choose vertex $x \in S^q$.

$$P = Q \cup \{x\} * \partial Q,$$

where $Q = (S^p - \text{st}(x, S^p), S^q - \text{st}(x, S^q))$.

Lemma

$$\mathbf{BallThm}_{p,q} \Rightarrow \mathbf{SphereThm}_{p,q}$$

Proof.

$P = (S^p, S^q)$. Choose vertex $x \in S^q$.

$$P = Q \cup \{x\} * \partial Q,$$

where $Q = (S^p - \text{st}(x, S^p), S^q - \text{st}(x, S^q))$.

Extend $Q \cong \Gamma^{p,q}$ to

a homeomorphism $P \cong \partial \Gamma^{p+1,q+1}$.



Overview of the induction

✓ Prove **BallThm** _{$p,0$} by hand.

BallThm _{p,q} implies **SphereThm** _{p,q} .

BallThm _{$p-1,q-1$} and **SphereThm** _{$p-1,q-1$}
together imply **BallThm** _{p,q} .

Overview of the induction

- ✓ Prove **BallThm** _{$p,0$} by hand.
- ✓ **BallThm** _{p,q} implies **SphereThm** _{p,q} .
BallThm _{$p-1,q-1$} and **SphereThm** _{$p-1,q-1$}
together imply **BallThm** _{p,q} .

The last step

BallThm _{$p-1,q-1$} and **SphereThm** _{$p-1,q-1$}



BallThm _{p,q}

This will require more machinery,
building on simplicial collapse
and regular neighborhoods.

BallThm _{$p-1, q-1$} and **SphereThm** _{$p-1, q-1$}



BallThm _{p, q}

BallThm _{$p-1, q-1$} and **SphereThm** _{$p-1, q-1$}



BallThm _{p, q}

Lemma

Assuming **BallThm** _{$p-1, q-1$} and **SphereThm** _{$p-1, q-1$} ,
 (B^p, B^q) with $p - q \geq 3$ is unknotted
provided $B^p \searrow B^q$.

BallThm _{$p-1, q-1$} and **SphereThm** _{$p-1, q-1$}



BallThm _{p, q}

Lemma

Assuming **BallThm** _{$p-1, q-1$} and **SphereThm** _{$p-1, q-1$} ,
 (B^p, B^q) with $p - q \geq 3$ is unknotted
provided $B^p \searrow B^q$.

Lemma

If $p - q \geq 3$ and (B^p, B^q) is any ball pair,
then $B^p \searrow B^q$.

But first...

Before we can proceed,
we will prove a couple of helpful lemmas.

Lemma

Q_1 and Q_2 are unknotted (p, q) -ball pairs.

Any homeomorphism $h : \partial Q_1 \xrightarrow{\cong} \partial Q_2$
extends to a homeomorphism $h' : Q_1 \xrightarrow{\cong} Q_2$.

Lemma

Q_1 and Q_2 are unknotted (p, q) -ball pairs.

Any homeomorphism $h : \partial Q_1 \xrightarrow{\cong} \partial Q_2$
extends to a homeomorphism $h' : Q_1 \xrightarrow{\cong} Q_2$.

Proof (via *Alexander trick*).

y = point in interior of Δ^q ;

then since Q_i is unknotted, we have maps

$$f_i : Q_i \xrightarrow{\cong} \{y\} * \partial \Gamma^{p,q}$$

Lemma

Q_1 and Q_2 are unknotted (p, q) -ball pairs.

Any homeomorphism $h : \partial Q_1 \xrightarrow{\cong} \partial Q_2$
extends to a homeomorphism $h' : Q_1 \xrightarrow{\cong} Q_2$.

Proof (via Alexander trick).

y = point in interior of Δ^q ;

then since Q_i is unknotted, we have maps

$$f_i : Q_i \xrightarrow{\cong} \{y\} * \partial \Gamma^{p,q}$$

$$g : \partial \Gamma^{p,q} \xrightarrow{f_1^{-1}} \partial Q_1 \xrightarrow{h} \partial Q_2 \xrightarrow{f_2} \partial \Gamma^{p,q}$$

Lemma

Q_1 and Q_2 are unknotted (p, q) -ball pairs.

Any homeomorphism $h : \partial Q_1 \xrightarrow{\cong} \partial Q_2$
extends to a homeomorphism $h' : Q_1 \xrightarrow{\cong} Q_2$.

Proof (via *Alexander trick*).

y = point in interior of Δ^q ;

then since Q_i is unknotted, we have maps

$$f_i : Q_i \xrightarrow{\cong} \{y\} * \partial \Gamma^{p,q}$$

$$g : \partial \Gamma^{p,q} \xrightarrow{f_1^{-1}} \partial Q_1 \xrightarrow{h} \partial Q_2 \xrightarrow{f_2} \partial \Gamma^{p,q}$$

$$h' : Q_1 \xrightarrow{f_1} \{y\} * \partial \Gamma^{p,q} \xrightarrow{\text{cone } g} \{y\} * \partial \Gamma^{p,q} \xrightarrow{f_2^{-1}} Q_2 \quad \square$$

In short, **radial extension**.

Lemma (assume **BallThm** _{$p-1, q-1$})

*If Q_1, Q_2 are unknotted (p, q) -ball pairs,
and $Q_3 = Q_1 \cap Q_2$ is a face,
then $Q_1 \cup Q_2$ is unknotted.*

Lemma (assume **BallThm** _{$p-1, q-1$})

*If Q_1, Q_2 are unknotted (p, q) -ball pairs,
and $Q_3 = Q_1 \cap Q_2$ is a face,
then $Q_1 \cup Q_2$ is unknotted.*

Proof.

Choose $Q_3 \xrightarrow{\cong} \Gamma^{p-1, q-1}$, extend over $\partial Q_1 - \text{int } Q_3$
to $h : \partial Q_1 \xrightarrow{\cong} \Gamma^{p-1, q-1} \cup C\partial\Gamma^{p-1, q-1}$,

Lemma (assume **BallThm** _{$p-1, q-1$})

*If Q_1, Q_2 are unknotted (p, q) -ball pairs,
and $Q_3 = Q_1 \cap Q_2$ is a face,
then $Q_1 \cup Q_2$ is unknotted.*

Proof.

Choose $Q_3 \xrightarrow{\cong} \Gamma^{p-1, q-1}$, extend over $\partial Q_1 - \text{int } Q_3$
to $h : \partial Q_1 \xrightarrow{\cong} \Gamma^{p-1, q-1} \cup C\partial\Gamma^{p-1, q-1}$,
 h extends to $Q_1 \rightarrow C\Gamma^{p-1, q-1}$

Lemma (assume **BallThm** _{$p-1, q-1$})

*If Q_1, Q_2 are unknotted (p, q) -ball pairs,
and $Q_3 = Q_1 \cap Q_2$ is a face,
then $Q_1 \cup Q_2$ is unknotted.*

Proof.

Choose $Q_3 \xrightarrow{\cong} \Gamma^{p-1, q-1}$, extend over $\partial Q_1 - \text{int } Q_3$
to $h : \partial Q_1 \xrightarrow{\cong} \Gamma^{p-1, q-1} \cup C\partial\Gamma^{p-1, q-1}$,

h extends to $Q_1 \rightarrow C\Gamma^{p-1, q-1}$

Similarly produce $Q_2 \rightarrow C\Gamma^{p-1, q-1}$

Lemma (assume **BallThm** _{$p-1, q-1$})

*If Q_1, Q_2 are unknotted (p, q) -ball pairs,
and $Q_3 = Q_1 \cap Q_2$ is a face,
then $Q_1 \cup Q_2$ is unknotted.*

Proof.

Choose $Q_3 \xrightarrow{\cong} \Gamma^{p-1, q-1}$, extend over $\partial Q_1 - \text{int } Q_3$
to $h : \partial Q_1 \xrightarrow{\cong} \Gamma^{p-1, q-1} \cup C\partial\Gamma^{p-1, q-1}$,

h extends to $Q_1 \rightarrow C\Gamma^{p-1, q-1}$

Similarly produce $Q_2 \rightarrow C\Gamma^{p-1, q-1}$

Glue together $Q_1 \cup Q_2 \cong S\Gamma^{p-1, q-1}$.



Regular neighborhoods

M , an n -manifold, $X \subset M$ a polyhedron
a **regular neighborhood** of X in M
is a subpolyhedron $N \subset M$ such that

- ▶ N is a closed neighborhood of X
- ▶ N is an n -manifold
- ▶ $N \searrow X$.

Theorem

*If N_1 and N_2 are regular neighborhoods of $X \subset M$,
there's a homeomorphism $N_1 \rightarrow N_2$ keeping X fixed.*

Lemma

Assuming **BallThm** _{$p-1, q-1$} and **SphereThm** _{$p-1, q-1$} ,
 (B^p, B^q) with $p - q \geq 3$ is unknotted
provided $B^p \searrow B^q$.

Lemma

Assuming **BallThm** _{$p-1, q-1$} and **SphereThm** _{$p-1, q-1$} ,
 (B^p, B^q) with $p - q \geq 3$ is unknotted
provided $B^p \searrow B^q$.

Warning: The lemma is false if $p - q = 2$.

If $(B^4, B^2) = \text{cone}(S^3, S^1)$,

then $B^4 \searrow B^2$ because cones collapse to a subcone

Lemma

$B^p \searrow B^q \Rightarrow (B^p, B^q) \text{ unknotted}$

Lemma

$B^p \searrow B^q \Rightarrow (B^p, B^q) \text{ unknotted}$

Pick regular neighborhood N of B^q ;

Lemma

$B^p \searrow B^q \Rightarrow (B^p, B^q) \text{ unknotted}$

Pick regular neighborhood N of B^q ;
then $(B^p, B^q) \cong (N, B^q)$.

Lemma

$B^p \searrow B^q \Rightarrow (B^p, B^q) \text{ unknotted}$

Pick regular neighborhood N of B^q ;
then $(B^p, B^q) \cong (N, B^q)$.

$$B^q = K_k \searrow K_{k-1} \searrow \cdots \searrow K_0 = \{x\}$$

Lemma

$B^p \searrow B^q \Rightarrow (B^p, B^q)$ *unknotted*

Pick regular neighborhood N of B^q ;

then $(B^p, B^q) \cong (N, B^q)$.

$B^q = K_k \searrow K_{k-1} \searrow \cdots \searrow K_0 = \{x\}$

Q_i = simplicial neighborhood of K_i in (B^p, B^q) .

Lemma

$B^p \searrow B^q \Rightarrow (B^p, B^q)$ *unknotted*

Pick regular neighborhood N of B^q ;

then $(B^p, B^q) \cong (N, B^q)$.

$B^q = K_k \searrow K_{k-1} \searrow \cdots \searrow K_0 = \{x\}$

Q_i = simplicial neighborhood of K_i in (B^p, B^q) .

Proceed by induction.

Lemma

$B^p \searrow B^q \Rightarrow (B^p, B^q)$ *unknotted*

Pick regular neighborhood N of B^q ;

then $(B^p, B^q) \cong (N, B^q)$.

$$B^q = K_k \searrow K_{k-1} \searrow \cdots \searrow K_0 = \{x\}$$

Q_i = simplicial neighborhood of K_i in (B^p, B^q) .

Proceed by induction.

$$Q_0 = \{x\} * L \text{ where } L = (\text{lk}(x, B^p), \text{lk}(x, B^q)),$$

Lemma

$B^p \searrow B^q \Rightarrow (B^p, B^q)$ *unknotted*

Pick regular neighborhood N of B^q ;

then $(B^p, B^q) \cong (N, B^q)$.

$B^q = K_k \searrow K_{k-1} \searrow \cdots \searrow K_0 = \{x\}$

Q_i = simplicial neighborhood of K_i in (B^p, B^q) .

Proceed by induction.

$Q_0 = \{x\} * L$ where $L = (\text{lk}(x, B^p), \text{lk}(x, B^q))$,
and L is unknotted by either **BallThm** _{$p-1, q-1$} or
SphereThm _{$p-1, q-1$} .

Suppose Q_{i-1} is unknotted.

Suppose Q_{i-1} is unknotted.

$K_i - K_{i-1}$ consists of a principal simplex A with a free face C .

Suppose Q_{i-1} is unknotted.

$K_i - K_{i-1}$ consists of a principal simplex A with a free face C .

Pick $a \in \text{int } A$ and $c \in \text{int } C$.

Suppose Q_{i-1} is unknotted.

$K_i - K_{i-1}$ consists of a principal simplex A with a free face C .

Pick $a \in \text{int } A$ and $c \in \text{int } C$.

$Q_a = \{a\} * (\text{lk}(a, B^p), \text{lk}(a, B^q))$ is unknotted.

Suppose Q_{i-1} is unknotted.

$K_i - K_{i-1}$ consists of a principal simplex A with a free face C .

Pick $a \in \text{int } A$ and $c \in \text{int } C$.

$Q_a = \{a\} * (\text{lk}(a, B^p), \text{lk}(a, B^q))$ is unknotted.

$Q_c = \{c\} * (\text{lk}(c, B^p), \text{lk}(c, B^q))$ is unknotted.

Suppose Q_{i-1} is unknotted.

$K_i - K_{i-1}$ consists of a principal simplex A with a free face C .

Pick $a \in \text{int } A$ and $c \in \text{int } C$.

$Q_a = \{a\} * (\text{lk}(a, B^p), \text{lk}(a, B^q))$ is unknotted.

$Q_c = \{c\} * (\text{lk}(c, B^p), \text{lk}(c, B^q))$ is unknotted.

$Q_i = Q_{i-1} \cup Q_a \cup Q_c$ is union of unknotted ball pairs along faces; therefore, Q_i is unknotted.

Suppose Q_{i-1} is unknotted.

$K_i - K_{i-1}$ consists of a principal simplex A with a free face C .

Pick $a \in \text{int } A$ and $c \in \text{int } C$.

$Q_a = \{a\} * (\text{lk}(a, B^p), \text{lk}(a, B^q))$ is unknotted.

$Q_c = \{c\} * (\text{lk}(c, B^p), \text{lk}(c, B^q))$ is unknotted.

$Q_i = Q_{i-1} \cup Q_a \cup Q_c$ is union of unknotted ball pairs along faces; therefore, Q_i is unknotted. □

All that remains

If we could prove

Lemma

*If $p - q \geq 3$ and (B^p, B^q) is any ball pair,
then $B^p \searrow B^q$.*

we would finish the argument.

All that remains

If we could prove

Lemma

*If $p - q \geq 3$ and (B^p, B^q) is any ball pair,
then $B^p \searrow B^q$.*

we would finish the argument.

Hitherto, no use of the codimension assumption.

All that remains

If we could prove

Lemma

*If $p - q \geq 3$ and (B^p, B^q) is any ball pair,
then $B^p \searrow B^q$.*

we would finish the argument.

Hitherto, no use of the codimension assumption.

Proof Technique.

Sunny collapse.



