

Lecture 36: Harmonic functions

Math 660—Jim Fowler

Tuesday, August 10, 2010

Example

$$\int_0^\pi \log \sin \theta \, d\theta = -\pi \log 2$$

Single-valued branch

Define a single-valued branch of $\sqrt{1-x^2}$ for any region Ω , so that ± 1 are in the same component of $\mathbb{C} - \Omega$.

Harmonic functions

Harmonic functions

A function $u : \mathbb{R}^2 \rightarrow \mathbb{R}$ is **harmonic** if u is continuous, with continuous second partial derivatives, and

$$\Delta u = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0.$$

Example of harmonic functions

Example of harmonic functions

$$u(x, y) = ax + by.$$

Example of harmonic functions

$$u(x, y) = ax + by.$$

The real part of an analytic function.

Harmonic functions in polar coordinates

Harmonic functions in polar coordinates

$$r \frac{\partial}{\partial r} \left(r \frac{\partial u}{\partial r} \right) + \frac{\partial^2 u}{\partial \theta^2} = 0$$

Harmonic functions in polar coordinates

$$r \frac{\partial}{\partial r} \left(r \frac{\partial u}{\partial r} \right) + \frac{\partial^2 u}{\partial \theta^2} = 0$$

So if $\frac{\partial u}{\partial \theta} = 0$, then $u = a \log r + b$.

Integrals of harmonic functions

If u is harmonic, with conjugate harmonic function v ,

Integrals of harmonic functions

If u is harmonic, with conjugate harmonic function v ,
then $f(x, y) = u(x, y) + i v(x, y)$ is analytic,

Integrals of harmonic functions

If u is harmonic, with conjugate harmonic function v , then $f(x, y) = u(x, y) + i v(x, y)$ is analytic, and we can find $\int f dz$.

Integrals of harmonic functions

If u is harmonic, with conjugate harmonic function v , then $f(x, y) = u(x, y) + i v(x, y)$ is analytic, and we can find $f dz$.

$$dv = -\frac{\partial u}{\partial y} dx + \frac{\partial u}{\partial x} dy.$$

Integrals of harmonic functions

If u is harmonic, with conjugate harmonic function v , then $f(x, y) = u(x, y) + i v(x, y)$ is analytic, and we can find $f dz$.

$$dv = -\frac{\partial u}{\partial y} dx + \frac{\partial u}{\partial x} dy.$$

But in practice, there may be no conjugate harmonic function, so we take as a definition

$$*du = -\frac{\partial u}{\partial y} dx + \frac{\partial u}{\partial x} dy.$$

Hodge theory

For any cycle γ ,

$$\int_{\gamma} du = 0$$

Hodge theory

For any cycle γ ,

$$\int_{\gamma} du = 0$$

And if γ is homologous to zero, then

$$\int_{\gamma} f dz = 0$$

Hodge theory

For any cycle γ ,

$$\int_{\gamma} du = 0$$

And if γ is homologous to zero, then

$$\int_{\gamma} f dz = 0$$

And so $\int_{\gamma} *du = 0$ for cycles homologous to zero.

Simply connected regions

In a simply connected regions, $\int_{\gamma} *du = 0$ for all cycles, so u has a single-valued conjugate function v .

For every cycle γ homologous to zero,

$$\int_{\gamma} u_1 * du_2 - u_2 * du_1 = 0$$

For every cycle γ homologous to zero,

$$\int_{\gamma} u_1 * du_2 - u_2 * du_1 = 0$$

Proof: If conjugate harmonic functions exist (e.g., in a simply connected region like a rectangle), then

$$\begin{aligned} u_1 * du_2 - u_2 * du_1 &= u_1 dv_2 - u_2 dv_1 \\ &= u_1 dv_2 + v_1 du_2 - d(u_2 v_1) \\ &= (\operatorname{im}(u_1 + iv_1)(du_2 + idv_2)) - d(u_2 v_1), \end{aligned}$$

For every cycle γ homologous to zero,

$$\int_{\gamma} u_1 * du_2 - u_2 * du_1 = 0$$

Proof: If conjugate harmonic functions exist (e.g., in a simply connected region like a rectangle), then

$$\begin{aligned} u_1 * du_2 - u_2 * du_1 &= u_1 dv_2 - u_2 dv_1 \\ &= u_1 dv_2 + v_1 du_2 - d(u_2 v_1) \\ &= (\operatorname{im}(u_1 + iv_1)(du_2 + idv_2)) - d(u_2 v_1), \end{aligned}$$

so the integral vanishes.

