

# Lecture 25: Partial fractions

Math 153 Section 57

Monday November 24, 2008

## Overview

Recall that the limit comparison test allows us to determine whether

$$\sum \frac{\text{polynomial}}{\text{polynomial}}$$

converges. These quotients of polynomials are called rational functions.

Rational functions are maybe the “next simplest” class of functions after polynomials (which we understand very well).

Derivative of a rational function is a rational function, so we can take all the derivatives we would like—no problem!

But to antidifferentiate a rational function? For this, we use **partial fractions**. This is, in fact, how the computers do these integrals.

## Theory

We can integrate polynomials.

What are some rational functions we can integrate?

$$\int \frac{1}{ax+b} dx = \frac{\log(ax+b)}{a} + C$$

or somewhat more generally, for  $n > 1$ ,

$$\int \frac{1}{(ax+b)^n} dx = \frac{1}{a} \cdot \frac{(ax+b)^{-n+1}}{-n+1} + C$$

For quadratics:

$$\int \frac{x}{x^2+b^2} dx = \frac{1}{2} \log(x^2+b^2) + C$$

or even

$$\int \frac{x-a}{(x-a)^2+b^2} dx = \frac{1}{2} \log((x-a)^2+b^2) + C$$

We can also do

$$\int \frac{1}{(x-a)^2 + b^2} dx = \frac{1}{b} \arctan\left(\frac{x-a}{b}\right) + C$$

We can also do these if the denominators are raised to a power (with more difficulty).

**These functions suffice to do all rational functions.** But we need a machine for transforming the general rational function into something of this form. What would that be?

## Enter partial fractions

[ “I would like to see a proof of such and such.” “How many alphabets are you familiar with?” ]

The theorem. Let  $f(x) = p(x)/q(x)$ , a ratio of polynomials. We can assume that  $q$ 's leading term has a 1 coefficient.

The fundamental theorem of algebra: factor denominator as

$$q(x) = \prod_{i=1}^m (x - a_i)^{j_i} \cdot \prod_{i=1}^n (x^2 + b_i x + c_i)^{k_i}$$

that is, into linear and quadratic terms.

Then we can write  $f(x)$  as

$$f(x) = \frac{p(x)}{q(x)} = P(x) + \sum_{i=1}^m \sum_{r=1}^{j_i} \frac{A_{ir}}{(x - a_i)^r} + \sum_{i=1}^n \sum_{r=1}^{k_i} \frac{B_{ir}x + C_{ir}}{(x^2 + b_i x + c_i)^r}$$

This seems pretty awful. How can we ever hope to find all these constants?

## An example

Suppose

$$\frac{1}{(x-3)^2(x-2)} = \frac{A}{x-3} + \frac{B}{(x-3)^2} + \frac{C}{x-2}$$

The least clever (and most involved) way is to multiply through by the denominator:

$$1 = A(x-2)(x-3) + B(x-2) + C(x-3)^2$$

This gives

$$1 = (A+C)x^2 + (-5A+B-6C)x + (6A-2B+9C)$$

So  $A+C=0$ , and  $-5A+B-6C=0$ , and  $6A-2B+9C=1$ . Solve the three equations in three unknowns (the resulting equations will always have a unique solution). Get  $A=-C$ . So  $5C+B-6C=0$ , so  $B-C=0$ , so  $B=C$ . Then,  $-6C-2C+9C=1$ , so  $C=1$ , and therefore  $A=-1$  and  $B=1$ .

Check!

$$\frac{1}{(x-3)^2(x-2)} = \frac{-1}{x-3} + \frac{1}{(x-3)^2} + \frac{1}{x-2}$$

This is true.

This lets us do the integral

$$\int \frac{1}{(x-3)^2(x-2)} dx$$

because you can do

$$\int \left( \frac{-1}{x-3} + \frac{1}{(x-3)^2} + \frac{1}{x-2} \right) dx$$

## So many problems

The problems involved here are many.

You are not likely to see  $(x-3)^2(x-2)$ . You are more likely to see

$$x^3 - 8x^2 + 21x - 18$$

How are you supposed to **solve the factorization problem?**

## A tool in our toolbox, not a rule for solving all problems!

It is silly for me to give you lists of rules.

## A second example

$$f(x) = \frac{x^3 + 16}{x^3 - 4x^2 + 8x}$$

Do long division

$$f(x) = 1 + \frac{4x^2 - 8x + 16}{x^3 - 4x^2 + 8x} = 1 + \frac{4x^2 - 8x + 16}{x(x^2 - 4x + 8)}$$

$$\frac{4x^2 - 8x + 16}{x(x^2 - 4x + 8)} = \frac{A}{x} + \frac{Bx + C}{x^2 - 4x + 8}$$

Multiply through by  $x(x^2 - 4x + 8)$  to get

$$4x^2 - 8x + 16 = A(x^2 - 4x + 8) + (Bx + C)x$$

So  $A = 2$ . So  $B = 2$ . And  $C = 0$ . Therefore,

$$f(x) = 1 + 2 \left( \frac{1}{x} + \frac{x}{x^2 - 4x + 8} \right)$$

## Some tricks

Multiply the partial fraction decomposition by  $(x - a_i)$  and take the limit  $x \rightarrow a_i$ . Then only the term  $A_{i,1}$  survives, so

$$A_{i,1} = \lim_{x \rightarrow a_i} (x - a_i) f(x)$$

In the case of

$$f(x) = \frac{1}{(x-3)^2(x-2)}$$

this tells us that

$$\lim_{x \rightarrow 2} (x-2) \cdot \frac{1}{(x-3)^2(x-2)} = 1$$

but of course is useless for to find the coefficients on  $1/(x-3)$  and  $1/(x-3)^2$ .

We can find

$$\lim_{x \rightarrow 3} (x-3)^2 \cdot \frac{1}{(x-3)^2(x-2)} = 1$$

to discover that the coefficient on  $1/(x-3)^2$  is 1.

Then we know that

$$\frac{1}{(x-3)^2(x-2)} = \frac{1}{x-2} + \frac{1}{(x-3)^2} + \frac{A}{x-3}$$

We could plug in a value like  $x = 4$  to find:

$$\frac{1}{(4-3)^2(4-2)} = \frac{1}{4-2} + \frac{1}{(4-3)^2} + \frac{A}{4-3}$$

which becomes

$$\frac{1}{2} = -\frac{1}{2} + \frac{1}{1^2} + \frac{A}{1}$$

so  $A = -1$ .

## Underview

Some some tricks include: taking limits, and plugging in values.