# Lecture 44: Partial fractions and Infinite Products

Math 660—Jim Fowler

Friday, August 20, 2010

#### Final exam

 $11{:}30\mathrm{A.M.}{-}1{:}18\mathrm{P.M.}$  on Tuesday, August 24, 2010.

#### **Review session**

During lecture on Monday.

#### **Partial fractions**

$$\frac{\pi^2}{\sin^2(\pi z)} = \sum_{n \in \mathbb{Z}} \frac{1}{(z-n)^2}$$

#### **Infinite products**

Usually, define  $\prod_{n=1}^{\infty} p_n$  by taking the limit of the "partial products," provided the limit exists and is nonzero.

#### Infinite products

Usually, define  $\prod_{n=1}^{\infty} p_n$  by taking the limit of the "partial products," provided the limit exists and is nonzero.

By convention, the infinite product  $\prod_{n=1}^{\infty} p_n$  converges if only finitely many factors are zero, and if the product converges (in the above sense) after removing those zeroes.

#### Some basic theorems

 $\sum a_n$  converges implies  $\lim a_n = 0$ .

#### Some basic theorems

$$\sum a_n$$
 converges implies  $\lim a_n = 0$ .

$$\prod p_n$$
 converges implies  $\lim p_n = 1$ .

#### Some basic theorems

$$\sum a_n$$
 converges implies  $\lim a_n = 0$ .

$$\prod p_n$$
 converges implies  $\lim p_n = 1$ .

So we may write  $p_n = 1 + a_n$  with  $\lim a_n = 0$ .

To compute  $\prod (1 + a_n)$ , perhaps consider  $\sum \log(1 + a_n)$ .

To compute  $\prod (1 + a_n)$ , perhaps consider  $\sum \log(1 + a_n)$ .

Choose principal branch of log.

To compute  $\prod (1 + a_n)$ , perhaps consider  $\sum \log(1 + a_n)$ .

Choose principal branch of log.

Convergence of  $\sum \log(1+a_n)$  implies convergence of  $\prod (1+a_n)$ .

To compute  $\prod (1 + a_n)$ , perhaps consider  $\sum \log(1 + a_n)$ .

Choose principal branch of log.

Convergence of  $\sum \log(1+a_n)$  implies convergence of  $\prod (1+a_n)$ . Why?

### Logarithms can get confused

Convergence of  $\prod (1 + a_n)$  implies convergence of  $\sum \log(1 + a_n)$ ,

## Logarithms can get confused

Convergence of  $\prod (1 + a_n)$  implies convergence of  $\sum \log(1 + a_n)$ , although  $\sum \log(1 + a_n)$  may not converge to the principal branch log of  $\prod (1 + a_n)$ , it converges to some logarithm of  $\prod (1 + a_n)$ .

### Infinite products analyzed as series

In fact,  $\sum \log(1+a_n)$  converges absolutely iff  $\sum |a_n|$  converges.

#### **Theorem**

A nowhere vanishing entire function f(z) can be written as  $f(z) = e^{g(z)}$  for some analytic function g.

#### **Theorem**

A nowhere vanishing entire function f(z) can be written as  $f(z) = e^{g(z)}$  for some analytic function g.

**Proof:** 

#### **Theorem**

A nowhere vanishing entire function f(z) can be written as  $f(z) = e^{g(z)}$  for some analytic function g.

**Proof:** f'(z)/f(z) is analytic in the whole plane, so it is the derivative of some g(z).

#### **Theorem**

A nowhere vanishing entire function f(z) can be written as  $f(z) = e^{g(z)}$  for some analytic function g.

**Proof:** f'(z)/f(z) is analytic in the whole plane, so it is the derivative of some g(z). But  $f(z)e^{-g(z)}$  has zero derivative,

#### **Theorem**

A nowhere vanishing entire function f(z) can be written as  $f(z) = e^{g(z)}$  for some analytic function g.

**Proof:** f'(z)/f(z) is analytic in the whole plane, so it is the derivative of some g(z). But  $f(z)e^{-g(z)}$  has zero derivative, so f(z) is a multiple of  $e^{g(z)}$ .

#### **Theorem**

A nowhere vanishing entire function f(z) can be written as  $f(z) = e^{g(z)}$  for some analytic function g.

**Proof:** f'(z)/f(z) is analytic in the whole plane, so it is the derivative of some g(z). But  $f(z)e^{-g(z)}$  has zero derivative, so f(z) is a multiple of  $e^{g(z)}$ .

#### **Function with finitely many zeroes**

If f(z) has finitely many zeroes  $a_1, \ldots, a_N$  and a zero of order m at the origin, then we can write

$$f(z) = z^m e^{g(z)} \prod_{n=1}^{N} \left(1 - \frac{z}{a_n}\right).$$

### Function with finitely many zeroes

If f(z) has finitely many zeroes  $a_1, \ldots, a_N$  and a zero of order m at the origin, then we can write

$$f(z) = z^m e^{g(z)} \prod_{n=1}^N \left(1 - \frac{z}{a_n}\right).$$

What if f(z) has infinitely many zeroes?

### Function with finitely many zeroes

If f(z) has finitely many zeroes  $a_1, \ldots, a_N$  and a zero of order m at the origin, then we can write

$$f(z) = z^m e^{g(z)} \prod_{n=1}^N \left(1 - \frac{z}{a_n}\right).$$

What if f(z) has infinitely many zeroes?

If  $\sum_{n=1}^{\infty} 1/|a_n|$  converges, then

$$\prod_{n=1}^{\infty} \left( 1 - \frac{z}{a_n} \right)$$

converges absolutely, uniformly on disks  $|z| \leq R$ .

If  $\sum_{n=1}^{\infty} 1/|a_n|$  converges, then

$$\prod_{n=1}^{\infty} \left( 1 - \frac{z}{a_n} \right)$$

converges absolutely, uniformly on disks |z| < R.

In general, need to include extra "convergence producing" factors.

#### Weierstrass showed that

$$f(z) = z^m e^{\phi(z)} \prod_{n=1}^{\infty} \left(1 - \frac{z}{u_n}\right) e^{\frac{z}{u_n} + \frac{1}{2} \left(\frac{z}{u_n}\right)^2 + \dots + \frac{1}{\lambda_n} \left(\frac{z}{u_n}\right)^{\lambda_n}}$$

Weierstrass showed that

$$f(z) = z^m e^{\phi(z)} \prod_{n=1}^{\infty} \left(1 - \frac{z}{u_n}\right) e^{\frac{z}{u_n} + \frac{1}{2}\left(\frac{z}{u_n}\right)^2 + \dots + \frac{1}{\lambda_n}\left(\frac{z}{u_n}\right)^{\lambda_n}}$$

#### Corollary

Every function meromorphic in the whole plane is a quotient of entire functions.

 $\sin \pi z = \pi z \prod_{n=1}^{\infty} \left( 1 - \frac{z^2}{n^2} \right)$ 

$$\frac{\sin(x)}{x} = \left(1 - \frac{x^2}{\pi^2}\right) \left(1 - \frac{x^2}{4\pi^2}\right) \left(1 - \frac{x^2}{9\pi^2}\right) \dots = \prod_{n=1}^{\infty} \left(1 - \frac{x^2}{n^2\pi^2}\right)$$

with 
$$x = \pi/2$$
, give

 $\frac{2}{\pi} = \left(1 - \frac{1}{2^2}\right) \left(1 - \frac{1}{2^2 \cdot 4}\right) \left(1 - \frac{1}{2^2 \cdot 9}\right) \cdots = \prod_{n=1}^{\infty} \left(1 - \frac{1}{4n^2}\right),$ 

and taking the reciprocal,

 $\frac{\pi}{2} = \prod_{n=1}^{\infty} \left( \frac{4n^2}{4n^2 - 1} \right)$ 

 $=\prod_{n=0}^{\infty}\frac{(2n)(2n)}{(2n-1)(2n+1)}=\frac{2}{1}\cdot\frac{2}{3}\cdot\frac{4}{3}\cdot\frac{4}{5}\cdot\frac{6}{5}\cdot\frac{6}{7}\cdot\frac{8}{7}\cdot\frac{8}{9}\cdots$ 

with 
$$x = \pi/2$$
, gives

with  $x = \pi/2$ , gives