

Lecture 35: Some integrals

Math 660—Jim Fowler

Monday, August 9, 2010

Generalizing argument principle

If $g(z)$ is analytic in Ω , then $g(z)\frac{f'(z)}{f(z)}$ has

Generalizing argument principle

If $g(z)$ is analytic in Ω , then $g(z) \frac{f'(z)}{f(z)}$ has
residue $h g(a)$ at a zero a of order h

Generalizing argument principle

If $g(z)$ is analytic in Ω , then $g(z)\frac{f'(z)}{f(z)}$ has
residue $h g(a)$ at a zero a of order h and
residue $-h g(a)$ at a pole a of order h .

Generalizing argument principle

If $g(z)$ is analytic in Ω , then $g(z)\frac{f'(z)}{f(z)}$ has
residue $h g(a)$ at a zero a of order h and
residue $-h g(a)$ at a pole a of order h .

Therefore,

$$\frac{1}{2\pi i} \int_{\gamma} g(z) \frac{f'(z)}{f(z)} dz = \sum_j n(\gamma, a_j) g(a_j) - \sum_j n(\gamma, b_k) g(b_k)$$

$f(z) = w$ has n solutions z_1, \dots, z_n in disk
 $|z - z_0| < \epsilon$.

$f(z) = w$ has n solutions z_1, \dots, z_n in disk
 $|z - z_0| < \epsilon$.

$$\sum_j z_j(w) = \frac{1}{2\pi i} \int_{|z-z_0|=\epsilon} \frac{f'(z)}{f(z) - w} z \, dz$$

$f(z) = w$ has n solutions z_1, \dots, z_n in disk
 $|z - z_0| < \epsilon$.

$$\sum_j z_j(w) = \frac{1}{2\pi i} \int_{|z-z_0|=\epsilon} \frac{f'(z)}{f(z) - w} z \, dz$$

$$\sum_j z_j(w)^m = \frac{1}{2\pi i} \int_{|z-z_0|=\epsilon} \frac{f'(z)}{f(z) - w} z^m \, dz$$

$f(z) = w$ has n solutions z_1, \dots, z_n in disk
 $|z - z_0| < \epsilon$.

$$\sum_j z_j(w) = \frac{1}{2\pi i} \int_{|z-z_0|=\epsilon} \frac{f'(z)}{f(z) - w} z \, dz$$

$$\sum_j z_j(w)^m = \frac{1}{2\pi i} \int_{|z-z_0|=\epsilon} \frac{f'(z)}{f(z) - w} z^m \, dz$$

so $z_j(w)$ are roots of a polynomial with coefficients depending analytically on w .

Integration tricks!

- ▶ The integrals we care about are usually over intervals;

- ▶ The integrals we care about are usually over intervals;
our residue technique requires a closed curve;

- ▶ The integrals we care about are usually over intervals;
our residue technique requires a closed curve;
need tricks for converting integrals over intervals to integrals over curves.

- ▶ The integrals we care about are usually over intervals;
our residue technique requires a closed curve;
need tricks for converting integrals over intervals to integrals over curves.
- ▶ The integrals we care about might be for real-valued functions;

- ▶ The integrals we care about are usually over intervals;
our residue technique requires a closed curve;
need tricks for converting integrals over intervals to integrals over curves.
- ▶ The integrals we care about might be for real-valued functions;
thankfully, many real integrals are actually of analytic functions.

Rational functions of trig functions

$$\int_0^{2\pi} R(\cos \theta, \sin \theta) d\theta$$

Rational functions of trig functions

$$\int_0^{2\pi} R(\cos \theta, \sin \theta) d\theta$$

Make substitution $z = e^{i\theta}$, and get

Rational functions of trig functions

$$\int_0^{2\pi} R(\cos \theta, \sin \theta) d\theta$$

Make substitution $z = e^{i\theta}$, and get

$$-i \int_{\text{unit circle}} R\left(\frac{1}{2}\left(z + \frac{1}{z}\right), \frac{1}{2i}\left(z - \frac{1}{z}\right)\right) \frac{dz}{z}$$

Example 1

$$\int_0^\pi \frac{d\theta}{a + \cos \theta} \text{ for } a > 1.$$

Example 2

For n even, compute

$$\int_0^{2\pi} \cos^n \theta \, d\theta$$

Example 2

For n even, compute

$$\int_0^{2\pi} \cos^n \theta \, d\theta = -i \int_{\text{circle}} \frac{1}{2^n} \left(z + \frac{1}{z} \right)^n \frac{dz}{z}$$

Example 2

For n even, compute

$$\begin{aligned}\int_0^{2\pi} \cos^n \theta \, d\theta &= -i \int_{\text{circle}} \frac{1}{2^n} \left(z + \frac{1}{z} \right)^n \frac{dz}{z} \\ &= -i \int_{\text{circle}} \frac{1}{2^n} \left(\sum_m \binom{n}{m} z^{2m-n-1} \right) dz\end{aligned}$$

Example 2

For n even, compute

$$\begin{aligned}\int_0^{2\pi} \cos^n \theta \, d\theta &= -i \int_{\text{circle}} \frac{1}{2^n} \left(z + \frac{1}{z} \right)^n \frac{dz}{z} \\ &= -i \int_{\text{circle}} \frac{1}{2^n} \left(\sum_m \binom{n}{m} z^{2m-n-1} \right) dz \\ &= -i \int_{\text{circle}} \frac{1}{2^n} \binom{n}{n/2} z^{-1} dz\end{aligned}$$

Example 2

For n even, compute

$$\begin{aligned}\int_0^{2\pi} \cos^n \theta \, d\theta &= -i \int_{\text{circle}} \frac{1}{2^n} \left(z + \frac{1}{z} \right)^n \frac{dz}{z} \\ &= -i \int_{\text{circle}} \frac{1}{2^n} \left(\sum_m \binom{n}{m} z^{2m-n-1} \right) dz \\ &= -i \int_{\text{circle}} \frac{1}{2^n} \binom{n}{n/2} z^{-1} dz \\ &= \pi \frac{1}{2^{n-1}} \binom{n}{n/2}\end{aligned}$$

Rational functions

By integrating over large semicircles, compute

$$\frac{1}{2\pi i} \int_{-\infty}^{\infty} R(x) dx = \sum_{\substack{z=a \\ \operatorname{Im} z > 0}} \operatorname{Res} R(z)$$

Rational functions times e^{iz}

By integrating over large semicircles, compute

$$\frac{1}{2\pi i} \int_{-\infty}^{\infty} R(x) e^{iz} dx = \sum_{\operatorname{Im} z > 0} \operatorname{Res}_{z=a} R(z) e^{iz}$$

provided $R(z)$ has a zero of order two at infinity.

Rational functions times e^{iz}

Integrate over rectangle with vertices $x_2, x_2 + iy, -x_1 + iy, -x_1$.

$$\frac{1}{2\pi i} \int_{-\infty}^{\infty} R(x) e^{iz} dx = \sum_{\operatorname{Im} z > 0} \operatorname{Res}_{z=a} R(z) e^{iz}$$

which holds provided $R(z)$ has a zero at infinity.

Powers of z

$$\int_0^\infty x^\alpha R(x) dx = 2 \int_0^\infty t^{2\alpha+1} R(t^2) dt$$

and then integrate over semicircle minus semicircle.

Example

$$\int_0^{\pi} \log \sin \theta \, d\theta$$