Lecture 40: Weierstrass' theorem

Math 660—Jim Fowler

Monday, August 16, 2010


Pictures of harmonic functions


The Poisson integral produces a harmonic function, given values on the boundary.


Pictures of harmonic functions


The Poisson integral produces a harmonic function, given values on the boundary.


Let's see some examples!


























Holomorphic means complex differentiable.

Holomorphic means complex differentiable.

Analytic means has a power series.

Holomorphic means complex differentiable.

Analytic means has a power series.

Now we show that holomorphic implies analytic.

Convergence of analytic functions

Suppose $f_n(z)$ is analytic in Ω_n and $f_n(z) \to f(z)$ in a region Ω , uniformly on compact subsets of Ω . Then f(z) is analytic in Ω , and $f'_n(z) \to f'(z)$ uniformly on compact subsets.

Convergence of analytic functions

Suppose $f_n(z)$ is analytic in Ω_n and $f_n(z) \to f(z)$ in a region Ω , uniformly on compact subsets of Ω . Then f(z) is analytic in Ω , and $f'_n(z) \to f'(z)$ uniformly on compact subsets.

Proof:

Convergence of analytic functions

Suppose $f_n(z)$ is analytic in Ω_n and $f_n(z) \to f(z)$ in a region Ω , uniformly on compact subsets of Ω . Then f(z) is analytic in Ω , and $f'_n(z) \to f'(z)$ uniformly on compact subsets.

Proof: Morera's theorem.

Zeroes and convergence

If $f_n(z) \neq 0$ in a region Ω , and $f_n(z) \rightarrow f(z)$ uniformly on compact subsets of Ω then f(z) is either everywhere or nowhere zero.



Taylor series