

Lecture 34: Argument principle

Math 660—Jim Fowler

Friday, August 6, 2010

Example 1

$$f(z) = \frac{1}{z^4 + 4}$$

Compute $\int_{\gamma} f \, dz$ for γ a circle of radius two and center 1.

Counting zeroes

Counting zeroes

Let $D^2 = \{z \in \mathbb{C} : |z| < 1\}$ and $f : D^2 \rightarrow \mathbb{C}$ analytic.
Assume $f(z) \not\equiv 0$.

Counting zeroes

Let $D^2 = \{z \in \mathbb{C} : |z| < 1\}$ and $f : D^2 \rightarrow \mathbb{C}$ analytic.
Assume $f(z) \not\equiv 0$.

Assume $f(z)$ has only finitely many zeroes, z_1, \dots, z_n ,
with multiplicity.

Counting zeroes

Let $D^2 = \{z \in \mathbb{C} : |z| < 1\}$ and $f : D^2 \rightarrow \mathbb{C}$ analytic.
Assume $f(z) \not\equiv 0$.

Assume $f(z)$ has only finitely many zeroes, z_1, \dots, z_n ,
with multiplicity.

$f(z) = (z - z_1) \cdots (z - z_n)g(z)$ for $g(z)$ analytic and
nowhere zero.

Counting zeroes

Let $D^2 = \{z \in \mathbb{C} : |z| < 1\}$ and $f : D^2 \rightarrow \mathbb{C}$ analytic.
Assume $f(z) \not\equiv 0$.

Assume $f(z)$ has only finitely many zeroes, z_1, \dots, z_n ,
with multiplicity.

$f(z) = (z - z_1) \cdots (z - z_n)g(z)$ for $g(z)$ analytic and
nowhere zero.

$$\frac{f'(z)}{f(z)} = \frac{1}{z - z_1} + \cdots + \frac{1}{z - z_n} + \frac{g'(z)}{g(z)}$$

Now $\int_{\gamma} \frac{g'(z)}{g(z)} dz = 0$, so

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f'(z)}{f(z)} dz = n(\gamma, z_1) + \cdots + n(\gamma, z_n)$$

Now $\int_{\gamma} \frac{g'(z)}{g(z)} dz = 0$, so

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f'(z)}{f(z)} dz = n(\gamma, z_1) + \cdots + n(\gamma, z_n)$$

In fact, this is true even if there are infinitely many zeroes in D^2 .

Counting zeroes, again

Theorem ()

If $f(z)$ is meromorphic in Ω , with zeroes a_j and poles b_k , then

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f'(z) dz}{f(z)} = \sum_j n(\gamma, a_j) - \sum_k n(\gamma, b_k)$$

for every γ homologous to zero in Ω .

Counting zeroes, again

Theorem (Argument principle)

If $f(z)$ is meromorphic in Ω , with zeroes a_j and poles b_k , then

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f'(z) dz}{f(z)} = \sum_j n(\gamma, a_j) - \sum_k n(\gamma, b_k)$$

for every γ homologous to zero in Ω .

Corollary (Rouché's theorem)

*Suppose $\gamma \in \Omega$ with $[\gamma] = [0] \in H_1(\Omega)$,
and $n(\gamma, a) = 0$ or 1 for $a \notin \omega$.*

*Suppose $f(z)$ and $g(z)$ are analytic in Ω ,
and satisfy $|f(z) - g(z)| < |f(z)|$ on γ .*

*Then $f(z)$ and $g(z)$ have the same number of zeroes
inside γ .*

Finding zeroes via Taylor's theorem

Suppose we want to know how many zeroes $f(z)$ has inside $B_R(0)$.

Finding zeroes via Taylor's theorem

Suppose we want to know how many zeroes $f(z)$ has inside $B_R(0)$.

Use Taylor's theorem to write $f(z) = p(z) + z^n f_n(z)$ for a polynomial $p(z)$ with degree $n - 1$.

Finding zeroes via Taylor's theorem

Suppose we want to know how many zeroes $f(z)$ has inside $B_R(0)$.

Use Taylor's theorem to write $f(z) = p(z) + z^n f_n(z)$ for a polynomial $p(z)$ with degree $n - 1$.

If we can prove $R^n |f_n(z)| < |p(z)|$, then we need only determine solutions of $p(z) = 0$.

Example

How many roots does $z^7 - 2z^5 + 6z^3 - z + 1 = 0$ have inside the disk $|z| < 1$?

Generalizing argument principle

If $g(z)$ is analytic in Ω , then $g(z)\frac{f'(z)}{f(z)}$ has

Generalizing argument principle

If $g(z)$ is analytic in Ω , then $g(z) \frac{f'(z)}{f(z)}$ has
residue $h g(a)$ at a zero a of order h

Generalizing argument principle

If $g(z)$ is analytic in Ω , then $g(z)\frac{f'(z)}{f(z)}$ has
residue $h g(a)$ at a zero a of order h and
residue $-h g(a)$ at a pole a of order h .

Generalizing argument principle

If $g(z)$ is analytic in Ω , then $g(z)\frac{f'(z)}{f(z)}$ has
residue $h g(a)$ at a zero a of order h and
residue $-h g(a)$ at a pole a of order h .

Therefore,

$$\frac{1}{2\pi i} \int_{\gamma} g(z) \frac{f'(z)}{f(z)} dz = \sum_j n(\gamma, a_j) g(a_j) - \sum_j n(\gamma, b_k) g(b_k)$$

$f(z) = w$ has n solutions z_1, \dots, z_n in disk
 $|z - z_0| < \epsilon$.

$f(z) = w$ has n solutions z_1, \dots, z_n in disk
 $|z - z_0| < \epsilon$.

$$\sum_j z_j(w) = \frac{1}{2\pi i} \int_{|z-z_0|=\epsilon} \frac{f'(z)}{f(z) - w} z \, dz$$

$f(z) = w$ has n solutions z_1, \dots, z_n in disk
 $|z - z_0| < \epsilon$.

$$\sum_j z_j(w) = \frac{1}{2\pi i} \int_{|z-z_0|=\epsilon} \frac{f'(z)}{f(z) - w} z \, dz$$

$$\sum_j z_j(w)^m = \frac{1}{2\pi i} \int_{|z-z_0|=\epsilon} \frac{f'(z)}{f(z) - w} z^m \, dz$$

$f(z) = w$ has n solutions z_1, \dots, z_n in disk
 $|z - z_0| < \epsilon$.

$$\sum_j z_j(w) = \frac{1}{2\pi i} \int_{|z-z_0|=\epsilon} \frac{f'(z)}{f(z) - w} z \, dz$$

$$\sum_j z_j(w)^m = \frac{1}{2\pi i} \int_{|z-z_0|=\epsilon} \frac{f'(z)}{f(z) - w} z^m \, dz$$

so $z_j(w)$ are roots of a polynomial with coefficients depending analytically on w .