

Lecture 25: Zeroes and poles

Math 660—Jim Fowler

Tuesday, July 26, 2011

Removable singularities

$f : \Omega - \{a\} \rightarrow \mathbb{C}$ analytic
can be extended to $\Omega \rightarrow \mathbb{C}$
iff $\lim_{z \rightarrow a} f(z)(z - a) = 0$.

Removable singularities

$f : \Omega - \{a\} \rightarrow \mathbb{C}$ analytic
can be extended to $\Omega \rightarrow \mathbb{C}$
iff $\lim_{z \rightarrow a} f(z)(z - a) = 0$.

Why? Cauchy's formula is valid.

Apply this trick to

$$F(z) = \frac{f(z) - f(a)}{z - a}$$

and get $f(z) = f(a) + (z - a)F(z)$.

Apply this trick to

$$F(z) = \frac{f(z) - f(a)}{z - a}$$

and get $f(z) = f(a) + (z - a)F(z)$.
Rinse, repeat.

Consequently . . .

If $f : \Omega \rightarrow \mathbb{C}$ is analytic,
for any $a \in \Omega$, we can write

$$f(z) = f(a) + \sum_{n=1}^k \frac{f^{(n)}(a)}{n!}(z-a)^n + F(z)(z-a)^{k+1}$$

for some analytic $F : \Omega \rightarrow \mathbb{C}$.

Say more about F

If $f : \Omega \rightarrow \mathbb{C}$ is analytic,
for any $a \in \Omega$, we can write

$$f(z) = f(a) + \sum_{n=1}^k \frac{f^{(n)}(a)}{n!} (z - a)^n + F(z)(z - a)^{k+1}$$

for the analytic $F : \Omega \rightarrow \mathbb{C}$.

$$F(z) = \frac{1}{2\pi i \eta(z, \gamma)} \int_{\gamma} \frac{f(w)}{(w - a)^n (w - z)} dw.$$

Zeroes and poles

If all derivatives of f vanish at a ,
then $f(z) = F(z)(z - a)^n$

Zeroes and poles

If all derivatives of f vanish at a ,
then $f(z) = F(z)(z - a)^n$
Estimate the size of $F(z)$.

Zeroes and poles

If all derivatives of f vanish at a ,

$$\text{then } f(z) = F(z)(z - a)^n$$

Estimate the size of $F(z)$.

$f(z)$ vanishes on Ω .

Zero of order k

If $f^{(n)}(a) = 0$ for $n \in \{0, \dots, k - 1\}$,
but $f^{(k)}(a) \neq 0$,
we say f has a **zero of order k** .

Zero of order k

If $f^{(n)}(a) = 0$ for $n \in \{0, \dots, k - 1\}$,
but $f^{(k)}(a) \neq 0$,
we say f has a **zero of order k** .
It's like a polynomial!

Zeroes are isolated

Zeroes of analytic function are isolated.

Zeroes are isolated

Zeroes of analytic function are isolated.

So if $f(z) = g(z)$ for $z \in S$,
(what sort of set is S ?)

then $f \equiv g$ on Ω .

Classifying singularities

Δ a disk, with center a .

Suppose $f : \Delta - \{a\} \rightarrow \mathbb{C}$ is analytic.

Classifying singularities

Δ a disk, with center a .

Suppose $f : \Delta - \{a\} \rightarrow \mathbb{C}$ is analytic.

If $\lim_{z \rightarrow a} f(z) = \infty$,

we call a a **pole**.

Consider $1/f(z)$,

which has a zero at a of order k ,

now called the order of the pole.

Classifying singularities

Δ a disk, with center a .

Suppose $f : \Delta - \{a\} \rightarrow \mathbb{C}$ is analytic.

If $\lim_{z \rightarrow a} f(z) = \infty$,

we call a a **pole**.

Consider $1/f(z)$,

which has a zero at a of order k ,

now called the order of the pole.

Meromorphic means holomorphic except for poles.

Meromorphic functions

Meromorphic functions

Quotients of analytic functions

Meromorphic functions

Quotients of analytic functions

Poles of meromorphic functions are isolated

Meromorphic functions

Quotients of analytic functions

Poles of meromorphic functions are isolated

Think of the relationship between \mathbb{Z} and \mathbb{Q} .

$$Z = \{\alpha \in \mathbb{R} : \lim_{z \rightarrow a} |(z-a)^\alpha f(z)| = 0\}$$

$$P = \{\alpha \in \mathbb{R} : \lim_{z \rightarrow a} |(z-a)^\alpha f(z)| = \infty\}$$

$$Z = \{\alpha \in \mathbb{R} : \lim_{z \rightarrow a} |(z - a)^\alpha f(z)| = 0\}$$

$$P = \{\alpha \in \mathbb{R} : \lim_{z \rightarrow a} |(z - a)^\alpha f(z)| = \infty\}$$

If $\alpha \in Z$ and $\beta > \alpha$, then $\beta \in Z$.

If $\alpha \in P$ and $\beta < \alpha$, then $\beta \in P$.

$$Z = \{\alpha \in \mathbb{R} : \lim_{z \rightarrow a} |(z - a)^\alpha f(z)| = 0\}$$

$$P = \{\alpha \in \mathbb{R} : \lim_{z \rightarrow a} |(z - a)^\alpha f(z)| = \infty\}$$

If $\alpha \in Z$ and $\beta > \alpha$, then $\beta \in Z$.

If $\alpha \in P$ and $\beta < \alpha$, then $\beta \in P$.

- ▶ $Z = \mathbb{R}$ and $P = \emptyset$,
- ▶ $Z = (n, \infty)$ and $P = (-\infty, n)$,
- ▶ $Z = \emptyset$ and $P = \emptyset$.

Partial fractions redux

If $f(z)$ has a pole of order k , consider

$$(z-a)^k f(z) = a_0 + a_1(z-a) + \cdots + a_{k-1}(z-a)^{k-1} + F(z)(z-a)^k$$

and divide by $(z - a)^k$ to get the **singular part**.

$Z = \emptyset$ and $P = \emptyset$

These are **essential singularities**.

Theorem (Weierstrass)

$f : \Omega - \{a\} \rightarrow \mathbb{C}$ analytic
with essential singularity at a .

For every $w \in \mathbb{C}$ and $\epsilon > 0$,
there is $z \in B_\epsilon(a) \cap \Omega$
so that $f(z) = w$.

Singularities at infinity

To determine the behavior of $f(z)$ at infinity,
consider instead $f(1/z)$.

Singularities at infinity

To determine the behavior of $f(z)$ at infinity,
consider instead $f(1/z)$.

What sort of singularity does $\sin z$ have near infinity?



