


Topology of Piecewise-Linear Manifolds

Jim Fowler

Lecture 2
Summer 2010

Goal

Definitions?





A **simplicial complex** K is
a collection of finite sets (called the “simplexes”),
with the property that

if $\sigma \in K$, and $\tau \subset \sigma$, then $\tau \in K$.

A **simplicial complex** K is
a collection of finite sets (called the “simplexes”),
with the property that

$$\text{if } \sigma \in K, \text{ and } \tau \subset \sigma, \quad \text{then } \tau \in K.$$

This definition is pure combinatorics,
but we will think of this as a geometric object.

Examples

$$= \{\emptyset, \{0\}, \{1\}, \{0, 1\}\}$$

Examples

$$I = \{\emptyset, \{0\}, \{1\}, \{0, 1\}\}$$

Examples

$$\begin{aligned} I &= \{\emptyset, \{0\}, \{1\}, \{0, 1\}\} \\ &= \{\emptyset, \{0\}, \{1\}\} \end{aligned}$$

Examples

$$I = \{\emptyset, \{0\}, \{1\}, \{0, 1\}\}$$

$$S^0 = \{\emptyset, \{0\}, \{1\}\}$$

Examples

$$I = \{\emptyset, \{0\}, \{1\}, \{0, 1\}\}$$

$$S^0 = \{\emptyset, \{0\}, \{1\}\}$$

$$= \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}, \{a, c\}\}$$

Examples

$$I = \{\emptyset, \{0\}, \{1\}, \{0, 1\}\}$$

$$S^0 = \{\emptyset, \{0\}, \{1\}\}$$

$$S^1 = \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}, \{a, c\}\}$$

Examples

$$I = \{\emptyset, \{0\}, \{1\}, \{0, 1\}\}$$

$$S^0 = \{\emptyset, \{0\}, \{1\}\}$$

$$\begin{aligned} S^1 &= \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}, \{a, c\}\} \\ &= \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}\} \end{aligned}$$

Examples

$$I = \{\emptyset, \{0\}, \{1\}, \{0, 1\}\}$$

$$S^0 = \{\emptyset, \{0\}, \{1\}\}$$

$$S^1 = \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}, \{a, c\}\}$$

$$V = \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}\}$$

Examples

$$I = \{\emptyset, \{0\}, \{1\}, \{0, 1\}\}$$

$$S^0 = \{\emptyset, \{0\}, \{1\}\}$$

$$S^1 = \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}, \{a, c\}\}$$

$$V = \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}\}$$

Shouldn't $V \cong I$?

Non-examples

Non-examples

$$\{\emptyset, \{0\}, \{0, 1\}\}$$

Non-examples

$$\{\emptyset, \{0\}, \{0, 1\}\}$$

$$\{\{0\}, \{1\}, \{0, 1\}\}$$

Examples

The n -simplex Δ^n is a complex: label the $n + 1$ vertices of Δ^n using the set $V = \{0, 1, 2, \dots, n\}$, then the simplexes of Δ^n are all 2^{n+1} subsets of V .

The n -sphere S^n consists of all simplexes in Δ^n , except for the top dimensional simplex $\{0, 1, \dots, n\}$.

Examples

The n -simplex Δ^n is a complex: label the $n + 1$ vertices of Δ^n using the set $V = \{0, 1, 2, \dots, n\}$, then the simplexes of Δ^n are all 2^{n+1} subsets of V .

The n -sphere S^n consists of all simplexes in Δ^n , except for the top dimensional simplex $\{0, 1, \dots, n\}$.

Problem

Calculate $\chi(S^n)$.

Simplicial Maps

A **simplicial map** $f : K \rightarrow L$ is a function $f : \text{vert}(K) \rightarrow \text{vert}(L)$ so that, whenever $\sigma \in K$, then $f(\sigma) \in L$.

Simplicial Maps

A **simplicial map** $f : K \rightarrow L$ is a function $f : \text{vert}(K) \rightarrow \text{vert}(L)$ so that, whenever $\sigma \in K$, then $f(\sigma) \in L$.

This can crush simplexes

Simplicial Maps

A **simplicial map** $f : K \rightarrow L$ is a function $f : \text{vert}(K) \rightarrow \text{vert}(L)$ so that, whenever $\sigma \in K$, then $f(\sigma) \in L$.

This can crush simplexes (e.g., a trivial example?)

Simplicial Maps

A **simplicial map** $f : K \rightarrow L$ is a function $f : \text{vert}(K) \rightarrow \text{vert}(L)$ so that, whenever $\sigma \in K$, then $f(\sigma) \in L$.

This can crush simplexes (e.g., a trivial example?)

Problem

Find a simplicial map $f : T^2 \rightarrow S^2$ which doesn't crush any edges (i.e., edges are sent to edges, not to vertices).

Simplicial Maps

A **simplicial map** $f : K \rightarrow L$ is a function $f : \text{vert}(K) \rightarrow \text{vert}(L)$ so that, whenever $\sigma \in K$, then $f(\sigma) \in L$.

This can crush simplexes (e.g., a trivial example?)

Problem

Find a simplicial map $f : T^2 \rightarrow S^2$ which doesn't crush any edges (i.e., edges are sent to edges, not to vertices).

Think "4-color the vertices of T^2 ."

Conversely...

Problem

Find a simplicial map $f : S^2 \rightarrow T^2$ which doesn't crush any edges.

Conversely...

Problem

Find a simplicial map $f : S^2 \rightarrow T^2$ which doesn't crush any edges.

Think "Skewer and fold."

Problem

Do there exist simplicial maps $f : T^2 \rightarrow S^2$ and $g : S^2 \rightarrow T^2$ which are inverses of each other?

Problem

Do there exist simplicial maps $f : T^2 \rightarrow S^2$ and $g : S^2 \rightarrow T^2$ which are inverses of each other?

No. **And why not?**

Problem

Do there exist simplicial maps $f : T^2 \rightarrow S^2$ and $g : S^2 \rightarrow T^2$ which are inverses of each other?

No. **And why not?**

- ▶ Euler characteristic.

Problem

Do there exist simplicial maps $f : T^2 \rightarrow S^2$ and $g : S^2 \rightarrow T^2$ which are inverses of each other?

No. **And why not?**

- ▶ Euler characteristic.
- ▶ Separation via curves.

Problem

Suppose $f : S^1 \rightarrow S^2$ is an injective simplicial map (i.e., distinct vertices are sent to distinct vertices). Does the image of f necessarily separate S^2 into two pieces?

Problem

Suppose $f : S^1 \rightarrow S^2$ is an injective simplicial map (i.e., distinct vertices are sent to distinct vertices). Does the image of f necessarily separate S^2 into two pieces?

Yes.

Problem

Suppose $f : S^1 \rightarrow S^2$ is an injective simplicial map (i.e., distinct vertices are sent to distinct vertices). Does the image of f necessarily separate S^2 into two pieces?

Yes.

Proof

A problem with our definitions: we want to talk about S^2 , but there are so many simplicial complexes which deserve to be called S^2 .

Our notion of simplicial complex is too rigid to be the right notion topologically.

Example

Let K be a circle with three arcs
and L be a circle with four arcs
(i.e., the boundary of a square).

Then K and L are not simplicially isomorphic.

Fixing the theory

Need to define a few things first. . .

- ▶ star
- ▶ closure
- ▶ link
- ▶ subdivision

Star

Definition

Let K be a complex, and $\sigma \in K$ a simplex.

The **star** of σ in K ,

written $\text{st}(\sigma, K)$,

is defined by

$$\text{st}(\sigma, K) = \{\tau \in K : \sigma < \tau\},$$

i.e., the star of σ includes all the simplexes having σ as a face.

Star

Definition

Let K be a complex, and $\sigma \in K$ a simplex.

The **star** of σ in K ,

written $\text{st}(\sigma, K)$,

is defined by

$$\text{st}(\sigma, K) = \{\tau \in K : \sigma < \tau\},$$

i.e., the star of σ includes all the simplexes having σ as a face.

Problem

Is the star of a simplex a complex?

Closure

Definition

Let S be a collection of simplexes in K .

The **closure** of S ,

written as $\text{cl}(S)$,

is the smallest subcomplex of K

containing the simplexes in S .

Closure

Definition

Let S be a collection of simplexes in K .

The **closure** of S ,

written as $\text{cl}(S)$,

is the smallest subcomplex of K

containing the simplexes in S .

Problem

Relate $\text{cl}(\text{cl}(S))$ and $\text{cl}(S)$.

Link

Definition

Let K be a complex, and $\sigma \in K$ a simplex.

The **link** of $\sigma \in K$,

written $\text{lk}(\sigma, K)$,

consists of those simplexes in K which are in $\text{cl}(\text{st}(\sigma, K))$ but not touching σ ; in other words,

$$\text{lk}(\sigma, K) = \{\tau \in \text{cl}(\text{st}(\sigma, K)) : \tau \cap \sigma = \emptyset\}.$$

Link

Definition

Let K be a complex, and $\sigma \in K$ a simplex.

The **link** of $\sigma \in K$,

written $\text{lk}(\sigma, K)$,

consists of those simplexes in K which are in $\text{cl}(\text{st}(\sigma, K))$ but not touching σ ; in other words,

$$\text{lk}(\sigma, K) = \{\tau \in \text{cl}(\text{st}(\sigma, K)) : \tau \cap \sigma = \emptyset\}.$$

Problem

Is the link of a simplex a complex?

Definition

Let K be a complex, and $\sigma \in K$ a simplex.

The **stellar subdivision** of K at σ is a new complex K_σ with:

- ▶ the vertices of K along with a brand new vertex v .
- ▶ the simplexes of K not in $\text{st}(\sigma, K)$, along with the simplexes in $v * (\partial\sigma) * \text{lk}(\sigma, K)$.

We might say:

$$K_\sigma := (K - \text{st}(\sigma, K)) \cup (v * (\partial\sigma) * \text{lk}(\sigma, K))$$

Definition

Let K be a complex, and $\sigma \in K$ a simplex.

The **stellar subdivision** of K at σ is a new complex K_σ with:

- ▶ the vertices of K along with a brand new vertex v .
- ▶ the simplexes of K not in $\text{st}(\sigma, K)$, along with the simplexes in $v * (\partial\sigma) * \text{lk}(\sigma, K)$.

We might say:

$$K_\sigma := (K - \text{st}(\sigma, K)) \cup (v * (\partial\sigma) * \text{lk}(\sigma, K))$$

Problem

Stellar subdivision of a complex is a complex?

subdivision = repeated stellar subdivision

Definition

Let K, L be complexes.

If K can be produced through a (possibly empty) sequence of stellar subdivisions of L , we say that K is a **subdivision** of L , and write $K \triangleleft L$.

subdivision = repeated stellar subdivision

Definition

Let K, L be complexes.

If K can be produced through a (possibly empty) sequence of stellar subdivisions of L , we say that K is a **subdivision** of L , and write $K \triangleleft L$.

As we will see, the *real* definition of subdivision is more general than this.

Piecewise linear maps

Definition

Let K, K', L, L' be complexes, with $K' \triangleleft K$ and $L' \triangleleft L$. If $f : K' \rightarrow L'$ is a simplicial map, we call $f : K \rightarrow L$ a **piecewise linear map** (or a **PL map** for short). We call $f : K' \rightarrow L'$ an **underlying simplicial map**.

Piecewise linear maps

Definition

Let K, K', L, L' be complexes, with $K' \triangleleft K$ and $L' \triangleleft L$. If $f : K' \rightarrow L'$ is a simplicial map, we call $f : K \rightarrow L$ a **piecewise linear map** (or a **PL map** for short). We call $f : K' \rightarrow L'$ an **underlying simplicial map**.

We write $f : K \rightarrow L$ for a PL map, but such a map does not send simplexes in K to simplexes in L

Piecewise linear maps

Definition

Let K, K', L, L' be complexes, with $K' \triangleleft K$ and $L' \triangleleft L$. If $f : K' \rightarrow L'$ is a simplicial map, we call $f : K \rightarrow L$ a **piecewise linear map** (or a **PL map** for short). We call $f : K' \rightarrow L'$ an **underlying simplicial map**.

The *real* definition of PL map is more general than this.

Going back, rethinking everything...

Are S^2 and T^2 the same?

Going back, rethinking everything...

Are S^2 and T^2 the same?

We will check that χ is well-defined.