

# Lecture 6: Abel's limit theorem

Math 660—Jim Fowler

Monday, June 27, 2011

# Homework questions

How did it go?

## Abel's limit theorem

Suppose  $\sum a_n$  converges and let  $f(z) = \sum a_n z^n$ .  
Then  $\lim_{z \rightarrow 1} f(z) = f(1)$  provided  $z \rightarrow 1$  so that

$$\frac{|1 - z|}{1 - |z|}$$

remains bounded.

Wlog  $\sum a_n = 0$  by changing  $a_0$ .

$$\begin{aligned}s_n(z) &= a_0 + a_1z + \cdots + a_nz^n \\&= s_0 + (s_1 - s_0)z + \cdots + (s_n - s_{n-1})z^n \\&= s_0(1 - z) + s_1(z - z^2) + \cdots + s_{n-1}(z^{n-1} - z^n) + s_nz^n \\&= (1 - z)(s_0 + \cdots + s_{n-1}z^{n-1}) + s_nz^n\end{aligned}$$

Since  $s_nz^n \rightarrow 0$ , we conclude

$$f(z) = (1 - z) \sum s_nz^n$$

Then

$$|f(z)| \leq |1 - z| \left| \sum_{n=0}^{m-1} s_nz^n \right| + M\epsilon$$

# Applications

$$\sum_{k=0}^{\infty} (-1)^n / (n+1) = \log 2$$

$$\sum_{k=0}^{\infty} (-1)^n / (2n+1) = \pi/4$$

Armed with a good theorem,  
we can extend the objects under consideration.

Armed with a good theorem,  
we can extend the objects under consideration.  
Abel summability

Let  $\sum a_n$  be a series—**convergent or not!**

$f(z) = \sum a_n z^n$  is more likely to converge  
for  $|z| < 1$  than  $\sum a_n$ .

If  $\sum a_n z^n$  converges for  $|z| < 1$ ,  
and  $\lim_{x \rightarrow 1^-} f(x) = L$ ,

say that  $a_n$  is Abel summable, with Abel sum  $L$



Let  $\sum a_n$  be a series—**convergent or not!**

$f(z) = \sum a_n z^n$  is more likely to converge  
for  $|z| < 1$  than  $\sum a_n$ .

If  $\sum a_n z^n$  converges for  $|z| < 1$ ,  
and  $\lim_{x \rightarrow 1^-} f(x) = L$ ,

say that  $a_n$  is Abel summable, with Abel sum  $L$

How does Abel summability relate to usual summability?

# Quiz

# Tauber's theorem

Suppose  $a_n$  is Abel summable, and  $\lim_{n \rightarrow \infty} n a_n = 0$ .  
Then  $\sum a_n$  converges in the usual sense.

# Tauber's theorem

Suppose  $a_n$  is Abel summable, and  $\lim_{n \rightarrow \infty} n a_n = 0$ .  
Then  $\sum a_n$  converges in the usual sense.

There are other such theorems,  
known as Tauberian theorems  
(and so-called Abelian theorems,  
which go the other way).

# Divergent series

A **summability method** associates to  $\sum a_n$  a sum  $L$ .

# Divergent series

A **summability method** associates to  $\sum a_n$  a sum  $L$ .

A summability method is *regular*  
if it yields the honest value for convergent series.

# Divergent series

A **summability method** associates to  $\sum a_n$  a sum  $L$ .

A summability method is *regular*  
if it yields the honest value for convergent series.

Abel's theorem: Abel summability is regular.

Use Abel summation to “sum”

$$1 - 2 + 3 - 4 + 5 - \dots =$$



Use Abel summation to “sum”

$$1 - 2 + 3 - 4 + 5 - \dots =$$

Amusingly, consider  $(1 - 1 + 1 - 1 + \dots)^2$ .

Define  $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$ .

Define  $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$ .

$$2^{-s}\zeta(s) = \sum (2n)^{-s}.$$

Define  $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$ .

$$2^{-s}\zeta(s) = \sum (2n)^{-s}.$$

$$(1 - 2 \cdot 2^{-s})\zeta(s) = \sum (-1)^{n+1} n^{-s}.$$

Define  $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$ .

$$2^{-s}\zeta(s) = \sum (2n)^{-s}.$$

$$(1 - 2 \cdot 2^{-s})\zeta(s) = \sum (-1)^{n+1} n^{-s}.$$

When  $s = -1$ , this is

$$(1 - 2 \cdot 2^1)\zeta(-1) = 1 - 2 + 3 - 4 + \cdots = 1/4$$

Define  $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$ .

$$2^{-s}\zeta(s) = \sum (2n)^{-s}.$$

$$(1 - 2 \cdot 2^{-s})\zeta(s) = \sum (-1)^{n+1} n^{-s}.$$

When  $s = -1$ , this is

$$(1 - 2 \cdot 2^1)\zeta(-1) = 1 - 2 + 3 - 4 + \cdots = 1/4$$

so  $1 + 2 + 3 + 4 + \cdots = -1/12$ .

So what about  $1 + 2 + 3 + 4 + \dots$ ?

So what about  $1 + 2 + 3 + 4 + \cdots$ ? Define  $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$  when the real part of  $s$  is larger than 1, but there is an analytic function agreeing with  $\zeta(s)$  with  $\zeta(-1) = -1/12$ .



What about  $1 + 2 + 4 + 8 + 16 + \dots$ ?

What about  $1 + 2 + 4 + 8 + 16 + \dots$ ?

Consider  $f(x) = 1/(1 - 2x)$ .

What about  $1 + 2 + 4 + 8 + 16 + \dots$ ?

Consider  $f(x) = 1/(1 - 2x)$ .

So  $\sum = -1$ ? Does this make any sense?