

Lecture 27: Zeroes and poles

Math 660—Jim Fowler

Wednesday, July 28, 2010

Theorem

*Suppose f and g are analytic functions.
If $f(z) = g(z)$ for $z \in A \subset \Omega$,
and A has an accumulation point in Ω ,
then $f = g$.*

Theorem

*Suppose f and g are analytic functions.
If $f(z) = g(z)$ for $z \in A \subset \Omega$,
and A has an accumulation point in Ω ,
then $f = g$.*

Analytic functions are determined by their values on sets with an accumulation point.

Singularities

Suppose $f(z)$ is analytic in a neighborhood of a , but perhaps not at a . (I.e., analytic in a region $0 < |z - a| < \delta$).

Singularities

Suppose $f(z)$ is analytic in a neighborhood of a , but perhaps not at a . (I.e., analytic in a region $0 < |z - a| < \delta$).

Then a is an isolated singularity of f .

Singularities

Suppose $f(z)$ is analytic in a neighborhood of a , but perhaps not at a . (I.e., analytic in a region $0 < |z - a| < \delta$).

Then a is an isolated singularity of f .

If $\lim_{z \rightarrow a} f(z) = \infty$, then we say that a is a pole of $f(z)$; we can define the order of a pole by considering $1/f(z)$, which has a removable singularity at a .

Singularities

Suppose $f(z)$ is analytic in a neighborhood of a , but perhaps not at a . (I.e., analytic in a region $0 < |z - a| < \delta$).

Then a is an isolated singularity of f .

If $\lim_{z \rightarrow a} f(z) = \infty$, then we say that a is a pole of $f(z)$; we can define the order of a pole by considering $1/f(z)$, which has a removable singularity at a .

Poles are isolated, because zeroes are.

Meromorphic functions

A function f which is analytic in Ω , except at points which are poles, is called *meromorphic* in Ω .

Meromorphic functions

A function f which is analytic in Ω , except at points which are poles, is called *meromorphic* in Ω .

Add, subtract, multiply, and divide (by nonzero) meromorphic functions to get new meromorphic functions.

Meromorphic functions

A function f which is analytic in Ω , except at points which are poles, is called *meromorphic* in Ω .

Add, subtract, multiply, and divide (by nonzero) meromorphic functions to get new meromorphic functions.

Think of a meromorphic function as a holomorphic function from Ω to S^2 , the Riemann sphere.

Singularities

If $\lim_{z \rightarrow a} |z - a|^\alpha |f(z)| = 0$ for some α ,

Singularities

If $\lim_{z \rightarrow a} |z - a|^\alpha |f(z)| = 0$ for some α ,
then the same is true for α replaced by a larger value,

Singularities

If $\lim_{z \rightarrow a} |z - a|^\alpha |f(z)| = 0$ for some α ,
then the same is true for α replaced by a larger value,
so the same is true for $\alpha = \text{integer}$.

Singularities

If $\lim_{z \rightarrow a} |z - a|^\alpha |f(z)| = 0$ for some α ,
then the same is true for α replaced by a larger value,
so the same is true for $\alpha = \text{integer}$.

Then $(z - a)^m f(z)$ has a removable singularity,

Singularities

If $\lim_{z \rightarrow a} |z - a|^\alpha |f(z)| = 0$ for some α ,
then the same is true for α replaced by a larger value,
so the same is true for $\alpha = \text{integer}$.

Then $(z - a)^m f(z)$ has a removable singularity,
so either $f(z) \equiv 0$ or $(z - a)^m f(z)$ has a zero of order k .

Singularities

If $\lim_{z \rightarrow a} |z - a|^\alpha |f(z)| = 0$ for some α ,
then the same is true for α replaced by a larger value,
so the same is true for $\alpha = \text{integer}$.

Then $(z - a)^m f(z)$ has a removable singularity,
so either $f(z) \equiv 0$ or $(z - a)^m f(z)$ has a zero of order k .

So for all $\alpha > m - k$, the limit vanishes,
and if $\alpha < m - k$, the limit equals infinity.

If $\lim_{z \rightarrow a} |z - a|^\alpha |f(z)| = \infty$ for some α ,

If $\lim_{z \rightarrow a} |z - a|^\alpha |f(z)| = \infty$ for some α ,
then the same is true for α replaced by a smaller value,

If $\lim_{z \rightarrow a} |z - a|^\alpha |f(z)| = \infty$ for some α ,
then the same is true for α replaced by a smaller value,
so the same is true for $\alpha = \text{integer}$.

If $\lim_{z \rightarrow a} |z - a|^\alpha |f(z)| = \infty$ for some α ,
then the same is true for α replaced by a smaller value,
so the same is true for $\alpha = \text{integer}$.

Then $(z - a)^m f(z)$ has a removable singularity,

If $\lim_{z \rightarrow a} |z - a|^\alpha |f(z)| = \infty$ for some α ,
then the same is true for α replaced by a smaller value,
so the same is true for $\alpha = \text{integer}$.

Then $(z - a)^m f(z)$ has a removable singularity,
and $(z - a)^m f(z)$ has a pole of order ℓ .

If $\lim_{z \rightarrow a} |z - a|^\alpha |f(z)| = \infty$ for some α ,
then the same is true for α replaced by a smaller value,
so the same is true for $\alpha = \text{integer}$.

Then $(z - a)^m f(z)$ has a removable singularity,
and $(z - a)^m f(z)$ has a pole of order ℓ .

So for all $\alpha < m + \ell$, the limit equals infinity,
and if $\alpha > m + \ell$, the limit vanishes.

Summarizing

So either $f(z) \equiv 0$,

or there is some h so that

if $\alpha > h$, then $\lim_{z \rightarrow a} |z - a|^\alpha |f(z)| = 0$

if $\alpha < h$, then $\lim_{z \rightarrow a} |z - a|^\alpha |f(z)| = \infty$

or maybe neither limit holds for any α .

Algebraic order

in the second case, the cut-off α is an integer.

Algebraic order

in the second case, the cut-off α is an integer.

Apply Taylor's theorem to $(z - a)^h f(z)$, to find

$$(z - a)^h f(z) = B_h + B_{h-1}(z - a) + \cdots + B_1(z - a)^{h-1} + \varphi(z)(z - a)^h$$

with $\varphi(z)$ analytic at $z = a$.

Algebraic order

in the second case, the cut-off α is an integer.

Apply Taylor's theorem to $(z - a)^h f(z)$, to find

$$(z - a)^h f(z) = B_h + B_{h-1}(z - a) + \cdots + B_1(z - a)^{h-1} + \varphi(z)(z - a)^h$$

with $\varphi(z)$ analytic at $z = a$.

Divide both sides by $(z - a)^h$, to get

$$f(z) = \frac{B_h}{(z - a)^h} + \frac{B_{h-1}}{(z - a)^{h-1}} + \cdots + \varphi(z)$$

The part before $\varphi(z)$ is the **singular part** of f .

Algebraic order

in the second case, the cut-off α is an integer.

Apply Taylor's theorem to $(z - a)^h f(z)$, to find

$$(z - a)^h f(z) = B_h + B_{h-1}(z - a) + \cdots + B_1(z - a)^{h-1} + \varphi(z)(z - a)^h$$

with $\varphi(z)$ analytic at $z = a$.

Divide both sides by $(z - a)^h$, to get

$$f(z) = \frac{B_h}{(z - a)^h} + \frac{B_{h-1}}{(z - a)^{h-1}} + \cdots + \varphi(z)$$

The part before $\varphi(z)$ is the **singular part** of f .

A pole comes with a well-defined singular part, and if you subtract off the singular part, you are left with an analytic function.

Essential singularity

If there is no such α satisfying those limits, then we say it is an essential singularity.

Theorem

An analytic function comes arbitrarily close to every complex value in a neighborhood of an essential singularity.

