



The people



Christine Tedijanto



Ed Goldstein



Yonatan Grad



Marc Lipsitch

Science

Contents ▾News ▾Careers ▾Journals ▾

SHARE

REPORT

f

Twitter

in

Reddit

Email

Projecting the transmission dynamics of SARS-CoV-2 through the postpandemic period

ID Stephen M. Kissler^{1,*}, ID Christine Tedijanto^{2,*}, ID Edward Goldstein², ID Yonatan H. Grad^{1,†,‡}, ID Marc Lipsitch^{2,†,‡}

+ See all authors and affiliations

Science 22 May 2020:
Vol. 368, Issue 6493, pp. 860-868
DOI: 10.1126/science.abb5793

Article

Figures & Data

Info & Metrics

eLetters


PDF

What happens next?


Four months into the severe acute respiratory syndrome–coronavirus 2 (SARS-CoV-2) outbreak, we still do not know enough about postrecovery immune protection and environmental and seasonal influences on transmission to predict transmission dynamics accurately. However, we do know that humans are seasonally afflicted by other, less severe coronaviruses. Kissler *et al.* used existing data to build a deterministic model of multiyear interactions between existing coronaviruses, with a focus on the United States, and used this to project the potential epidemic dynamics and pressures on critical care capacity over the next 5 years. The long-term dynamics of SARS-CoV-2 strongly depends on immune responses and immune cross-reactions between the coronaviruses, as well as the timing of introduction of the new virus into a population. One scenario is that a resurgence in SARS-CoV-2 could occur as far into the future as 2025.

Science, this issue p. 860

The people



Christine Tedijanto



Ed Goldstein



Yonatan Grad



Marc Lipsitch

Science

Contents ▾News ▾Careers ▾Journals ▾

SHARE

REPORT

f

t

in

g+

e

Projecting the transmission dynamics of SARS-CoV-2 through the postpandemic period

ID

Stephen M. Kissler^{1,*},

ID

Christine Tedijanto^{2,*},

ID

Edward Goldstein²,

ID

Yonatan H. Grad^{1,†,‡},

ID

Marc Lipsitch^{2,†,‡}

+ See all authors and affiliations

Science 22 May 2020:
Vol. 368, Issue 6493, pp. 860-868
DOI: 10.1126/science.abb5793

Article

Figures & Data

Info & Metrics

eLetters

PDF

What happens next?

Four months into the severe acute respiratory syndrome–coronavirus 2 (SARS-CoV-2) outbreak, we still do not know enough about postrecovery immune protection and environmental and seasonal influences on transmission to predict transmission dynamics accurately. However, we do know that humans are seasonally afflicted by other, less severe coronaviruses. Kissler *et al.* used existing data to build a deterministic model of multiyear interactions between existing coronaviruses, with a focus on the United States, and used this to project the potential epidemic dynamics and pressures on critical care capacity over the next 5 years. The long-term dynamics of SARS-CoV-2 strongly depends on immune responses and immune cross-reactions between the coronaviruses, as well as the timing of introduction of the new virus into a population. One scenario is that a resurgence in SARS-CoV-2 could occur as far into the future as 2025.

Science, this issue p. 860

Pre-modeling – Model building – Model analysis