NAVAL DIABETES
CLINIC & RESEARCH CENTER
(LABORATORY SERVICES)

Name	: **Mrs. SUJATHA**
Age/Gender	: **55 Y / Female**
Patient No	: **0026**
Refered By	: **Dr.REVINSELVAN**
Bill No	: **0564**
Collection Date	: **13-Oct-2026 06:20:00 AM**
Reported Date	: **13-Oct-2026 05:59:48 PM**
Report Printed on	: **16-Oct-2026 10:55:54 AM**

Test Description	RESULT	UNITS	Reference Range
	BIOCHEMISTRY		
FASTING BLOOD SUGAR (FBS)			
Glucose Fasting	**158.0** ↑	mg/dl	60 - 110
Method : GOD POD			
Interpretation : Fasting Blood Sugar more than 126 mg/dl on more than one occasion can indicate Diabetes Mellitus.			
BLOOD SUGAR POST PRANDIAL (BSPP)			
Glucose Post Prandial	**174.0** ↑	mg/dl	80 - 140
Method : GOD POD-Endpoint			
Interpretation : A postprandial glucose reading of 140-199 mg/dl indicates prediabetes. A postprandial reading over 200 mg/dl indicates diabetes.			
GLYCOSYLATED HAEMOGLOBIN (GHB / HBA1c)			
HbA1C	**13.4** ↑	%	Below 6.0% - Normal Value 6.0% - 7.0% - Good Control 7.0% - 8.0% - Fair Control 8.0% - 10% - Unsatisfactory Control Above 10% - Poor Control
AVERAGE BLOOD GLUCOSE (ABG)	337.75	mg/dl	90 - 120 mg/dl : Excellent Control 121 - 150 mg/dl : Good Control 151 - 180 mg/dl : Average Control 181 - 210 mg/dl : Action Suggested > 211 mg/dl : Panic Value

Note

Glycosylated hemoglobin(hemoglobin A1c,HbA1c,A1C,or Hb1c, sometimes also HbA1c) is a form of hemoglobin used primary to identify the average plasma glucose concentration over prolonged periods of time. It is formed in a non-enzymatic pathway by hemoglobin's normal exposure to high plasma levels of glucose. Glycation of hemoglobin has been associated with cardiovascular disease, nephropathy and retinopathy in diabetes mellitus. Monitoring the Hb1C in type-1 diabetic patients may improve treatment. HbA1c is a weighted average of blood glucose levels during the preceding 120 days, which is the average life span of red blood cells. A large change in mean blood glucose can increase HbA1c levels within 1-2 weeks. Sudden changes in HbA1c may occur because recent changes in blood glucose levels contribute relatively more to the final HbA1c levels than earlier' events. For instance, mean blood

A. Sharmila

Dr.A.Sharmila, MBBS,MD,DNB(Pathology)
Post doctoral fellowship in molecular pathology
(CMC,Vellore)

69/25, Naval Complex, S.P.Pudur,
Paramathi Road, Namakkal - 637 001
91502 40797
revinprakash14@gmail.com

NAVAL DIABETES
CLINIC & RESEARCH CENTER
(LABORATORY SERVICES)

Name	: **Mrs. SUJATHA**
Age/Gender	: **55 Y / Female**
Patient No	: **0026**
Refered By	: **Dr.REVINSELVAN**
Bill No	: **0564**
Collection Date	: **13-Oct-2026 06:20:00 AM**
Reported Date	: **13-Oct-2026 05:59:48 PM**
Report Printed on	: **16-Oct-2026 10:55:54 AM**

glucose levels in the 30 days immediately preceding blood sampling contribute 50% to the HbA1c level, whereas glucose levels in the preceding 90-120 day period contribute only 10%. Thus, it does not make 120 days to detect a clinically meaningful change in HbA1c following a significant change in mean plasma glucose level.

**** END OF REPORT ****

Dr.A.Sharmila, MBBS,MD,DNB(Pathology)
Post doctoral fellowship in molecular pathology
(CMC,Vellore)