

# On the classical and quantum dynamics of a class of nonpolynomial oscillators

V. Chithiika Ruby and M. Lakshmanan

*Department of Nonlinear Dynamics, School of Physics,  
Bharathidasan University, Tiruchirapalli - 620 024, India.*

## Abstract

We consider two one dimensional nonlinear oscillators, namely (i) Higgs oscillator and (ii) a  $k$ -dependent nonpolynomial rational potential, where  $k$  is the constant curvature of a Riemannian manifold. Both the systems are of position dependent mass form,  $m(x) = \frac{1}{(1 + kx^2)^2}$ , belonging to the quadratic Liénard type nonlinear oscillators. They admit different kinds of motions at the classical level. While solving the quantum versions of the systems, we consider a generalized position dependent mass Hamiltonian in which the ordering parameters of the mass term are treated as arbitrary. We observe that the quantum version of the Higgs oscillator is exactly solvable under appropriate restrictions of the ordering parameters, while the second nonlinear system is shown to be quasi exactly solvable using the Bethe ansatz method in which the arbitrariness of ordering parameters also plays an important role to obtain quasi-polynomial solutions. We extend the study to three dimensional generalizations of these nonlinear oscillators and obtain the exact solutions for the classical and quantum versions of the three dimensional Higgs oscillator. The three dimensional generalization of the quantum counterpart of the  $k$ -dependent nonpolynomial potential is found out to be quasi exactly solvable.

## I. INTRODUCTION

The classical nonlinear oscillator introduced by Mathews and Lakshmanan and characterized by the Lagrangian,

$$L = \frac{1}{2} \frac{(\dot{x}^2 - \omega_0^2 x^2)}{(1 + \lambda x^2)}, \quad (1)$$

where  $\omega_0$  and  $\lambda$  are parameters, has drawn considerable interest in the literature due to its unique simple harmonic oscillatory behavior [1, 2]. The nonlinear system (1) and its generalizations are continuously studied for many aspects such as 3-dimensional [3] and  $d$ -dimensional generalizations [4, 5] and rational extensions of the potentials [6]. Higgs geometrically interpreted the above nonlinear system which emerged while generalizing the Cartesian coordinates of Euclidean geometry. He also obtained a non-polynomial Lagrangian while defining the harmonic oscillator in generalized coordinates which arises on the gnomonic projection onto the tangent plane from the center of the sphere [7]. The one-dimensional version of the Higgs oscillator is described by the Lagrangian,

$$L = \frac{\dot{x}^2}{2(1 + kx^2)^2} - \frac{\omega_0}{2} x^2, \quad (2)$$

where  $k$  is a parameter related to the curvature. The Higgs oscillator has also attracted considerable attention since its introduction to the literature [4, 8].

Both the above nonlinear systems are unique examples for the dynamics of systems in curved space. Interestingly, Carinena et al. obtained the two nonlinear systems while studying the dynamics of harmonic oscillator on the two dimensional Riemann spaces ( $M_k^2$ ) of constant curvature (that is, sphere  $S^2$  and hyperbolic plane  $\mathcal{H}^2$ ). In this unified approach, the curvature  $k$  was treated as a parameter and the corresponding Lagrangian of the harmonic oscillator potential in terms of geodesic polar coordinates  $(R, \Phi)$  on  $M_k^2$  is expressed as [8, 9]

$$\mathcal{L}(k) = \frac{1}{2} (v_R^2 + S_k^2(R) v_\Phi^2) - \frac{1}{2} \omega_0^2 T_k^2(R), \quad (3)$$

where  $T_k = \frac{S_k}{C_k}$ ,  $S_k$  and  $C_k$  are  $k$ -dependent trigonometric functions. They are defined as

$$C_k = \begin{cases} \cos(\sqrt{k} x), & k > 0 \\ 1, & k = 0 \\ \cosh(\sqrt{-k} x), & k < 0 \end{cases} \quad (4)$$

$$S_k = \begin{cases} \frac{1}{\sqrt{k}} \sin(\sqrt{k} x), & k > 0 \\ x, & k = 0 \\ \frac{1}{\sqrt{-k}} \sinh(\sqrt{-k} x), & k < 0 \end{cases} \quad (5)$$

Here  $k > 0$  denotes the spherical plane, whereas  $k < 0$  denotes hyperbolic plane and  $k = 0$  denotes the Euclidean plane.

(1) On the transformation  $(R, \Phi) \rightarrow (r', \phi)$  as  $r' = T_k(R)$ ,  $\Phi = \phi$ , the Lagrangian  $\mathcal{L}(k)$  becomes

$$\mathcal{L}_H(k) = \frac{1}{2} \left( \frac{v_{r'}^2}{(1 + kr'^2)^2} + \frac{v_\phi^2}{(1 + kr'^2)} \right) - \frac{1}{2} \omega_0^2 r'^2, \quad (6)$$

which is known as the two dimensional Higgs oscillator [7].

(2) Similarly, on the transformation  $(R, \Phi) \rightarrow (r, \phi)$  as  $r = S_k(R)$ ,  $\Phi = \phi$  with  $k = -\lambda$ , the Lagrangian  $\mathcal{L}(k)$  becomes

$$\mathcal{L}_M(\lambda) = \frac{1}{2} \left( \frac{v_r^2}{(1 + \lambda r^2)^2} + r^2 v_\phi^2 \right) - \frac{\omega_0^2}{2} \frac{r^2}{(1 + \lambda r^2)}, \quad (7)$$

which is known as the two dimensional Mathews-Lakshmanan oscillator [1].

Classically, the kinetic energy term of the Mathews-Lakshmanan oscillator (MLO) depicts the dynamics of a nonlinear  $SU(2)XSU(2)$ - chirally invariant Lagrangian expressed in Gasiorowicz-Geffen co-ordinates [10],

$$\mathcal{L} = \frac{1}{2} \left[ (\partial_\mu \Phi) \cdot (\partial_\mu \Phi) + \frac{\lambda (\Phi \cdot \partial_\mu \Phi) (\Phi \cdot \partial_\mu \Phi)}{(1 - \lambda \Phi^2)} \right], \quad (8)$$

where  $\lambda = \text{constant}$ , and is expressed in flat space  $\Phi \rightarrow \mathbf{q}$ .

In another coordinate system, namely Schwinger coordinates, the same nonlinear chiral system (8) is expressed as

$$\mathcal{L} = \frac{1}{2} \left[ (\partial_\mu \Phi) \cdot (\partial_\mu \Phi) - \frac{k}{(1 + k\Phi^2)} \left( \Phi^2 (\partial_\mu \Phi) \cdot (\partial_\mu \Phi) + \frac{(\Phi \cdot \partial_\mu \Phi) (\Phi \cdot \partial_\mu \Phi)}{(1 + k\Phi^2)} \right) \right], \quad (9)$$

where  $k$  is constant [10].

In this work, we are interested to study the nonlinear systems associated with the Lagrangian (9), in the flat space, since it is related to the Higgs oscillator (2). In flat space, the Lagrangian takes the form

$$L = \frac{1}{2} \left[ \dot{\mathbf{q}}^2 - \frac{k}{(1 + kq^2)} \left( q^2 \dot{\mathbf{q}}^2 + \frac{(\mathbf{q} \cdot \dot{\mathbf{q}})^2}{(1 + kq^2)} \right) \right]. \quad (10)$$

We consider the one-dimensional version of the nonlinear chiral system (10) corresponding to two different additional potentials,  $V_{1,2}(x)$ ,

$$L = \frac{1}{2} \left[ \frac{\dot{x}^2}{(1 + kx^2)^2} \right] - V_{1,2}(x), \quad (11)$$

where

$$V_1(x) = \frac{\omega_0^2 x^2}{2}, \quad V_2(x) = \frac{\omega_0^2 x^2}{2(1 + kx^2)^2}. \quad (12)$$

Here, the potential  $V_1(x)$  corresponds to the Higgs oscillator, while  $V_2(x)$  is a non-polynomial potential that includes merely the mass term in its potential as in the case of Mathews-Lakshmanan oscillator (1).

The conjugate momentum for these systems (11) is

$$p = \frac{\partial L}{\partial \dot{x}} = \frac{\dot{x}}{(1 + kx^2)^2}, \quad (13)$$

and the corresponding Hamiltonian can be written as

$$H = \frac{(1 + kx^2)^2}{2} p^2 + V_{1,2}(x). \quad (14)$$

The above Hamiltonians are of the form of position dependent mass systems,  $H = \frac{p^2}{2m(x)} + V(x)$ , where  $m(x)$  is the position dependent mass term.

In this work, we intend to study both the classical and quantum solvability of these one-dimensional systems. Both the above systems with the potentials  $V_1(x)$  and  $V_2(x)$  are classically of quadratic Liénard-type one-dimensional nonlinear oscillators. They can be exactly solved classically. Here, the mass of the systems is position dependent,  $m(x) = \frac{1}{(1 + kx^2)^2}$ , where  $k$  is constant. We have first employed semiclassical treatment to analyze the quantum solvability of the nonlinear systems. Recently, we studied the problem of removal of ordering ambiguity in position dependent mass quantum systems characterized by a generalized position dependent mass Hamiltonian which generalizes a number of Hermitian as well as non-Hermitian ordered forms of the Hamiltonian [11, 12]. Motivated by this study, while

analyzing the quantum dynamics of the nonlinear systems endowed with one-dimensional potentials, we use a generalized  $2N$ -parameter kinetic energy operator in which we keep the ordering parameters as arbitrary. Interestingly the above two systems can also be generalized to higher dimensions as well. Particularly we will consider the three dimensional generalizations of these systems both classically and quantum mechanically.

The manuscript is structured into three major segments as the classical, semiclassical and quantum studies of the above two one-dimensional nonlinear systems and their corresponding three dimensional generalizations. In sec. II, we study the classical dynamics of the one dimensional systems. The quantum solvability of these systems is examined through semiclassical technique, namely the modified Bohr-Sommerfeld quantization rule in Sec. III. In Sec. IV, we use the form of general ordered position dependent mass Hamiltonian and analyze the solvability of the nonlinear systems. We further extend our study to the three dimensional generalization of the two nonlinear systems in sections VB and VC. In Sec. VI, we present a summary of the results on the two systems.

## II. CLASSICAL DYNAMICS OF THE ONE DIMENSIONAL NONLINEAR CHIRAL MODELS

Even though the two nonlinear systems have the same kinetic energy term, the difference in their potentials will lead to different solutions. We first consider the Higgs oscillator and then the new nonlinear oscillator system.

### A. One dimensional Higgs oscillator

The Lagrangian of the Higgs oscillator is of the form,

$$L = \frac{\dot{x}^2}{2(1+kx^2)^2} - \frac{\omega_0^2}{2}x^2, \quad (2)$$

where  $k$  and  $\omega_0$  are parameters.

The classical dynamics of the system (2) is described by the equation of motion,

$$\ddot{x} - \frac{2kx}{(1+kx^2)}\dot{x}^2 + \omega_0^2(1+kx^2)^2x = 0. \quad (15)$$

It is of the form of the quadratic Liénard type nonlinear equation,

$$y'' + f(y)y'^2 + g(y) = 0. \quad (16)$$

By using the multiplication factor,  $\dot{x}$ , Eq. (15) is integrated to be

$$\frac{\dot{x}^2}{(1 + kx^2)^2} + \omega_0^2 x^2 = \epsilon = \text{constant}. \quad (17)$$

On implementation of the simple transformation,

$$y = \frac{x}{\sqrt{1 + kx^2}}, \quad (18)$$

equation (17) is reduced to the form

$$\dot{y}^2 + (\omega_0^2 + \epsilon k)y^2 = \epsilon. \quad (19)$$

When  $k > 0$ ,  $y$  is real and bounded between 0 and  $\frac{1}{\sqrt{k}}$  for all values of  $x$ , that is  $-\infty < x < \infty$  whereas for  $k < 0$ ,  $y \in (0, \infty)$  in the region  $-\frac{1}{\sqrt{|k|}} < x < \frac{1}{\sqrt{|k|}}$ .

On integrating (19), we obtain

$$\int \frac{dy}{\sqrt{1 - \frac{(\omega_0^2 + \epsilon k)}{\epsilon} y^2}} = \sqrt{\epsilon} t + C, \quad (20)$$

where  $C$  is a constant. Consequently, we get

$$y(t) = A \sin(\Omega t + C), \quad (21)$$

where  $A^2 = \frac{\epsilon}{(\omega_0^2 + \epsilon k)}$  is the square of the amplitude and considered to be real and  $\Omega^2 = \frac{\omega_0^2}{(1 - kA^2)}$  is the square of the frequency. The solution of equation (15) can be obtained from (18) and (21) as

$$x(t) = \frac{A \sin(\Omega t + C)}{\sqrt{1 - kA^2 \sin^2(\Omega t + C)}}, \quad (22)$$

where  $|A| < \frac{1}{\sqrt{k}}$  for  $k > 0$  and  $x(t)$  is periodic for all the values of  $A$  when  $k < 0$ . The explicit expression (22) of the one dimensional Higgs oscillator consisting of trigonometric function depicts not only the solvability of the nonlinear system classically but also motivates us to study the exact solvability of its quantum version. The variable  $x(t)$  is plotted in figure 1.

## B. One dimensional nonlinear oscillator, $V_2(x)$

Let us consider the Lagrangian of the potential,  $V_2(x) = \frac{\omega_0^2 x^2}{2(1 + kx^2)^2}$ , which is

$$L = \frac{\dot{x}^2 - \omega_0^2 x^2}{2(1 + kx^2)^2}, \quad (23)$$




FIG. 1: The plot of  $x(t)$  obtained in (22) of the system (2) for  $k = 0.5$ ,  $A = 1$  and  $\Omega = 0.5$ .

where  $k$  and  $\omega_0$  are system parameters. The corresponding equation of motion is

$$\ddot{x} - \frac{2kx}{(1+kx^2)}\dot{x}^2 + \omega_0^2 \frac{(1-kx^2)}{(1+kx^2)}x = 0, \quad (24)$$

which is also of the form of the quadratic Liénard type nonlinear equation, Eq. (16). Hence, the multiplication factor,  $\frac{\dot{x}}{(1+kx^2)^2}$ , reduces Eq. (24) to the form

$$\frac{\dot{x}^2}{(1+kx^2)^2} + \frac{\omega_0^2 x^2}{(1+kx^2)^2} = \epsilon = \text{constant}. \quad (25)$$

To solve Eq. (25), we use the transformation  $\tau = \Omega t$  and  $x(t) = AX(\tau)$ , so that it can be reexpressed as

$$\dot{X}^2(\tau) = \frac{\epsilon}{A^2 \Omega^2} + \frac{(2k\epsilon - \omega_0^2)}{\Omega^2} X^2 + \frac{\epsilon k^2 A^2}{\Omega^2} X^4, \quad (26)$$

which results on integration to

$$x(t) = A \operatorname{sn} \left( \frac{\omega_0}{1+kA^2} t, m \right) \quad (27)$$

with

$$\epsilon = \frac{\omega_0^2 A^2}{(1+kA^2)^2}, \quad m = kA^2. \quad (28)$$

Using the identity [17],  $\operatorname{sn}(u, m) = \frac{(1+m_1') \operatorname{sn}(u_1, m_1) \operatorname{cn}(u_1, m_1)}{\operatorname{dn}(u_1, m_1)}$ , where  $u = (1+m_1') u_1$  and  $m = \frac{1-m_1'}{1+m_1'}$ , and by substituting  $u_1 = \frac{\omega_0 t}{2}$  and  $m_1' = \frac{1-kA^2}{1+kA^2}$ , we can express the solution (27) as

$$x(t) = \frac{2 A \operatorname{sn}(\frac{\omega_0}{2} t, m_1) \operatorname{cn}(\frac{\omega_0}{2} t, m_1)}{(1+k A^2) \operatorname{dn}(\frac{\omega_0}{2} t, m_1)}, \quad (29)$$

where the modulus  $m_1 = \frac{2\sqrt{k}A^2}{(1+kA^2)}$  lies between  $0 \leq m_1 \leq 1$ .

The solutions (27) and (29) include the Jacobian elliptic functions  $sn$ ,  $cn$  and  $dn$  which ensure us that the quantum counterpart of system may be quasi exactly solvable. When  $k \rightarrow 0$ , the modulus  $m = 0$  which reduces  $x(t)$  (vide Eq. (27) and (29)) to be

$$x(t) = A \sin(\omega_0 t), \quad (30)$$

the solution of the linear harmonic oscillator as expected. The dynamics of the nonlinear system is dependent on  $m_1$  which is depicted in Figure. 2.




FIG. 2: The plot of  $x(t)$ , (27), of the system (23) with different values of  $m_1$  (a)  $m_1 = 0$ , (b)  $m_1 = 0.5$  and (c)  $m_1 = 1$  for  $A = 1$  and  $\Omega = 0.5$ .

### III. SEMICLASSICAL QUANTIZATION

The semiclassical study often acts as a bridge between the study of classical dynamics and full quantum investigation of a system and often helps in the possibility of solving the quantum version of the system exactly. To quantize the system semiclassically, we use the modified Bohr-Sommerfeld quantization rule [13], namely

$$\oint p dx = \left(n + \frac{1}{2}\right) h, \quad n = 0, 1, 2, 3, \dots, \quad (31)$$

where  $h$  is the Planck's constant and  $n$  is any non-negative integer and the integration is carried out over a closed orbit in the  $(x, p)$  space.



### A. Higgs oscillator

We can express the canonically conjugate momentum of the system (2) by using the solution (22) as

$$p = A\Omega \cos(\Omega t + C) \left(1 - kA^2 \sin(\Omega t + C)\right)^{\frac{1}{2}}. \quad (32)$$

We first determine the energy of the system  $E = \frac{\epsilon}{2}$  from (17) or (19) using the solution (22) as

$$E = \frac{\omega_0^2 A^2}{1 - kA^2}. \quad (33)$$

Using  $x(t)$ , vide (22) in (13), we can express the integral in (31) to be of the form,

$$I = \epsilon \oint \frac{dt}{(1 + (1 + kA^2) \tan^2(\Omega t + C))} \quad (34)$$

$$= \frac{-\omega_0}{k} \left(1 - \frac{1}{\sqrt{1 - kA^2}}\right), \quad (35)$$

which on being substituted in (31) yields

$$\frac{\omega_0^2 A^2}{(1 - kA^2)} = 2 \left(n + \frac{1}{2}\right) \hbar\omega_0 + \left(n^2 + n + \frac{1}{4}\right) \hbar^2 k. \quad (36)$$

On substituting (36) in (33), we obtain the semiclassical energy spectrum of the Higgs oscillator (2) as

$$E_n = \left(n + \frac{1}{2}\right) \hbar\omega_0 + \left(n^2 + n + \frac{1}{4}\right) \frac{\hbar^2 k}{2}, \quad n = 0, 1, 2, 3, \dots \quad (37)$$

Here the energy levels are quadratic as obtained by Higgs for the two dimensional case of the system (2) [7]. In the limit  $k \rightarrow 0$ , the energy levels (37) become the energy spectrum of the harmonic oscillator as the system (2) is reduced to that of the linear harmonic oscillator.

### B. Semiclassical method for the second system (23)

We now analyze the solvability of the system (23) semiclassically. To do so, we use the momentum of the system (13) as

$$p = \frac{\partial L}{\partial \dot{x}} = \frac{\dot{x}}{(1 + kx^2)^2}.$$

By using the solution (27), we can evaluate the integral in (31) as

$$I = \frac{\omega_0^2 A^2}{(1 + kA^2)^2} \int_0^{4K(m_1)} cn^2(\omega_0 t, m_1) dt, \quad (38)$$

$$= \frac{\omega_0}{4k} (E(m_1) - K(m_1)), \quad (39)$$

where  $y = cn(\omega_0 t, m_1)$  and  $K(m_1)$  and  $E(m_1)$  are complete elliptic integrals of the first kind and second kind, respectively. In terms of  $m$ , the relation (39) can be expressed as

$$I = \frac{\omega_0}{2k} \left( \frac{E(m)}{(1+m)} - K(m) \right). \quad (40)$$

Then,  $I$  can be related with quantum number  $n$  through (31) as

$$\frac{\omega_0}{2k} \left( \frac{E(m)}{(1+m)} - K(m) \right) = \left( n + \frac{1}{2} \right) h, \quad n = 0, 1, 2, 3, \dots \quad (41)$$

This relation (41) implies that one has to solve for  $m$  (and then  $m_1$ ) and so for  $A$  implicitly, which implies that only a numerical solution can be given but not an exact solution in general.

#### IV. QUANTUM SOLVABILITY OF THE NONLINEAR SYSTEMS

Most of the nonlinear oscillators defined in curved space are equivalently observed as position dependent mass systems on quantum transition. They also include the quadratic Liénard type nonlinear oscillators, for example Mathews-Lakshmanan (ML) oscillator [2], its generalizations [14], its extensions [15] and Higgs oscillator [7].

The classical Hamiltonian  $H(x, p)$  corresponding to (14) is

$$H = (1 + kx^2)^2 \frac{p^2}{2} + V(x), \quad (42)$$

where  $p$  is the momentum (13).

In this section, we consider the two potentials, (1)  $V_1(x) = \frac{\omega_0^2 x^2}{2}$ , the Higgs oscillator and (2) the nonlinear system,  $V_2(x) = \frac{\omega_0^2 x^2}{2(1 + kx^2)^2}$ . The associated Hamiltonians are of the forms

$$H_1 = (1 + kx^2)^2 \frac{p^2}{2} + \frac{\omega_0^2 x^2}{2}, \quad (43)$$

and

$$H_2 = (1 + kx^2)^2 \frac{p^2}{2} + \frac{\omega_0^2 x^2}{2(1 + kx^2)^2}, \quad (44)$$

respectively.

We observe that the Hamiltonian  $H(x, p)$ , specified by (42), is of position dependent mass type. To solve the system quantum mechanically, we consider the most general ordered form of the Hamiltonian operator that allows many number of possible mixtures of the fundamental term,  $m^\alpha \hat{p} m^\beta \hat{p} m^\gamma$ , where  $\hat{p}$  is momentum operator,  $\hat{p} = -i \hbar \frac{d}{dx}$ . It reads as [16]

$$\hat{H} = \frac{1}{2} \sum_{i=1}^N w_i m^{\alpha_i} \hat{p} m^{\beta_i} \hat{p} m^{\gamma_i} + V(x), \quad (45)$$

where  $N$  is an arbitrary positive integer, and the ordering parameters  $(\alpha_i, \beta_i, \gamma_i)$  should satisfy the constraint  $\alpha_i + \beta_i + \gamma_i = -1$ ,  $i = 1, 2, 3, \dots, N$ , and  $w_i$ 's are real weights which are summed to be 1. The above form globally connects all the Hermitian orderings and also provides a complete classification of Hermitian and non-Hermitian orderings [16]. The kinetic energy operator of  $\hat{H}$  in (45) possesses  $2N$  free ordering parameters.

The operator (45) is not Hermitian in general and can be related with its Hermitian counterpart through a similarity transformation. The Hermitian Hamiltonian  $\hat{H}_{her}$  is deduced by performing the transformation

$$\hat{H}_{her} = m^\eta \hat{H}_{non} m^{-\eta}, \quad (46)$$

with  $2\eta = \bar{\gamma} - \bar{\alpha}$ , where  $\bar{X} = (\bar{\alpha}, \bar{\beta}, \bar{\gamma})$  denotes the weighted mean value,  $\bar{X} = \sum_{i=1}^N w_i X_i$  and  $-\bar{\beta} = 1 + \bar{\alpha} + \bar{\gamma}$  (ref. [11]). The Hermitian Hamiltonian (46) for a potential  $V(x)$  can be expressed as

$$\hat{H}_{her} = \frac{1}{2} \hat{p} \frac{1}{m} \hat{p} + \frac{\hbar^2}{2} \left[ \left( \frac{\bar{\alpha} + \bar{\gamma}}{2} \right) \frac{d^2}{dx^2} \left( \frac{1}{m} \right) + \left( \bar{\alpha} \bar{\gamma} + \frac{1}{4} (\bar{\gamma} - \bar{\alpha})^2 \right) \left( \frac{d}{dx} \left( \frac{1}{m} \right) \right)^2 m \right] + V(x). \quad (47)$$

Since  $[\hat{x}, \hat{p}] = i\hbar$ , we assume the coordinate representation of  $\hat{p} = -i\hbar \frac{d}{dx}$ .

To solve the quantum system (42), we start with the time-independent Schrödinger equation for the Hamiltonian (47),

$$\psi'' - \frac{m'}{m} \psi' + \left( \left( \frac{\bar{\alpha} + \bar{\gamma}}{2} \right) \frac{m''}{m} - \left( \bar{\alpha} \bar{\gamma} + \bar{\gamma} + \bar{\alpha} + \frac{1}{4} (\bar{\gamma} - \bar{\alpha})^2 \right) \frac{m'^2}{m^2} \right) \psi + \frac{2m}{\hbar^2} (E - V(x)) \psi = 0, \quad (48)$$

where  $' = \frac{d}{dx}$  and obtain the solution for arbitrary values of the ordering parameters  $\bar{\alpha}$  and  $\bar{\beta}$ .

### A. Higgs oscillator, $H_1$

For Higgs oscillator, the Hamiltonian is of the form,

$$H = (1 + kx^2)^2 \frac{p^2}{2} + \frac{\omega_0^2}{2} x^2. \quad (43)$$

The generalized time-independent Schrödinger equation (48) for the Hamiltonian (43) becomes

$$\psi'' + \frac{4kx}{(1 + kx^2)} \psi' + \left( \frac{2\eta_1 k - \frac{\omega_0^2}{\hbar^2 k}}{(1 + kx^2)} + \frac{\frac{2E}{\hbar^2} + 4\eta_2 k + \frac{\omega_0^2}{\hbar^2 k}}{(1 + kx^2)^2} \right) \psi = 0, \quad \left( ' = \frac{d}{dx} \right) \quad (49)$$

where

$$\eta_1 = 5(\bar{\alpha} + \bar{\gamma}) - 8 \left( \bar{\alpha}\bar{\gamma} + \bar{\alpha} + \bar{\gamma} + \frac{1}{4}(\bar{\gamma} - \bar{\alpha})^2 \right), \quad (50)$$

$$\eta_2 = -3(\bar{\alpha} + \bar{\gamma}) + 4 \left( \bar{\alpha}\bar{\gamma} + \bar{\alpha} + \bar{\gamma} + \frac{1}{4}(\bar{\gamma} - \bar{\alpha})^2 \right). \quad (51)$$

We solve the equation (49) for both the cases (i)  $k < 0$  and (ii)  $k > 0$ .

#### (i) Positive values of $k$

On using the transformation,

$$z = \frac{k x^2}{1 + k x^2}, \quad \psi(z) = (1 - z)^d \phi(z), \quad (52)$$

we can express Eq. (49) as

$$z(1 - z)\phi''(z) + \left( \frac{1}{2} - 2dz \right) \phi'(z) + \left[ \frac{\frac{\eta_1}{2} + d \left( d - \frac{3}{2} \right) - \frac{\mu^2}{4}}{1 - z} - \xi \right] \phi(z) = 0, \quad \left( ' = \frac{d}{dz} \right) \quad (53)$$

where

$$\xi = \frac{E}{2\hbar^2 k} + \eta_2 + \frac{\mu^2}{4} - d(d - 1), \quad (54)$$

$$\mu = \frac{\omega_0}{\hbar k}. \quad (55)$$

With  $d = \frac{3}{4} \pm \frac{1}{2} \sqrt{\mu^2 - 2\eta_1 + \frac{9}{4}}$  and  $\tilde{\mu} = \sqrt{\mu^2 - 2\eta_1 + \frac{9}{4}}$ , we can reduce Eq. (53) to the form

$$z(1 - z)\phi''(z) + \left( \frac{1}{2} - \left( \pm \tilde{\mu} + \frac{3}{2} \right) z \right) \phi'(z) - \nu \left( -\nu \pm \tilde{\mu} + \frac{1}{2} \right) \phi(z) = 0, \quad (56)$$

where  $\xi$  is now defined as

$$\xi = -\nu \left( -\nu \pm \tilde{\mu} + \frac{1}{2} \right), \quad \nu \in \mathbb{Z}. \quad (57)$$

Equation (56) is of the form of a hypergeometric equation,

$$y(1-y)w''(y) + (c - (a+b+1)y)w'(y) - abw(y) = 0, \quad |y| < 1, \quad (58)$$

which admits solution near  $y = 0$  as

$$w(y) = C_1 {}_2F_1(a, b; c; y) + C_2 y^{1-c} {}_2F_1(a-c+1, b-c+1; 2-c; y). \quad (59)$$

On comparing (53) and (56) we have

$$a = \nu, \quad b = -\nu \pm \tilde{\mu} + \frac{1}{2}, \quad c = \frac{1}{2}. \quad (60)$$

Hence the general solution of (56) can be written as

$$\phi(z) = C_1 {}_2F_1\left(\nu, -\nu \pm \tilde{\mu} + \frac{1}{2}; \frac{1}{2}; z\right) + C_2 z^{1/2} {}_2F_1\left(\nu + \frac{1}{2}, -\nu \pm \tilde{\mu} + 1; \frac{3}{2}; z\right), \quad |z| < 1. \quad (61)$$

Here,  $C_1$  and  $C_2$  are arbitrary constants.

On using (61) in (52), we can express the solution for (49) in terms of  $z$  as

$$\psi(z) = (1-z)^{\frac{3}{4} \pm \frac{\tilde{\mu}}{2}} \left[ C_1 {}_2F_1\left(\nu, -\nu \pm \tilde{\mu} + \frac{1}{2}; \frac{1}{2}; z\right) + C_2 z^{1/2} {}_2F_1\left(\nu + \frac{1}{2}, -\nu \pm \tilde{\mu} + 1; \frac{3}{2}; z\right) \right]. \quad (62)$$

To get bound state solutions, we apply the boundary condition  $\psi(z) = 0$  at  $z = 1$  on (62) which implies that  $d$  should be positive, that is,  $\frac{3}{4} + \frac{\tilde{\mu}}{2}$ . To obtain polynomial solution, we use the identity [17],

$$\mathcal{N} {}_2F_1\left(2a, 2b; a+b+\frac{1}{2}, \frac{1-\sqrt{z}}{2}\right) = C_1 {}_2F_1\left(a, b; \frac{1}{2}; z\right) + C_2 z^{1/2} {}_2F_1\left(a+\frac{1}{2}, b+\frac{1}{2}; \frac{3}{2}; z\right), \quad (63)$$

where  $C_1$  and  $C_2$  take the values

$$C_1 = \frac{\Gamma(a+b+\frac{1}{2})\sqrt{\pi}}{\Gamma(a+\frac{1}{2})\Gamma(b+\frac{1}{2})} \mathcal{N}, \quad C_2 = \frac{\Gamma(a+b+\frac{1}{2})\sqrt{\pi}}{\Gamma(b)\Gamma(a)} \mathcal{N}. \quad (64)$$

Here  $\mathcal{N}$  is the normalization constant. Then, the solution (62) can be expressed as

$$\psi(z) = \mathcal{N} (1-z)^{\frac{3}{4} \pm \frac{\tilde{\mu}}{2}} {}_2F_1\left(2\nu, -2\nu + 2\tilde{\mu} + 1; \tilde{\mu} + 1; \frac{1-\sqrt{z}}{2}\right). \quad (65)$$

The constants (64) are now of the forms,

$$C_1 = \frac{\Gamma(\tilde{\mu} + 1)\sqrt{\pi}}{\Gamma(\nu + \frac{1}{2})\Gamma(-\nu + \tilde{\mu} + 1)} \mathcal{N}, \quad C_2 = \frac{\Gamma(\tilde{\mu} + 1)\sqrt{\pi}}{\Gamma(\nu)\Gamma(-\nu + \tilde{\mu} + \frac{1}{2})} \mathcal{N}. \quad (66)$$

The auxiliary condition on the solution  $\psi(z)$  that it should be square integrable requires  $\psi(z)$  must be polynomial which can be achieved by choosing  $\nu = -\frac{n}{2}$ , where  $n$  is a positive definite integer. Hence, Eq. (62) becomes

$$\psi_n(z) = \mathcal{N}_n (1 - z)^{\frac{3}{4} + \frac{\tilde{\mu}}{2}} {}_2F_1 \left( -n, n + 2\tilde{\mu} + 1; 1 + \tilde{\mu}; \frac{1 - \sqrt{z}}{2} \right), \quad n = 0, 1, 2, 3, \dots \quad (67)$$

By using the transformation, [17]

$${}_2F_1(c - a, c - b, c, y) = (1 - y)^{a+b-c} {}_2F_1(a, b, c; y), \quad (68)$$

we can express the solution (67) as

$$\psi_n(z) = \mathcal{N}_n 2^{\tilde{\mu}} (1 - z)^{3/4} \frac{(1 - \sqrt{z})^{\tilde{\mu}/2}}{(1 + \sqrt{z})^{\tilde{\mu}/2}} {}_2F_1 \left( -n - \tilde{\mu}, n + \tilde{\mu} + 1; 1 + \tilde{\mu}; \frac{1 - \sqrt{z}}{2} \right). \quad (69)$$

Since the associated Legendre polynomial,  $P_\lambda^{-q}(\sqrt{z})$ , where  $\lambda$  and  $q$  are parameters, can be related with the hypergeometric function as

$$P_\lambda^{-q}(\sqrt{z}) = \frac{1}{\Gamma(1 + q)} \frac{(1 - \sqrt{z})^{q/2}}{(1 + \sqrt{z})^{q/2}} {}_2F_1 \left( -\lambda, \lambda + 1; 1 + q; \frac{1 - \sqrt{z}}{2} \right), \quad q > 0, \quad (70)$$

we can express the solution (69) in the form

$$\psi_n(z) = \mathcal{N}_n 2^{\tilde{\mu}} \Gamma(1 + \tilde{\mu}) (1 - z)^{\frac{3}{4}} P_{n+\tilde{\mu}}^{-\tilde{\mu}}(\sqrt{z}), \quad (71)$$

with  $\lambda = n + \tilde{\mu}$  and  $q = \tilde{\mu}$ . Then in terms of  $x$ , we have

$$\psi_n(x) = \mathcal{N}_n 2^{\tilde{\mu}} \Gamma(1 + \tilde{\mu}) (1 + k x^2)^{-\frac{3}{4}} P_{n+\tilde{\mu}}^{-\tilde{\mu}} \left( \frac{\sqrt{k} x}{\sqrt{1 + k x^2}} \right), \quad k > 0, \quad (72)$$

where  $0 < \frac{\sqrt{k} x}{\sqrt{1 + k x^2}} < 1$  provided that when  $k > 0$ ,  $-\infty < x < \infty$ . Here, the normalization constant  $\mathcal{N}_n$  can be found out as

$$1 = \int_{-\infty}^{\infty} \psi_n^*(x) \psi_n(x) dx, \quad k > 0 \quad (73)$$

$$\mathcal{N}_n = \left( \frac{\sqrt{k} \Gamma(n + 2\mu + 1) (2n + 2\mu + 1)}{n! 2} \right)^{1/2}. \quad (74)$$

The corresponding energy eigenvalues  $E_n$  can be obtained from (54) and (57) with  $\nu = -\frac{n}{2}$  as

$$\frac{E}{2\hbar^2 k} + \frac{\mu^2}{4} + \eta_2 - \left(\frac{\tilde{\mu}}{2} + \frac{3}{4}\right) \left(\frac{\tilde{\mu}}{2} - \frac{1}{4}\right) = -\nu \left(-\nu + \tilde{\mu} + \frac{1}{2}\right), \quad (75)$$

which yields

$$E_n = \left(n + \frac{1}{2}\right) \hbar \sqrt{\omega_0^2 + \hbar^2 k^2 \left(2\eta_1 - \frac{9}{4}\right)} + \left(n^2 + n + 2\bar{\alpha} + 2\bar{\gamma} + \frac{3}{2}\right) \frac{\hbar^2 k}{2}, \quad (76)$$

where  $\eta_1$  (vide Eq. (50)) is a function of the ordering parameters  $\bar{\alpha}$  and  $\bar{\gamma}$ . Here the presence of ordering parameters  $\bar{\alpha}$  and  $\bar{\gamma}$  in the energy eigenvalues (76) indicates that the different choices of ordering parameters results in different energy spectrum. Thus different choices of ordering parameters leads to different effective potentials of (43). Hence we have obtained a class of exactly solvable potentials by using this procedure.

In the limit  $k \rightarrow 0$ , we obtain the energy spectrum of the harmonic oscillator,  $E_n = \left(n + \frac{1}{2}\right) \hbar \omega_0$ , as the system (43) represents the simple harmonic oscillator in the same limit.

In the recent work [18], the general ordered non-Hermitian ordered form (45) of the Hamiltonian corresponding to the Mathews-Lakshmanan oscillator (1) has been solved exactly. The corresponding eigenfunctions are also expressed in terms of associated Legendre polynomials as

$$\psi_n = \begin{cases} N_n (1 + \lambda x^2)^{\frac{\bar{\gamma} - \bar{\alpha}}{2}} P_{n+\mu}^{-\mu}(\lambda^{\frac{1}{2}} x), & |x| < \lambda^{-\frac{1}{2}} \\ 0, & |x| > \lambda^{-\frac{1}{2}}, \quad n = 0, 1, 2, 3, \dots, \end{cases} \quad (77)$$

where  $\mu = \omega_0/\lambda\hbar$  and  $N_n$  is the normalization constant. We observe that the discrete solutions of both the systems (vide (72) and (77)) differ in their independent variables which implies that the solutions of the Higgs oscillator can be related with that of the Mathews-Lakshmanan oscillator by an appropriate transformation. The corresponding energy spectrum of the later system is

$$E_n = \left(n + \frac{1}{2}\right) \hbar \sqrt{\omega_0^2 + \hbar^2 \lambda^2 (4\bar{\alpha} - \bar{\gamma} + (\bar{\gamma} - \bar{\alpha})^2)} + (n^2 + n - \bar{\alpha} - \bar{\gamma}) \frac{\hbar^2 \lambda}{2}. \quad (78)$$

The bound-state energy levels of the Higgs oscillator are found to be a function of  $k$  similar to the Mathews-Lakshmanan oscillator (1) in which the energy levels are also quadratic in  $n$  and are linearly dependent on the parameter  $\lambda$ .

**(ii) Negative values of  $k$**

When  $k < 0$ , that is  $k = -|k|$ , the spatial region is divided into the following two regions:

$$\text{Region I :} \quad -\frac{1}{\sqrt{|k|}} \leq x \leq \frac{1}{\sqrt{|k|}} \quad (79)$$

$$\text{Region II :} \quad |x| > \frac{1}{\sqrt{|k|}}. \quad (80)$$

We analyze the solvability of (49) in accordance with the two regions. Eq. (49) then becomes

$$\psi'' - \frac{4|k|x}{(1 - |k|x^2)}\psi' + \left( \frac{-2\eta_1|k| + \frac{\omega_0^2}{\hbar^2|k|}}{(1 - |k|x^2)} + \frac{\frac{2E}{\hbar^2} - 4\eta_2|k| - \frac{\omega_0^2}{\hbar^2|k|}}{(1 - |k|x^2)^2} \right) \psi = 0, \quad (81)$$

where  $\eta_1$  and  $\eta_2$  are defined in (50) and (51).

**(a) Region I: Bound States**

On using the transformation,

$$z = |k| x^2, \quad \psi(z) = (1 - z)^d \phi(z), \quad (82)$$

we can express the equation (49) as

$$z(1 - z)\phi''(z) - \left[ \frac{1}{2} - \left( 2d + \frac{5}{2} \right) z \right] \phi'(z) - \left( d - \frac{\tilde{\mu}}{2} + \frac{3}{4} \right) \left( d + \frac{\tilde{\mu}}{2} + \frac{3}{4} \right) \phi(z) = 0 \quad (83)$$

with

$$\frac{E}{2\hbar^2|k|} - \frac{\mu^2}{4} - \eta_2 + d(d + 1) = 0, \quad (84)$$

$$\mu = \frac{\omega_0}{\hbar k}, \quad \text{and} \quad (85)$$

$$\tilde{\mu} = \sqrt{\mu^2 - 2\eta_1 + \frac{9}{4}}. \quad (86)$$

Equation (83) is of the form of the hypergeometric differential equation (58) and hence the solution,  $\psi(z)$  (vide (59)), in terms of  $x$  can be written as,

$$\begin{aligned} \psi(x) = (1 - |k|x^2)^d & \left[ C_1 {}_2F_1 \left( d + \frac{3}{4} - \frac{\tilde{\mu}}{2}, d + \frac{3}{4} + \frac{\tilde{\mu}}{2}; \frac{1}{2}; |k|x^2 \right) \right. \\ & \left. + C_2 \sqrt{|k|x} {}_2F_1 \left( d - \frac{1}{4} - \frac{\tilde{\mu}}{2}, d - \frac{1}{4} + \frac{\tilde{\mu}}{2}; \frac{3}{2}; |k|x^2 \right) \right], \end{aligned} \quad (87)$$



where  $C_1$  and  $C_2$  are parameters.

We consider positive values of  $d$  to obtain bounded solutions. And the hypergeometric function of (87) is reduced to polynomial form so as to normalize the solution as

$$\psi(x) = N 2^{d-1} \frac{(1 - \sqrt{|k|x})^d}{(1 + \sqrt{|k|x})^{d+1}} {}_2F_1 \left( \tilde{\mu} - \frac{1}{2}, \tilde{\mu} + \frac{1}{2}; 2d + 2; \frac{1 - \sqrt{|k|x}}{2} \right), \quad (88)$$

where we have used the identities subsequently,

$$\begin{aligned} {}_2F_1 \left( a, b; \frac{a+b+1}{2}; y \right) &= \frac{\sqrt{\pi} \Gamma(\frac{a+b+1}{2})}{\Gamma(\frac{a+1}{2}) \Gamma(\frac{b+1}{2})} {}_2F_1 \left( \frac{a}{2}, \frac{b}{2}; \frac{1}{2}; (2y-1)^2 \right) \\ &\quad + (2y-1) \frac{2\sqrt{\pi} \Gamma(\frac{a+b+1}{2})}{\Gamma(\frac{a}{2}) \Gamma(\frac{b}{2})} {}_2F_1 \left( \frac{a+1}{2}, \frac{b+1}{2}; \frac{3}{2}; (2y-1)^2 \right), \quad (89) \\ {}_2F_1(a, b; c; y) &= (1-y)^{c-a-b} {}_2F_1(c-a, c-b; c; y). \quad (90) \end{aligned}$$

Equation (88) is reduced to be polynomial when  $\tilde{\mu} + \frac{1}{2} - n = 2d + 2$ ,  $n = 0, 1, 2, \dots$  is integer. Hence, the value of  $d$  becomes  $d = \frac{1}{2} \left( -n + \tilde{\mu} - \frac{3}{2} \right)$  to get bounded and normalizable eigenfunction. Now we use another identity,

$$P_\nu^m(y) = \frac{1}{\Gamma(1-m)} \frac{(1+y)^{m/2}}{(1-y)^{m/2}} {}_2F_1 \left( -\nu, \nu + 1; 1 - m; \frac{1-y}{2} \right) \quad (91)$$

which transforms the solution (88) to

$$\psi_n(x) = N_n \frac{\Gamma(\frac{1}{2} - n + \tilde{\mu})}{\sqrt{(1 - |k|x)^2}} P_{\tilde{\mu} - \frac{1}{2}}^{n - \tilde{\mu} + \frac{1}{2}}(\sqrt{|k|x}), \quad (92)$$

where  $P_\nu^m(y)$  is known to be the associated Legendre polynomial with  $m$  and  $\nu$  as parameters. Here,  $n = 0, 1, 2, \dots, N$  and the upper limit  $N < 2\tilde{\mu} - 1$  and the normalization constant  $N_n$  is evaluated to be

$$N_n = \left( \frac{\sqrt{|k|} (n - \tilde{\mu} + \frac{1}{2}) \Gamma(2\tilde{\mu} - n)}{\Gamma(n+1) 2^{\frac{n}{4} - \frac{\tilde{\mu}}{4} + \frac{1}{8}}} \right)^{1/2}, \quad (93)$$

since

$$\int_{-1}^1 \frac{P_l^m(y) P_l^n(y)}{\sqrt{1-y^2}} dy = \begin{cases} 0, & m \neq n, \\ \frac{(l+m)!}{m(l-m)!}, & m = n \neq 0, \\ \infty, & m = n = 0. \end{cases} \quad (94)$$

Energy eigenvalues can be evaluated from (84) as

$$E_n = \left( n + \frac{1}{2} \right) \hbar \omega_0 \sqrt{1 + \frac{|k|^2 \hbar^2}{\omega_0^2} \left( \frac{9}{4} - 2\eta_1 \right)} - \left( n^2 + n + 2\bar{\alpha} + 2\bar{\gamma} + \frac{3}{2} \right) \frac{\hbar^2 |k|}{2}, \quad n = 0, 1, 2, 3, \dots, N. \quad (95)$$

The energy eigenvalues  $E_n$  are discrete and finite in number. And above the upper limit of  $n$ , that is  $N$ , the energy eigenvalue is continuous.

**(b) Region II: Continuous energy states**

We consider the region  $|x| > \frac{1}{\sqrt{|k|}}$ . Eq. (83) is in the form of hypergeometric equation (58) which admits solution near  $z = \infty$  as

$$w(z) = C_1(1-z)^{-a} {}_2F_1\left(a, c-b; a-b+1; \frac{1}{1-z}\right) + C_2(1-z)^{-b} {}_2F_1\left(b, c-a; b-a+1; \frac{1}{1-z}\right). \quad (96)$$

As  $a = d + \frac{\tilde{\mu}}{2} + \frac{3}{4}$ ,  $b = d - \frac{\tilde{\mu}}{2} + \frac{3}{4}$  and  $c = \frac{1}{2}$  in (83), we can express the solution of (81) near infinity, that is  $x = \infty$ , as

$$\begin{aligned} \psi(x) = (1 - |k|x^2)^d \left[ C_1(1 - |k|x^2)^{-d - \frac{\tilde{\mu}}{2} - \frac{3}{4}} {}_2F_1\left(d + \frac{\tilde{\mu}}{2} + \frac{3}{4}, -d - \frac{1}{4} + \frac{\tilde{\mu}}{2}; 1 + \tilde{\mu}; \frac{1}{1 - |k|x^2}\right) \right. \\ \left. + C_2(1 - |k|x^2)^{-d + \frac{\tilde{\mu}}{2} - \frac{3}{4}} {}_2F_1\left(d - \frac{\tilde{\mu}}{2} + \frac{3}{4}, -d - \frac{\tilde{\mu}}{2} + \frac{3}{4}; 1 - \tilde{\mu}; \frac{1}{1 - |k|x^2}\right) \right]. \end{aligned} \quad (97)$$

Now we use the relation of associated Legendre polynomial  $P_\nu^m(y)$  with hypergeometric function,

$$\begin{aligned} P_\nu^m(y) = \frac{\Gamma(-\nu - \frac{1}{2})}{2^{\nu+1} \sqrt{\pi} \Gamma(-\nu + \mu)} (y^2 - 1)^{-\frac{1+\nu}{2}} {}_2F_1\left(\frac{\nu + m + 1}{2}, \frac{\nu - m + 1}{2}; \nu + \frac{3}{2}; \frac{1}{1 - y^2}\right) \\ + \frac{2^\nu \Gamma(\nu + \frac{1}{2})}{\sqrt{\pi} \Gamma(\nu + m + 1)} (y^2 - 1)^{\frac{\nu}{2}} {}_2F_1\left(\frac{-\nu - m}{2}, \frac{-\nu + m}{2}; -\nu + \frac{1}{2}; \frac{1}{1 - y^2}\right), \quad |1 - y^2| > 1 \end{aligned} \quad (98)$$

and express the solution (97) in terms of associated Legendre polynomial in which we have chosen the arbitrary constants  $C_1 = \frac{\Gamma(-\nu - \frac{1}{2})}{2^{\nu+1} \sqrt{\pi} \Gamma(-\nu + \mu)}$ ,  $C_2 = \frac{2^\nu \Gamma(\nu + \frac{1}{2})}{\sqrt{\pi} \Gamma(\nu + m + 1)}$  and  $\nu = \tilde{\mu} - \frac{1}{2}$  and  $-m = 2d + 1$  to be

$$\psi(x) = N (1 - |k|x^2)^{-1/2} P_{\mu - \frac{1}{2}}^{-2d-1}(\sqrt{|k|x}), \quad |x| > \frac{1}{\sqrt{|k|}}. \quad (99)$$

Since the Hamiltonian is Hermitian, we consider  $2d = -1 + i\rho$ , where  $\rho$  is a real parameter and the solution (99) turns out to be

$$\psi_\rho(x) = (1 - |k|x^2)^{-1/2} P_{\mu - \frac{1}{2}}^{-i\rho}(\sqrt{|k|x}) \Theta(\pm \sqrt{|k|x} - 1), \quad (100)$$

with energy eigenvalues,

$$E_\rho = \left( \frac{\rho^2 + \mu^2 + 1}{2} + 2\eta_2 \right) \hbar^2 |k|, \quad (101)$$

where  $\Theta(\pm \sqrt{|k|x} - 1)$  is known as Heaviside step function, which is defined as  $\Theta(y) = 1$  for  $y > 0$  and  $\Theta(y) = 0$  for  $y < 0$ .

## B. Quantum solvability of the nonlinear system $H_2$

Let us next consider the Hamiltonian,

$$H_2 = (1+kx^2)^2 \frac{p^2}{2} + \frac{\omega_0^2 x^2}{2(1+kx^2)^2}. \quad (44)$$

The generalized time-independent Schrödinger equation for the Hamiltonian (44) becomes (vide Eq. (48))

$$\psi'' + \frac{4kx}{(1+kx^2)}\psi' + \left( \frac{2\eta_1 k}{(1+kx^2)} + \frac{\frac{2E}{\hbar^2} + 4\eta_2 k}{(1+kx^2)^2} - \frac{\omega_0^2 x^2}{\hbar^2 (1+kx^2)^4} \right) \psi = 0, \quad (102)$$

where

$$\eta_1 = 5(\bar{\alpha} + \bar{\gamma}) - 8 \left( \bar{\alpha}\bar{\gamma} + \bar{\alpha} + \bar{\gamma} + \frac{1}{4}(\bar{\gamma} - \bar{\alpha})^2 \right), \quad (50)$$

$$\eta_2 = -3(\bar{\alpha} + \bar{\gamma}) + 4 \left( \bar{\alpha}\bar{\gamma} + \bar{\alpha} + \bar{\gamma} + \frac{1}{4}(\bar{\gamma} - \bar{\alpha})^2 \right). \quad (51)$$

The asymptotic requirement  $\psi(x) \rightarrow 0$  as  $x \rightarrow \pm\infty$  and the presence of the inverse power of 4 term inside the bracket in (102) lead to the ansatz,

$$\psi(x) = \exp\left(\frac{-\mu k x^2}{2(1+kx^2)}\right) (1+kx^2)^d (kx^2)^l S(x), \quad (103)$$

which reduces (102) into,

$$S''(x) + \left( \frac{4l}{x} + \frac{4k(d+1)x}{1+kx^2} - \frac{2\mu kx}{(1+kx^2)^2} \right) S'(x) + \left( \frac{2l(2l-1)}{x^2} + \frac{4\tilde{\eta}k}{1+kx^2} + \frac{4\xi k}{(1+kx^2)^2} + \frac{4\mu dk}{(1+kx^2)^3} \right) S(x) = 0, \quad (104)$$

where

$$\tilde{\eta} = \frac{\eta_1}{2} + d \left( d + \frac{3}{2} \right) + 2l(d+1), \quad (105)$$

$$\xi = \frac{E}{2\hbar^2 k^2} + \eta_2 - \frac{\mu}{4} - \mu(l+d) - d(d+1). \quad (106)$$

with  $\mu = \frac{\omega_0}{\hbar k}$ . On using the transformation  $z = \frac{k x^2}{1+kx^2}$  in Eq.(104), we can reexpress it as

$$z(1-z)^2 S''(z) + \left[ 2l + \frac{1}{2} + \left( 2d - \mu - 2l - \frac{1}{2} \right) z + (-2d + 2\mu) z^2 - \mu z^3 \right] S'(z) + [\xi + \tilde{\eta} + \mu d - (\xi + 2\mu d)z + \mu d z^2] S(z) = 0, \quad |z| < 1. \quad (107)$$

This is of the form of confluent Heun equation which can be solved using Bethe ansatz method [19, 20].

### 1. Bethe-Ansatz method

To start with, we first give a brief description of the method which is a quasi-exact treatment and then use the results to solve equation (107), [19, 20]. Consider the differential equation of the form,

$$\sum_{j=0}^4 a_j z^j S''(z) + \sum_{j=0}^3 b_j z^j S'(z) + \sum_{j=0}^2 c_j z^j S(z) = 0, \quad (108)$$

where  $a_0, a_1, a_2, a_3, a_4, b_0, b_1, b_2, b_3, c_0, c_1$  and  $c_2$  are parameters.

Eq. (108) has a  $n$ -degree polynomial solution,

$$S(z) = \prod_{i=1}^n (z - z_i), \quad S(z) = 1 \quad \text{for } n = 0, \quad (109)$$

with the distinct roots  $z_1, z_2, \dots, z_n$ , satisfying the Bethe-ansatz equations,

$$\sum_{j \neq i}^n \frac{2}{z_i - z_j} = -\frac{b_3 z_i^3 + b_2 z_i^2 + b_1 z_i + b_0}{a_4 z_i^4 + a_3 z_i^3 + a_2 z_i^2 + a_1 z_i + a_0}, \quad (110)$$

provided the following restrictions on the parameters hold:

$$c_2 = -n(n-1)a_4 - nb_3, \quad (111)$$

$$c_1 = -nb_2 - n(n-1)a_3 - (2(n-1)a_4 + b_3) \sum_{i=1}^n z_i, \quad (112)$$

$$-c_0 = (2(n-1)a_4 + b_3) \sum_{i=1}^n z_i^2 + 2a_4 \sum_{i < k}^n z_i z_k + (2(n-1)a_3 + b_2) \sum_{i=1}^n z_i + n b_1. \quad (113)$$

On comparing Eq. (107) with (108) for the values of  $l = 0$  and  $l = \frac{1}{2}$ , we have  $a_4 = a_0 = 0, a_1 = 1, a_2 = -2, a_3 = 1$  and  $b_0 = 2l + \frac{1}{2}, b_1 = 2d - \mu - 2l - \frac{1}{2}, b_2 = 2\mu - 2d, b_3 = -\mu$  and  $c_0 = \xi + \tilde{\eta} + \mu d, c_1 = -\xi - 2\mu d, c_2 = \mu d$ . The relations (111), (112) and (113) imply that

$$d = n, \quad (114)$$

$$\xi = -\mu \sum_{i=1}^n z_i - n(n+1), \quad (115)$$

$$\tilde{\eta} = \mu \sum_{i=1}^n z_i^2 + (2 - \mu) \sum_{i=1}^n z_i - n \left( n - 2l - \frac{3}{2} \right) \quad (116)$$

with

$$\sum_{j \neq i}^n \frac{2}{z_i - z_j} = -\frac{\mu z_i^3 + \left( -\mu + 2n - 2l - \frac{1}{2} \right) z_i + (2\mu - 2n) z_i^2 + 2l + \frac{1}{2}}{z_i (1 - z_i)^2}. \quad (117)$$

Hence we can write down the eigenfunctions,  $\psi_n(x)$  by using (109) and then the resultant expression in (103) with  $z = \frac{kx^2}{1+kx^2}$  as

$$\psi_n^{(l)}(x) = N_n^{(l)} \exp\left(-\frac{\omega_0 x^2}{2\hbar(1+kx^2)}\right) (1+kx^2)^n (kx^2)^l \prod_{i=0}^n \left(\frac{kx^2}{1+kx^2} - z_i\right) \quad (118)$$

and the energy eigenvalues as

$$E_n^{(l)} = \left(2n + 2l + \frac{1}{2}\right) \hbar\omega_0 + \left(-2\mu \sum_{i=1}^n z_i + \bar{\gamma} + \bar{\alpha} + \eta_1\right) \hbar^2 k \quad (119)$$

with

$$\eta_1 = 2\mu \sum_i^n z_i^2 + 2(2-\mu) \sum_{i=1}^n z_i - 4n^2 - 4l. \quad (120)$$

Hence while obtaining the eigenfunctions (118) and eigenvalues (119), the term  $\eta_1$  in (120) containing ordering parameters is related with the quantum number  $n$  given in (120) in which the ordering parameters,  $\alpha_i + \beta_i + \gamma_i = -1$ ,  $i = 1, 2, 3, \dots, N$ , are considered to be arbitrary.

To describe the quasi-polynomial form of the solutions (118), we explicitly obtain the ground state. When  $l = 0$ , the ground state solution,  $\psi_0^{(0)}$ , can be expressed as

$$\psi_0^{(0)}(x) = \exp\left(-\frac{\omega_0 x^2}{2\hbar(1+kx^2)}\right), \quad (121)$$

$$E_0^{(0)} = \frac{\hbar\omega_0}{2} + (\bar{\alpha} + \bar{\gamma}) \hbar^2 k. \quad (122)$$

This solution (121) is not bounded when  $x \rightarrow \pm\infty$  for  $k > 0$ , whereas for  $k < 0$ , that is  $k = -|k|$ , the ground state solution,

$$\psi_0^{(0)}(x) = N_0^0 \exp\left(-\frac{\omega_0 x^2}{2\hbar(1-|k|x^2)}\right), \quad (123)$$

which is bounded within  $\frac{-1}{\sqrt{|k|}} < x < \frac{1}{\sqrt{|k|}}$ . To obtain the normalized solution, we consider the normalization condition,

$$1 = N_0^{(0)^2} \int_{-\frac{1}{\sqrt{|k|}}}^{\frac{1}{\sqrt{|k|}}} \exp\left(-\frac{\omega_0 x^2}{\hbar(1-|k|x^2)}\right) dx. \quad (124)$$

The integral (124) can be integrated with the factor  $m^{3/4}$  which will be produced when we use non-Hermitian ordered Hamiltonian in the following.

## 2. Non-Hermitian Hamiltonian

The general ordered one-dimensional non-Hermitian Hamiltonian (45) is of the expanded form as

$$\hat{H} = -\frac{\hbar^2}{2m} \left[ \frac{d^2}{dx^2} + (\bar{\gamma} - \bar{\alpha} - 1) \frac{m'}{m} + \left( \bar{\gamma} \frac{m''}{m} - (\bar{\alpha}\bar{\gamma} + 2\bar{\gamma}) \frac{m'^2}{m^2} \right) \right] + V(x). \quad (125)$$

The associated time-independent one dimensional generalized Schrödinger equation is

$$\Phi'' + \frac{4kx}{(1+kx^2)}(1+\bar{\alpha}-\bar{\gamma})\Phi' + \left( \frac{4\sigma_1 k}{(1+kx^2)} + \frac{\frac{2E}{\hbar^2} + 4\sigma_2 k}{(1+kx^2)^2} - \frac{\omega_0^2 x^2}{\hbar^2 (1+kx^2)^4} \right) \Phi = 0, \quad (126)$$

where

$$\sigma_1 = -4\bar{\alpha}\bar{\gamma} - 3\bar{\gamma}, \quad (127)$$

$$\sigma_2 = 4\bar{\alpha}\bar{\gamma} + 2\bar{\gamma}. \quad (128)$$

Here we use the same form of the ansatz as in the case of Hermitian ordering (103),

$$\Phi(x) = \exp\left(\frac{-\mu k x^2}{2(1+kx^2)}\right) (1+kx^2)^d (kx^2)^l S(x), \quad (129)$$

which reduces Eq. (126) into

$$S''(x) + \left( \frac{4l}{x} + \frac{4k(d+\bar{\alpha}-\bar{\gamma}+1)x}{1+kx^2} - \frac{2\mu kx}{(1+kx^2)^2} \right) S'(x) + \left( \frac{2l(2l-1)}{x^2} + \frac{4\sigma k}{1+kx^2} + \frac{4\epsilon k}{(1+kx^2)^2} + \frac{4\mu(d+\bar{\alpha}-\bar{\gamma})k}{(1+kx^2)^3} \right) S(x) = 0, \quad (130)$$

where

$$\sigma = \sigma_1 + d \left( d + \frac{3}{2} + 2l + 2\bar{\alpha} - 2\bar{\gamma} \right) + 2l(1+\bar{\alpha}-\bar{\gamma}), \quad (131)$$

$$\epsilon = \frac{E}{2\hbar^2 k^2} + \sigma_2 - \frac{\mu}{4}(1+4l+4d+4\bar{\alpha}-4\bar{\gamma}) - d(d+1+2\bar{\alpha}-2\bar{\gamma}). \quad (132)$$

with  $\mu = \frac{\omega_0}{\hbar k}$ . The transformation  $\sqrt{k}x = \frac{\sqrt{z}}{\sqrt{1-z}}$  reduces equation (130) into

$$z(1-z)^2 S''(z) + \left[ 2l + \frac{1}{2} + \left( 2d + 2\bar{\alpha} - 2\bar{\gamma} - \mu - 2l - \frac{1}{2} \right) z + (-2d - 2\bar{\alpha} + 2\bar{\gamma} + 2\mu) z^2 - \mu z^3 \right] S'(z) + [\epsilon + \sigma + \mu(d+\bar{\alpha}-\bar{\gamma}) - (\epsilon + 2\mu(d+\bar{\alpha}-\bar{\gamma}))z + \mu(d+\bar{\alpha}-\bar{\gamma})z^2] S(z) = 0, \quad (133)$$

$|z| < 1,$

On comparing Eq. (133) with (108) with  $l = 0$  and  $l = \frac{1}{2}$ , we have  $a_0 = a_4 = 0, a_1 = 1, a_2 = -2, a_3 = 1$  and  $b_0 = 2l + \frac{1}{2}, b_1 = 2d + 2\bar{\alpha} - 2\bar{\gamma} - \mu - 2l - \frac{1}{2}, b_2 = 2\mu - 2d - 2\bar{\alpha} + 2\bar{\gamma}, b_3 = -\mu$  and  $c_0 = \epsilon + \sigma + \mu(d + \bar{\alpha} - \bar{\gamma}), c_1 = -\epsilon - 2\mu(d + \bar{\alpha} - \bar{\gamma}), c_2 = \mu(d + \bar{\alpha} - \bar{\gamma})$ . The relations (111), (112) and (113) ensure that

$$d = n + \bar{\gamma} - \bar{\alpha}, \quad (134)$$

$$\epsilon = -\mu \sum_{i=1}^n z_i - n(n+1), \quad (135)$$

$$\sigma = \mu \sum_i z_i^2 + (2 - \mu) \sum_{i=1}^n z_i - n \left( n - 2l - \frac{3}{2} \right) \quad (136)$$

with

$$\sum_{j \neq i}^n \frac{2}{z_i - z_j} = \frac{\mu z_i^2 + (-\mu + 2n) z_i + 2l + \frac{1}{2}}{z_i(z_i - 1)}. \quad (137)$$

Hence we can write down the eigenfunctions,  $\Phi_n(x)$ , by using (109) in  $S(z)$  and then the resultant expression in (129) with  $z = \frac{kx^2}{1 + kx^2}$  as

$$\Phi_n^{(l)}(x) = N_n^{(l)} \exp \left( -\frac{\omega_0 x^2}{2\hbar(1 + kx^2)} \right) (1 + kx^2)^{n + \bar{\gamma} - \bar{\alpha}} (kx^2)^l \prod_{i=0}^n \left( \frac{kx^2}{1 + kx^2} - z_i \right) \quad (138)$$

and the energy eigenvalues are

$$E_n^{(l)} = \left( 2n + 2l + \frac{1}{2} \right) \hbar \omega_0 + \left( \sigma_1 + \bar{\gamma} - (\bar{\gamma} - \bar{\alpha})(\bar{\gamma} - \bar{\alpha} - 1) - \mu \sum_{i=1}^n z_i \right) 2\hbar^2 k \quad (139)$$

with

$$\sigma_1 = \mu \sum_i z_i^2 + (2 - \mu) \sum_{i=1}^n z_i - 2n^2 - 2l + (\bar{\gamma} - \bar{\alpha}) \left( \bar{\gamma} - \bar{\alpha} - \frac{3}{2} \right). \quad (140)$$

Here it is also noted that the term  $\sigma_1$  containing ordering parameters is related with the quantum number  $n$  given in (140) and hence the ordering parameters,  $\alpha_i + \beta_i + \gamma_i = -1, i = 1, 2, 3, \dots, N$ , are considered to be arbitrary.

**(a) Ground state:  $n = 0$  and  $l = 0$**

When  $l = 0$ , the ground state solution,  $\Phi_0^{(0)}$ , can be explicitly expressed as

$$\Phi_0^{(0)}(x) = N_0^{(0)} \exp \left( -\frac{\omega_0 x^2}{2\hbar(1 + kx^2)} \right) (1 + kx^2)^{\bar{\gamma} - \bar{\alpha}} \quad (141)$$

and the energy eigenvalues are

$$E_0^{(0)} = \frac{\hbar \omega_0}{2} + (\bar{\alpha} + \bar{\gamma}) \hbar^2 k \quad (142)$$

with a restriction on ordering parameters through (140) as

$$4\bar{\alpha}\bar{\gamma} = -(\bar{\gamma} - \bar{\alpha})^2 - \frac{3}{2}(\bar{\gamma} + \bar{\alpha}). \quad (143)$$

**(i) Boundness:**

For  $k > 0$ ,  $\psi_0^{(0)}(x) = 0$  at  $x = \pm\infty$  if  $\bar{\gamma} - \bar{\alpha} < 0$ , whereas for  $k < 0$ ,  $\psi_0^{(0)}(x) = 0$  at  $x = \pm \frac{1}{\sqrt{|k|}}$  for all values of  $\bar{\gamma} - \bar{\alpha}$ .

**(ii) Normalizability:**

We have already seen that non-Hermitian ordered Hamiltonian  $\hat{H}_{non}$  is related with the Hermitian ordered Hamiltonian  $\hat{H}_{her}$  with respect to  $m^\eta$  as given in (46). And the eigenfunction of  $\hat{H}_{her}$ , say  $\psi$ , is related with that of the non-Hermitian Hamiltonian  $\hat{H}_{non}$ , say  $\Phi$ , through the relation  $\Phi = m^{-\eta}\psi$ . Hence we can call non-hermitian ordered Hamiltonian  $\hat{H}_{non}$  as quasi-Hermitian. In general, the normalizability condition for the quasi-Hermitian Hamiltonian  $\hat{H}_{non}$  is generalized to be  $\langle m^\eta \Phi | m^\eta \Phi \rangle = 1$ . But in our study we observe that the presence of  $(1 + kx^2)^{\bar{\gamma} - \bar{\alpha}} (= m^{2\eta})$  is responsible for the solution (138) to be bounded as well as normalizable. Hence we discuss about the normalizability of the ground state solution for both the cases. For  $k < 0$ , the normalization condition becomes

$$\begin{aligned} 1 &= \int \Phi^{(0)*} \Phi^{(0)} dx \\ &= \int_{-\frac{1}{\sqrt{|k|}}}^{\frac{1}{\sqrt{|k|}}} \exp\left(-\frac{\omega_0 x^2}{\hbar(1 - |k|x^2)}\right) (1 - |k|x^2)^{2\bar{\gamma} - 2\bar{\alpha}} dx. \end{aligned} \quad (144)$$

On using the transformation  $z = \frac{\sqrt{|k|x}}{\sqrt{1 - |k|x^2}}$ , Eq. (144) becomes

$$1 = 2 \frac{N_0^{(0)2}}{\sqrt{|k|}} \int_0^\infty \exp\left(-\frac{\omega_0 z^2}{\hbar|k|}\right) (1 + z^2)^{2\bar{\alpha} - 2\bar{\gamma} - \frac{3}{2}} dz. \quad (145)$$

By considering  $\bar{\alpha} - \bar{\gamma} = \frac{3}{4}$ , the integral (145) leads to

$$N_0^{(0)} = \left(\frac{\omega_0}{\hbar \pi}\right)^{1/4}. \quad (146)$$

Similarly for  $k > 0$  with  $\bar{\alpha} - \bar{\gamma} = \frac{3}{4}$ , by using the transformation  $z = \frac{\sqrt{k}x}{\sqrt{1 + kx^2}}$ , we can evaluate the normalization constant,

$$N_0^{(0)} = \left(\frac{4 \omega_0}{\hbar \pi}\right)^{1/4} \frac{1}{\sqrt{\text{erf}\left(\sqrt{\frac{\omega_0}{\hbar k}}\right)}}, \quad (147)$$



where  $erf(a)$  is the error function. Here, we may choose  $\bar{\alpha} - \bar{\gamma} = s + \frac{3}{4}$ , where  $s$  is an integer. For the ground state, we have chosen  $s = 0$ . Essentially this choice leads to one more constraint on the ordering parameters, that is  $\bar{\alpha} - \bar{\gamma} = s + \frac{3}{4}$ ,  $s = 0, 1, 2, 3, \dots$ , besides  $\bar{\alpha} + \bar{\beta} + \bar{\gamma} = -1$ . Hence the solutions (141) and (142) now consist of only one arbitrary ordering parameter.

**(b) Ground state:  $n = 0$  and  $l = 1/2$**

When  $l = 1/2$ , the ground state solution,  $\Phi_0^{(1/2)}$ , can be explicitly expressed as

$$\Phi_0^{(1/2)}(x) = N_0^{(1/2)} \exp\left(-\frac{\omega_0 x^2}{2\hbar(1+kx^2)}\right)(1+kx^2)^{\bar{\gamma}-\bar{\alpha}}\sqrt{k} x, \quad (148)$$

$$E_0^{(1/2)} = \frac{3}{2}\hbar\omega_0 + (\bar{\alpha} + \bar{\gamma} - 2)\hbar^2k \quad (149)$$

with

$$4\bar{\alpha}\bar{\gamma} = -(\bar{\gamma} - \bar{\alpha})^2 - \frac{3}{2}(\bar{\gamma} + \bar{\alpha}) + 1. \quad (150)$$

As in the earlier case, we consider  $\bar{\gamma} - \bar{\alpha} = -\frac{5}{4}$  to make  $\Phi_0^{(1/2)}(x) = 0$  for both the cases  $k < 0$  and  $k > 0$  at their corresponding boundaries. The normalization constants can be obtained as

$$N_0^{(1/2)} = \left[-\left(\frac{\omega_0}{\hbar}\right)^{3/2} \frac{2}{|k|\sqrt{\pi}}\right]^{1/2}, \quad k < 0, \quad (151)$$

where  $\sqrt{-|k|}$  gets canceled by the same term in (148) and

$$N_0^{(1/2)} = \left(\frac{\mu^{3/2} \sqrt{k}}{\left(\frac{\sqrt{\pi}}{2} erf(\sqrt{\mu}) - \sqrt{\mu} e^{-\mu}\right)}\right)^{1/2}, \quad k > 0. \quad (152)$$

**(c) Bound state  $n = 1$  and  $l = 0$**

Now we evaluate the excited state  $n = 1$  and  $l = 0$ ,

$$\Phi_1^{(0)}(x) = N_1^{(0)} \exp\left(-\frac{\omega_0 x^2}{2\hbar(1+kx^2)}\right)(1+kx^2)^{1+\bar{\gamma}-\bar{\alpha}}\left(\frac{kx^2}{1+kx^2} - z_1\right) \quad (153)$$

and the energy eigenvalues,

$$E_1^{(0)} = \frac{5}{2}\hbar\omega_0 - (\mu z_1 - \sigma_1 - \bar{\gamma} + (\bar{\gamma} - \bar{\alpha})(\bar{\gamma} - \bar{\alpha} - 1))2\hbar^2k. \quad (154)$$

with

$$\sigma_1 = \mu z_1^2 + (2 - \mu)z_1 - 2 + (\bar{\gamma} - \bar{\alpha})\left(\bar{\gamma} - \bar{\alpha} - \frac{3}{2}\right). \quad (155)$$

And  $z_1$  can be evaluated with  $l = 0$  through the equation (137),

$$\mu z_1^2 + (2 - \mu)z_1 + \frac{3}{2} = 0, \quad (156)$$

whose roots are

$$z_1 = \left( \frac{1}{2} - \frac{1}{\mu} \right) \pm \frac{1}{2\mu} \sqrt{\mu^2 - 6\mu + 4}. \quad (157)$$

The state  $\Phi_1^{(0)}$  is normalizable with  $\bar{\alpha} - \bar{\gamma} = \frac{3}{4}$  for both the cases  $k < 0$  and  $k > 0$ . The normalization constant,  $N_1^{(0)}$ , reads as

$$N_1^{(0)} = \left( \frac{\mu^{5/2} \sqrt{|k|}}{\sqrt{\pi} \left( \frac{3}{4} + \mu z_1 + \mu^2 z_1^2 \right)} \right)^{1/2}, \quad k < 0, \quad (158)$$

$$N_1^{(0)} = \left( \frac{\mu^{5/2} \sqrt{k}}{2 \frac{erf(\sqrt{\mu})}{\sqrt{\mu}} \left( \frac{3}{4} - \mu z_1 + \mu^2 z_1^2 \right) - \frac{1}{\sqrt{\pi}} \left( \frac{3}{2} - 2\mu z_1 + \mu^2 z_1^2 \right) e^{-\mu}} \right)^{1/2}, \quad k > 0. \quad (159)$$

#### (d) Bound state $n = 1$ and $l = 1/2$

Similarly we evaluate the excited state  $n = 1$  and  $l = 1/2$ ,

$$\Phi_1^{(1/2)}(x) = N_1^{(1/2)} \exp \left( -\frac{\omega_0 x^2}{2\hbar(1+kx^2)} \right) \sqrt{k} x (1+kx^2)^{1+\bar{\gamma}-\bar{\alpha}} \left( \frac{kx^2}{1+kx^2} - z_1 \right) \quad (160)$$

and the energy eigenvalues,

$$E_1^{(1/2)} = \frac{7}{2} \hbar \omega_0 - [\mu z_1 - \sigma_1 - \bar{\gamma} + (\bar{\gamma} - \bar{\alpha})(\bar{\gamma} - \bar{\alpha} - 1)] 2\hbar^2 k \quad (161)$$

with

$$\sigma_1 = \mu z_1^2 + (2 - \mu)z_1 - 3 + (\bar{\gamma} - \bar{\alpha}) \left( \bar{\gamma} - \bar{\alpha} - \frac{3}{2} \right). \quad (162)$$

And  $z_1$  can be evaluated with  $l = \frac{1}{2}$  through the equation (137),

$$\mu z_1^2 + (2 - \mu)z_1 + \frac{3}{2} = 0, \quad (163)$$

whose roots are

$$z_1 = \left( \frac{1}{2} - \frac{1}{\mu} \right) \pm \frac{1}{2\mu} \sqrt{\mu^2 - 10\mu + 4}. \quad (164)$$

The state  $\Phi_1^{(1/2)}$  is normalizable with  $\bar{\alpha} - \bar{\gamma} = \frac{9}{4}$  for both the cases  $k < 0$  and  $k > 0$ . The

normalization constant,  $N_1^{(1/2)}$ , reads as

$$N_1^{(1/2)} = \left( -\frac{\mu^{7/2} \sqrt{|k|}}{\sqrt{\pi} \left( \frac{z_1^2 \mu^3}{2} + \frac{3}{2} \mu z_1 + \frac{15}{8} \mu^2 z_1^2 \right)} \right)^{1/2}, \quad k < 0, \quad (165)$$

$$N_1^{(1/2)} = \left( \frac{-\sqrt{k}}{\left( \frac{\partial^3}{\partial \mu^3} + 2z_1 \frac{\partial^2}{\partial \mu^2} + z_1^2 \frac{\partial}{\partial \mu} \right) \frac{\text{erf}(\sqrt{\mu})}{\sqrt{\mu}}} \right)^{1/2}, \quad k > 0, \quad (166)$$

where  $\mu = \frac{\omega_0}{\hbar k}$ . Note that in the above Eqs. (141) and (153),  $l = 0$  represents even states whereas  $l = \frac{1}{2}$  in (148) and (160) represents odd states. While normalizing these states, we fixed the value of  $\bar{\alpha} - \bar{\gamma}$ . In general, the value is considered to be

$$\bar{\alpha} - \bar{\gamma} = n + l + \frac{3}{4}. \quad (167)$$

When  $n > 1$ , we get the polynomial of  $z_i$  of order  $n+1$  which can not be solved analytically. Hence, we explicitly evaluated the first two states of the system and so it is quasi exactly solvable.

## V. THREE DIMENSIONAL GENERALIZATION OF THE NONLINEAR OSCILLATORS

We extend our study to the three dimensional generalization of the two nonlinear oscillators. To start with, we consider the Lagrangian corresponding to the Higgs oscillator (see (10)),

$$L = \frac{1}{2} \left[ \frac{\dot{\mathbf{q}}^2}{(1 + k\mathbf{q}^2)} - \frac{k (\mathbf{q} \cdot \dot{\mathbf{q}})^2}{(1 + k\mathbf{q}^2)^2} - \omega_0^2 \mathbf{q}^2 \right], \quad (168)$$

where  $\omega_0$  is the potential parameter, which results in the equation of motion,

$$\frac{\ddot{q}_i}{(1 + k\mathbf{q}^2)^2} - 2k \frac{(\mathbf{q} \cdot \dot{\mathbf{q}})}{(1 + k\mathbf{q}^2)^3} \dot{q}_i + \left[ \frac{2k^2 (\mathbf{q} \cdot \dot{\mathbf{q}})^2}{(1 + k\mathbf{q}^2)^3} + \omega_0^2 \right] q_i = 0, \quad i = 1, 2, 3. \quad (169)$$

We express the equation of motion (169) in polar coordinates  $(r, \theta, \phi)$ , as  $q_1 = r \sin \theta \cos \phi$ ,  $q_2 = r \sin \theta \sin \phi$ ,  $q_3 = r \cos \theta$ . The radial part (that is for  $r$ ) and angular parts (for  $(\theta, \phi)$ ) become,

$$\frac{\ddot{r}}{(1 + kr^2)^2} - \frac{2kr}{(1 + kr^2)^3} \dot{r}^2 + \omega_0^2 r - \frac{r}{(1 + kr^2)^2} \left( \dot{\theta}^2 + \sin^2(\theta) \dot{\phi}^2 \right) = 0, \quad (170)$$

$$\ddot{\theta} + \frac{2\dot{r}\dot{\theta}}{r(1 + kr^2)} - \sin \theta \cos \theta \dot{\phi}^2 = 0, \quad (171)$$

$$\frac{r^2 \sin^2 \theta \dot{\phi}}{(1 + kr^2)} = C_1, \quad (172)$$

where  $C_1$  is constant. On substituting (172) in (171) and (170), we obtain

$$\frac{r^4 \dot{\theta}^2}{2(1+kr^2)^2} + \frac{C_1^2}{2\sin^2(\theta)} = \frac{C_2^2}{2}, \quad (173)$$

$$\frac{\ddot{r}}{(1+kr^2)^2} - \frac{2kr\dot{r}^2}{(1+kr^2)^3} + \omega_0^2 r - \frac{C_2^2}{r^3} = 0. \quad (174)$$

Equation (174) can be expressed as

$$\frac{\dot{r}^2}{(1+kr^2)^2} + \frac{C_2^2(1+kr^2)}{r^2} + \omega_0^2 r^2 = C_3, \quad (175)$$

where  $C_3$  is a third integration constant.

We first consider the radial part (175). By using the transformation,

$$z = \frac{r^2}{1+kr^2}, \quad (176)$$

Eq. (174) becomes

$$\dot{z}^2 = -az^2 + bz - c, \quad (177)$$

where  $a = 4(k^2 C_2^2 + kC_3 + \omega_0^2)$ ,  $b = 4(C_3 + 2kC_2^2)$  and  $c = 4C_2^2$ . On further using the transformation,  $z = z' - \tau$ , equation (177) becomes

$$\dot{z}' = \sqrt{a}\sqrt{\Lambda^2 - z'^2}, \quad (178)$$

where  $\tau = -\frac{C_3 + 2kC_2^2}{2(k^2 C_2^2 + kC_3 + \omega_0^2)}$  and  $\Lambda = \frac{\sqrt{C_3^2 - 4\omega_0^2 C_2^2}}{2(k^2 C_2^2 + kC_3 + \omega_0^2)}$ . On solving Eq. (178), we can get

$$z'(t) = \Lambda \sin(\Omega t + \kappa) \quad (179)$$

so that using (176) we obtain

$$r(t) = A \left[ \frac{\eta + \sin(\Omega t + \kappa)}{1 - k\eta A^2 - kA^2 \sin(\Omega t + \kappa)} \right]^{1/2}, \quad (180)$$

where  $\kappa$  is a constant and

$$\begin{aligned} \Omega &= 2\sqrt{(k^2 C_2^2 + kC_3 + \omega_0^2)}, \\ A^2 &= \Lambda = \frac{\sqrt{C_3^2 - 4\omega_0^2 C_2^2}}{2(k^2 C_2^2 + kC_3 + \omega_0^2)}, \\ \eta &= -\frac{\tau}{\Lambda} = \frac{C_3 + 2kC_2^2}{\sqrt{C_3^2 - 4\omega_0^2 C_2^2}}. \end{aligned} \quad (181)$$

When  $k > 0$ , the solution (180) is periodic for  $C_3^2 - 4\omega_0^2 C_2^2 > 0$  and  $C_3 > 0$  which is shown in the figure 3. When  $k < 0$ , the periodic solutions exist for  $C_3^2 - 4\omega_0^2 C_2^2 > 0$  and  $\Omega > 0$ .




FIG. 3: The plot of  $r(t)$  Eq. (180) for  $C_3 = 1$ ,  $C_2 = 0.2$ ,  $k = 1$  and  $\omega_0 = 1$ .

Now we consider the Lagrangian corresponding to the nonlinear oscillator (via (10)),

$$L = \frac{1}{2} \left[ \frac{\dot{\mathbf{q}}^2}{(1 + k\mathbf{q}^2)} - \frac{k(\mathbf{q} \cdot \dot{\mathbf{q}})^2}{(1 + k\mathbf{q}^2)^2} - \frac{\omega_0^2 \mathbf{q}^2}{(1 + k\mathbf{q}^2)^2} \right], \quad (182)$$

where  $\omega_0$  is the potential parameter, which results in the equation of motion,

$$\frac{\ddot{q}_i}{(1 + k\mathbf{q}^2)^2} - 2k \frac{(\mathbf{q} \cdot \dot{\mathbf{q}})}{(1 + k\mathbf{q}^2)^2} \dot{q}_i + \left[ \frac{2k^2(\mathbf{q} \cdot \dot{\mathbf{q}})^2}{(1 + k\mathbf{q}^2)^3} - \frac{k(\mathbf{q} \cdot \dot{\mathbf{q}})}{(1 + k\mathbf{q}^2)^2} + \frac{\omega_0^2(1 - k\mathbf{q}^2)}{(1 + k\mathbf{q}^2)^3} \right] q_i = 0. \quad (183)$$

We again express the equation of motion (183) in polar coordinates  $(r, \theta, \phi)$  as before. The radial part and angular parts become,

$$\ddot{r} - \frac{2kr}{(1 + kr^2)} \dot{r}^2 + \omega_0^2 r \left( \frac{1 - kr^2}{1 + kr^2} \right) - r \left( \dot{\theta}^2 + \sin^2 \theta \dot{\phi}^2 \right) = 0, \quad (184)$$

$$\ddot{\theta} + \frac{2\dot{r}\dot{\theta}}{r(1 + kr^2)} - \sin \theta \cos \theta \dot{\phi}^2 = 0, \quad (185)$$

$$\frac{r^2 \sin^2 \theta \dot{\phi}}{(1 + kr^2)} = C_1, \quad (186)$$

where  $C_1$  is constant. On substituting (186) in (185) and (184), we can get

$$\frac{r^4 \dot{\theta}^2}{2(1 + kr^2)^2} + \frac{C_1^2}{2 \sin^2 \theta} = \frac{C_2^2}{2}, \quad (187)$$

$$\frac{\ddot{r}}{(1 + kr^2)^2} - \frac{2kr \dot{r}^2}{(1 + kr^2)^3} + \omega_0^2 r \left( \frac{1 - kr^2}{1 + kr^2} \right) - \frac{C_2^2}{r^3} = 0. \quad (188)$$

On simple transformation Eq. (188) becomes that of a quadratic polynomial. It will in principle lead to elliptic functions, though complicated to express explicitly. So we do not write the form here. Therefore we will study the quantum solvability of the nonlinear system (182) in Sec. V C.

### A. Semiclassical method

The conjugate momentum for the Lagrangian (168) is

$$p_i = \frac{\partial L}{\partial \dot{q}_i} = \frac{\dot{q}_i}{(1 + k\mathbf{q}^2)} - \frac{k(\mathbf{q} \cdot \dot{\mathbf{q}})q_i}{(1 + k\mathbf{q}^2)^2} \quad (189)$$

and

$$\mathbf{p} \cdot \mathbf{p} = \frac{\dot{\mathbf{q}}^2}{(1 + k\mathbf{q}^2)^2} - \frac{k(2 - k\mathbf{q}^2)(\mathbf{q} \cdot \dot{\mathbf{q}})^2}{(1 + k\mathbf{q}^2)^4}, \quad \mathbf{p} \cdot \mathbf{q} = \frac{\mathbf{q} \cdot \dot{\mathbf{q}}}{(1 + k\mathbf{q}^2)^2} \quad (190)$$

yield the Hamiltonian corresponding to the Lagrangian (168) as

$$H = \frac{1}{2} \left[ (1 + k\mathbf{q}^2)(\mathbf{p}^2 + k(\mathbf{p} \cdot \mathbf{q})^2) + \frac{\omega_0^2 \mathbf{q}^2}{2} \right]. \quad (191)$$

In polar coordinates, the Hamiltonian can be expressed as

$$H = \frac{1}{2} \left[ \frac{\dot{r}^2}{(1 + kr^2)^2} + \frac{r^2 (\dot{\theta}^2 + \sin^2 \theta \dot{\phi}^2)}{(1 + kr^2)} + \omega_0^2 r^2 \right]. \quad (192)$$

On substituting (186) and (187), we can reduce the Hamiltonian (192) as

$$H = \frac{1}{2} \left[ \frac{\dot{r}^2}{(1 + kr^2)^2} + \frac{C_2^2(1 + kr^2)}{r^2} + \omega_0^2 r^2 \right]. \quad (193)$$

Now using the solution (180), the energy for the classical system (191) can be obtained as

$$E = \frac{1}{2} \left[ \frac{\Omega^2}{4k} - \frac{\omega_0^2}{2k} \right]. \quad (194)$$

To obtain the semiclassical energy eigenvalues, we apply the Bohr-Sommerfeld quantization rule (31) to the radial part,

$$\oint p_r dr = \left( n_r + \frac{1}{2} \right) h = I, \quad n_r = 0, 1, 2, \dots \quad (195)$$

The momentum  $p_r$  can be expressed as

$$p_r = \frac{\dot{r}}{(1 + kr^2)^2}, \quad (196)$$

along with the quantization of angular parts

$$\oint p_\theta d\theta = lh, \quad l = 0, 1, 2, \dots n. \quad (197)$$

On substituting the solution (180) in (196) and using (175) and (194) in the integral (195), we can get

$$\begin{aligned} I &= \frac{1}{\Omega} \int_0^{2\pi} \left( C_3 + \frac{\omega_0^2}{k} - \frac{C_2^2}{A^2 (\eta + \sin(u))} - \frac{\omega_0^2}{k (1 - k\eta A^2 - kA^2 \sin(u))} \right) du, \\ &= \frac{2\pi}{\Omega} \left( C_3 + \frac{\omega_0^2}{k} - \frac{C_2^2}{A^2 \sqrt{\eta^2 - 1}} - \frac{\omega_0^2}{k \sqrt{(1 - kA^2\eta)^2 - k^2 A^4}} \right). \end{aligned} \quad (198)$$

where  $u = \Omega t + \kappa$ .

On evaluating (198), we can get

$$\frac{\Omega}{4k} - C_2 - \frac{\omega_0}{k} = (2n_r + 1)\hbar, \quad (199)$$

which sets the energy value (194) as

$$E_n = \left( 2n_r + l + \frac{3}{2} \right) \hbar \omega_0 + \left( 2n_r + l + \frac{3}{2} \right)^2 \frac{\hbar^2 k}{2}. \quad (200)$$

The semiclassical energy is also of the form of a quadratic in  $n_r$ . It confirms that the quantum counterpart of the system (191) can also be exactly solved.

We do not here apply the semiclassical method to the system (182) as one requires the explicit form of the classical solution of the system, as it is quite cumbersome as pointed out earlier. However, we will carry out the Bethe ansatz method to this system in the following.

## B. Quantum solvability of the three dimensional Higgs oscillator

The components of conjugate momentum in polar coordinates  $(p_r, p_\theta, p_\phi)$  can be obtained as

$$p_r = \frac{\dot{r}}{(1 + kr^2)^2}, \quad p_\theta = \frac{r^2}{(1 + kr^2)} \dot{\theta}, \quad \text{and} \quad p_\phi = \frac{r^2 \sin^2 \theta}{(1 + kr^2)} \dot{\phi}. \quad (201)$$

On substituting (201) in (192), we can get

$$H = \frac{1}{2} \left[ (1 + kr^2)^2 p_r^2 + \frac{(1 + kr^2)}{r^2} \left( p_\theta^2 + \frac{p_\phi^2}{\sin^2 \theta} \right) + \omega_0^2 r^2 \right]. \quad (202)$$

The Hamiltonian (202) consists of position ( $r$ ) term along with momentum term  $p_r$  and obviously corresponds to a position-dependent mass (PDM) quantum system. While quantizing the Hamiltonian, it is required that the ordering between momentum and mass operators in

the kinetic energy term is taken care of appropriately. In this three dimensional case also, we consider the most general ordered form of the Hamiltonian operator,

$$\hat{H} = \frac{1}{2} \left[ \sum_{i=1}^N w_i m^{\alpha_i} \hat{p}_r m^{\beta_i} \hat{p}_r m^{\gamma_i} + \frac{(1 + kr^2)}{r^2} \left( p_\theta^2 + \frac{p_\phi^2}{\sin^2 \theta} \right) + \omega_0^2 r^2 \right], \quad (203)$$

where  $N$  is an arbitrary positive integer, and the ordering parameters should satisfy the constraint  $\alpha_i + \beta_i + \gamma_i = -1$ ,  $i = 1, 2, 3, \dots, N$  and  $w_i$ 's are real weights which are summed to be 1. The Hermitian Hamiltonian (46) for a potential  $V(r) = \frac{\omega_0^2 r^2}{2}$  with mass  $m(r) = \frac{1}{(1 + kr^2)^2}$  can be expressed as,

$$\begin{aligned} \hat{H}_{her} = & \frac{1}{2} \left( \frac{1}{m} \hat{p}_r^2 - i\hbar \frac{d}{dr} \left( \frac{1}{m} \right) \hat{p}_r \right) + \frac{\hbar^2}{2} \left[ \left( \frac{\bar{\alpha} + \bar{\gamma}}{2} \right) \frac{d^2}{dr^2} \left( \frac{1}{m} \right) + \left( \bar{\alpha}\bar{\gamma} + \frac{1}{4}(\bar{\gamma} - \bar{\alpha})^2 \right) \left( \frac{d}{dr} \left( \frac{1}{m} \right) \right)^2 m \right. \\ & \left. + \frac{1}{\sqrt{m} r^2} \left( \hat{p}_\theta^2 + \frac{\hat{p}_\phi^2}{\sin^2 \theta} \right) \right] + V(r), \end{aligned} \quad (204)$$

where  $\hat{p}_r = -i\hbar \left( \frac{\partial}{\partial r} + \frac{1}{r} \right)$ ,  $\hat{p}_\theta = -i\hbar \left( \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \right)$  and  $\hat{p}_\phi = -i\hbar \frac{\partial}{\partial \phi}$ . The generalized Schrödinger equation can be written as

$$\begin{aligned} \Psi'' + \left( \frac{2}{r} - \frac{m'}{m} \right) \Psi' + \left[ \left( \frac{\bar{\alpha} + \bar{\gamma}}{2} \right) \left( \frac{m''}{m} \right) - \left( \bar{\alpha}\bar{\gamma} + \bar{\alpha} + \bar{\gamma} + \frac{1}{4}(\bar{\gamma} - \bar{\alpha})^2 \right) \left( \frac{m'^2}{m^2} \right) \right. \\ \left. - \frac{\hat{L}^2}{r^2 \sqrt{m}} + \frac{2m}{\hbar^2} (E - V(r)) \right] \Psi = 0, \end{aligned} \quad (205)$$

where the square of the angular momentum operator  $\hat{L}^2$  is

$$\hat{L}^2 = -\hbar^2 \left( \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left( \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \right) + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \phi^2} \right), \quad (206)$$

where  $' = \frac{\partial}{\partial r}$ . Considering the wavefunction to be of the form,  $\Psi(r, \theta, \phi) = \frac{\chi(r)}{r} Y(\theta, \phi)$ , with a spherically symmetric factor  $\frac{\chi(r)}{r}$  with  $m(r) = \frac{1}{(1 + kr^2)^2}$  and a generalized angular function  $Y(\theta, \phi)$ , Eq. (205) can be separated as follows,

$$\text{Angular part:} \quad L^2 Y(\theta, \phi) = \hbar^2 l(l+1) Y(\theta, \phi), \quad (207)$$

$$\text{Radial part:} \quad \chi'' + \frac{4kr}{(1 + kr^2)} \chi' + \left[ \frac{A}{(1 + kr^2)} + \frac{B}{(1 + kr^2)^2} - \frac{l(l+1)}{r^2} \right] \chi = 0, \quad (208)$$

where  $Y_{l,m}(\theta, \phi) = \sqrt{\frac{(l-m)!}{(l+m)!}} e^{im\phi} P_l^m(\cos \theta)$  are spherical harmonics. Here  $P_l^m$  is the associated Legendre polynomial,  $l$  is the angular momentum quantum number and  $m$  is the



magnetic quantum number. For every value of  $l$ ,  $m = -l, -(l-1), \dots, 0, \dots, l-1, l$ , that is  $m$  takes  $2l+1$  values. Here

$$A = 2\eta_1 - k\mu^2 + kl(l+1), \quad (209)$$

$$B = -4\eta_2 + k\mu^2 + \frac{2E}{\hbar^2}, \quad (210)$$

$$\mu = \frac{\omega_0}{\hbar k}. \quad (211)$$

As we did in the one dimensional case, we solve the radial part (208) for both  $k > 0$  and  $k < 0$ .

### (a) Positive values of $k$

When  $k > 0$ ,  $r$  spreads over from 0 to  $\infty$ . Let us consider the transformation

$$\sqrt{k} r = \frac{\sqrt{z}}{\sqrt{1-z}}. \quad (212)$$

Then Eq. (208) gets reduced to

$$z(1-z)\chi''(z) + \frac{1}{2}\chi'(z) + \left[ \frac{B}{4k} + \frac{A}{4k(1-z)} + \frac{l(l+1)}{4z} \right] \chi(z) = 0. \quad (213)$$

On assuming  $\chi(z) = z^{\frac{l+1}{2}}(1-z)^{\frac{3}{4}+\frac{\mu}{2}}\phi(z)$ , Eq. (213) reduces to

$$z(1-z)\phi''(z) + \left[ l + \frac{3}{2} - \left( l + \mu + \frac{3}{2} \right) \right] \phi'(z) + \left[ \frac{B}{4k} - (\eta+d)(\eta+d-1) \right] \phi(z) = 0. \quad (214)$$

This is of the form of a hypergeometric equation,

$$x(1-x)w''(x) + (c - (a+b+1)x)w'(x) - abw(x) = 0, \quad (215)$$

which admits the solution at  $x = 0$  as

$$w(x) = C_1 {}_2F_1(a, b; c, x) + C_2 x^{1-c} {}_2F_1(a-c+1, b-c+1; 2-c, x), \quad |x| < 1. \quad (216)$$

Then the solution for (214) can be written as

$$\phi(z) = C_1 {}_2F_1\left(b, -b+l+\mu+\frac{3}{2}; l+\frac{3}{2}; z\right) + C_2 z^{-l-\frac{1}{2}} {}_2F_1\left(b-l-\frac{1}{2}, -b+\mu+1; \frac{1}{2}-l, z\right), \quad |z| < 1. \quad (217)$$

It reduces to polynomial form when  $b = -n_r$ ,

$$\phi(z) = C_1 {}_2F_1\left(-n_r, n_r+l+\mu+\frac{3}{2}; l+\frac{3}{2}; z\right) + C_2 z^{-l-\frac{3}{2}} {}_2F_1\left(-n_r-l-\frac{1}{2}, n_r+\mu+1; \frac{1}{2}-l, z\right), \quad |z| < 1. \quad (218)$$

where  $C_1$  and  $C_2$  are integration constants, which sets the energy eigenvalues (210) as

$$E_{n_r, l} = \left(2n_r + l + \frac{3}{2}\right) \hbar\omega_0 \sqrt{1 + \frac{1}{\mu^2} \left(\frac{9}{4} - 2\eta_1\right)} + \left[\left(2n_r + l + \frac{3}{2}\right)^2 + 2\bar{\alpha} + 2\bar{\gamma} - \frac{1}{4}\right] \frac{\hbar^2 k}{2}. \quad (219)$$

The boundary condition, that is at  $z = 1$ ,  $\phi_n(z) = 0$  and the continuity condition,  $\phi_n(z) = \text{constant} = \mathcal{N}$  at  $z = 0$  relates the constants  $C_1$  and  $C_2$ , if we choose

$$C_1 = \frac{\Gamma(-l - \frac{1}{2})}{\Gamma(n_r + 1)\Gamma(-l - n_r - \frac{1}{2})} \mathcal{N}, \quad C_2 = -\frac{\sin(n_r\pi)\Gamma(l + \frac{1}{2})\Gamma(\mu + n_r + 1)}{\pi\Gamma(n_r + l + \mu + \frac{3}{2})} \mathcal{N}. \quad (220)$$

Using now the identity,

$$\begin{aligned} P_\nu^{(a,b)}(x) &= \frac{\Gamma(-b)}{\Gamma(\nu+1)\Gamma(-b-\nu)} {}_2F_1\left(-\nu, a+b+\nu+1, b+1, \frac{x+1}{2}\right) \\ &\quad - \frac{\sin(\nu\pi)\Gamma(b)\Gamma(a+\nu+1)}{\pi\Gamma(a+b+\nu+1)} {}_2F_1\left(a+\nu+1, -b-\nu; 1-b; \frac{x+1}{2}\right) \left(\frac{x+1}{2}\right)^{-b}, \\ &\quad b \in \text{Integers}, \end{aligned} \quad (221)$$

where  $P_\nu^{(a,b)}(x)$  are Jacobi polynomials and  $\nu, a, b$  are parameters, we can reduce the solution (218) as

$$\phi_n(z) = \mathcal{N}_{n_r, l} P_{n_r}^{(\mu, l+1/2)}(2z-1). \quad (222)$$

Here  $\mathcal{N}_{n_r, l}$  is the normalization constant. Then the solution of (213) becomes

$$\chi(z) = \mathcal{N}_{n_r, l} z^{\frac{l+1}{2}} (1-z)^{\frac{3}{4}+\frac{\mu}{2}} N_n P_{n_r}^{(\mu, l+1/2)}(2z-1). \quad (223)$$

Then, in terms of  $r$ ,

$$\chi_{n, l}(r) = \mathcal{N}_{n, l} \frac{(k r^2)^{\frac{l+1}{2}}}{(1 + k r^2)^{\frac{l}{2} + \frac{\mu}{2} + \frac{5}{4}}} P_{n_r}^{(\mu, l+1/2)}\left(\frac{k r^2 - 1}{k r^2 + 1}\right). \quad (224)$$

The eigenfunction can also be expressed in terms of associated Jacobi polynomials as

$$\Psi_{n_r, l, m}(r, \theta, \phi) = \mathcal{N}_{n_r, l} \frac{1}{\sqrt{k}} \frac{(k r^2)^{\frac{l}{2}}}{(1 + k r^2)^{\frac{l}{2} + \frac{\mu}{2} + \frac{5}{4}}} P_{n_r}^{(\mu, l+1/2)}\left(\frac{k r^2 - 1}{k r^2 + 1}\right) Y_{l, m}(\theta, \phi). \quad (225)$$

The normalization constant can be evaluated as

$$1 = \oint \psi_{n, l, m}^*(r, \theta, \phi) \psi_{n, l, m}(r, \theta, \phi) r^2 \sin \theta dr d\theta d\phi,$$

so that

$$1 = \mathcal{N}_{n_r, l}^2 \int_0^\infty \int_0^{2\pi} \int_{-\pi}^\pi \frac{|\chi(r)|^2}{r^2} |Y_{l, m}|^2 r^2 \sin \theta dr d\theta d\phi. \quad (226)$$

Since the spherical harmonics are already normalized, we are left with the radial part as

$$\begin{aligned}\mathcal{N}_{n_r, l}^{-2} &= \int_{-\infty}^{\infty} |\chi(r)|^2 dr, \\ &= \int_{-\infty}^{\infty} \frac{(k r^2)^{l+1}}{(1 + k r^2)^{l+\mu+\frac{5}{2}}} P_{n_r}^{(\mu, l+1/2)} \left( \frac{k r^2 - 1}{k r^2 + 1} \right) P_{m_r}^{(\mu, l+1/2)} \left( \frac{k r^2 - 1}{k r^2 + 1} \right) dr. \quad (227)\end{aligned}$$

By using the transformation,  $\frac{\tau+1}{2} = \frac{k r^2}{(1+k r^2)}$ , we can rewrite the integration as

$$\begin{aligned}\mathcal{N}_{n_r, l}^{-2} &= \int_{-\infty}^{\infty} |\chi(r)|^2 dr, \\ &= \frac{1}{4k\sqrt{k}} \frac{1}{2^{l+\mu+1/2}} \int_{-1}^1 (1-\tau)^\mu (1+\tau)^{l+\frac{1}{2}} P_{n_r}^{(\mu, l+1/2)}(\tau) P_{m_r}^{(\mu, l+1/2)}(\tau) d\tau, \\ \mathcal{N}_{n_r, l}^{-2} &= \frac{1}{2k\sqrt{k}} \frac{\Gamma(n_r + \mu + 1)\Gamma(n_r + l + 3/2)}{n_r!(2n_r + \mu + l + 3/2)\Gamma(n_r + l + \mu + 3/2)} \delta_{n_r, m_r}, \quad (228)\end{aligned}$$

where  $\delta_{n_r, m_r}$  is the Kronecker delta function. Hence the normalization constant becomes

$$\mathcal{N}_{n_r, l} = \left( \frac{2k\sqrt{k} n_r!(2n_r + \mu + l + 3/2)\Gamma(n_r + l + \mu + 3/2)}{\Gamma(n_r + \mu + 1)\Gamma(n_r + l + 3/2)} \right)^{\frac{1}{2}}. \quad (229)$$

### (b) Negative values of $k$

When  $k < 0$ , the radius is confined within the region  $\left(0, \frac{1}{\sqrt{|k|}}\right)$ . Hence we solve the system (208) in two regions,

$$\text{Region I : } 0 \leq r < \frac{1}{\sqrt{|k|}}, \quad (230)$$

$$\text{Region II : } |r| > \frac{1}{\sqrt{|k|}}, \quad (231)$$

We use transformation

$$z = |k| r^2, \text{ and } \chi(z) = z^s (1-z)^d \phi(z), \quad (232)$$

and Eq. (208) turns out to be

$$z(1-z)\phi''(z) + \left[2s + \frac{1}{2} - \left(2s + 2d + \frac{5}{2}\right)\right] \phi'(z) - \left(s + d + \frac{3}{4} + \frac{\tilde{\mu}}{2}\right) \left(s + d + \frac{3}{4} - \frac{\tilde{\mu}}{2}\right) \phi(z) = 0, \quad (233)$$

where  $\tilde{\mu} = \sqrt{\frac{9}{4} + \mu^2 - 2\eta_1}$  and

$$s \left(s + \frac{1}{2}\right) - \frac{l(l+1)}{4} = 0, \quad (234)$$

$$\frac{E}{2\hbar^2 |k|} - \eta_2 - \frac{\mu^2}{4} + d(d+1) = 0. \quad (235)$$

Eq. (234) results in  $s = \frac{l+1}{2}$  and  $s = \frac{l}{2}$  in which we consider  $s = \frac{l+1}{2}$  as we are seeking well-defined function of (208). Eq. (233) is of the form of hypergeometric equation and admits the solution at  $r = 0$  (via (59)) as

$$\phi(z) = C_1 {}_2F_1(a, b; c, z) + C_2 z^{1-c} {}_2F_1(a - c + 1, b - c + 1; 2 - c, z), \quad |z| < 1. \quad (236)$$

Then the solution for (214) can be written as

$$\begin{aligned} \phi(z) = & C_1 {}_2F_1\left(l + d + \frac{5}{4} + \frac{\tilde{\mu}}{2}, l + d + \frac{5}{4} - \frac{\tilde{\mu}}{2}; l + \frac{3}{2}; z\right) \\ & + C_2 z^{-l-\frac{1}{2}} {}_2F_1\left(d + l - \frac{3}{2} - \frac{\tilde{\mu}}{2}, d + l - \frac{3}{2} + \frac{\tilde{\mu}}{2}; \frac{1}{2} - l, z\right), \quad |z| < 1. \end{aligned} \quad (237)$$

It reduces to polynomial form when  $d = -n_r - \frac{l}{2} - \frac{5}{4} + \frac{\tilde{\mu}}{2}$ ,

$$\phi(z) = C_1 {}_2F_1\left(-n_r, n_r + \tilde{\mu}; l + \frac{3}{2}; z\right) + C_2 z^{-l-\frac{3}{2}} {}_2F_1(-n_r - l - \tilde{\mu}, -n_r - l; \frac{1}{2} - l, z), \quad |z| < 1. \quad (238)$$

where  $C_1$  and  $C_2$  are integration constants, which sets the energy eigenvalues (235) as

$$E_{n_r, l} = \left(2n_r + l + \frac{3}{2}\right) \hbar\omega_0 \sqrt{1 + \frac{1}{\mu^2} \left(\frac{9}{4} - 2\eta_1\right)} - \left[\left(2n_r + l + \frac{3}{2}\right)^2 + 2\bar{\alpha} + 2\bar{\gamma} - \frac{1}{4}\right] \frac{\hbar^2 |k|}{2}. \quad (239)$$

And the solution  $\chi_{n, l}(r)$  can be expressed as,

$$\begin{aligned} \chi_{n, l}(r) = & \mathcal{N}_{n, l} \frac{(|k| r^2)^{\frac{l+1}{2}}}{(1 - |k| r^2)^{n_r + \frac{l}{2} - \frac{\mu}{2} + \frac{5}{4}}} \left[ C_1 {}_2F_1\left(-n_r, n_r + \mu; l + \frac{3}{2}; |k| r^2\right) \right. \\ & \left. + C_2 (|k| r^2)^{-l-\frac{1}{2}} {}_2F_1(-n_r - l - \tilde{\mu}, -n_r - l; \frac{1}{2} - l, |k| r^2) \right]. \end{aligned} \quad (240)$$

The eigenstates are bounded if  $n_r + \frac{l}{2} + \frac{5}{4} > \frac{\mu}{2}$  and  $n_r = 0, 1, 2, \dots N$ .

Similarly for the region  $II$ , we can obtain the solution  $\chi_{n, l}(r)$  for  $|r| > \frac{1}{\sqrt{|k|}}$  as

$$\begin{aligned} \chi_{n, l}(r) = & (|k| r^2)^{\frac{l+1}{2}} \left[ C_1 (1 - |k| r^2)^{-\frac{l}{2} - \frac{5}{4} + \frac{\tilde{\mu}}{2}} {}_2F_1\left(d + \frac{l}{2} - \frac{\tilde{\mu}}{2} + \frac{5}{4}, -d - \frac{l}{2} + \frac{\tilde{\mu}}{2} + \frac{1}{4}; 1 - \tilde{\mu}; \frac{1}{1 - |k| r^2}\right) \right. \\ & \left. + C_2 (1 - |k| r^2)^{-\frac{l}{2} - \frac{5}{4} - \frac{\tilde{\mu}}{2}} {}_2F_1\left(d + \frac{l}{2} + \frac{\tilde{\mu}}{2} + \frac{5}{4}, -d - \frac{l}{2} + \frac{\tilde{\mu}}{2} + \frac{1}{4}; 1 + \tilde{\mu}; \frac{1}{1 - |k| r^2}\right) \right], \end{aligned} \quad (241)$$

where we have used (96). The states (241) are continuum states.

### C. Quantum solvability of nonlinear oscillator $V_2(r)$

The Hamiltonian corresponding to the Lagrangian (182) can be expressed as

$$H = \frac{1}{2} \left[ (1 + k\mathbf{q}^2)(\mathbf{p}^2 + k(\mathbf{p} \cdot \mathbf{q})^2) + \frac{\omega_0^2 \mathbf{q}^2}{2(1 + k\mathbf{q}^2)^2} \right]. \quad (242)$$

In polar coordinates, the Hamiltonian can be expressed as

$$H = \frac{1}{2} \left[ \frac{\dot{r}^2}{(1 + kr^2)^2} + \frac{r^2 (\dot{\theta}^2 + \sin^2 \theta \dot{\phi}^2)}{(1 + kr^2)} + \frac{\omega_0^2 r^2}{(1 + kr^2)^2} \right]. \quad (243)$$

On substituting the components of momentum (201) in (243), we can get

$$H = \frac{1}{2} \left[ (1 + kr^2)^2 p_r^2 + \frac{(1 + kr^2)}{r^2} \left( p_\theta^2 + \frac{p_\phi^2}{\sin^2 \theta} \right) + \frac{\omega_0^2 r^2}{(1 + kr^2)^2} \right]. \quad (244)$$

To get normalizable and bounded solutions for the system (244), we consider the non-Hermitian ordered form of the Hamiltonian (203), for the potential  $V(r) = \frac{\omega_0^2 r^2}{2(1 + kr^2)^2}$  with mass  $m(r) = \frac{1}{(1 + kr^2)^2}$  as,

$$\begin{aligned} \hat{H} = & \frac{1}{2} \left( \frac{1}{m} \hat{p}_r^2 - i\hbar(\bar{\alpha} - \bar{\gamma} + 1) \frac{d}{dr} \left( \frac{1}{m} \right) \hat{p}_r \right) + \frac{\hbar^2}{2} \left[ \bar{\gamma} \frac{d^2}{dr^2} \left( \frac{1}{m} \right) + \bar{\alpha}\bar{\gamma} \left( \frac{d}{dr} \left( \frac{1}{m} \right) \right)^2 \right] m \\ & + \frac{1}{\sqrt{m} r^2} \left( \hat{p}_\theta^2 + \frac{\hat{p}_\phi^2}{\sin^2 \theta} \right) \Big] + V(r), \end{aligned} \quad (245)$$

where  $\hat{p}_r = -i\hbar \left( \frac{\partial}{\partial r} + \frac{1}{r} \right)$ ,  $\hat{p}_\theta = -i\hbar \left( \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \right)$  and  $\hat{p}_\phi = -i\hbar \frac{\partial}{\partial \phi}$ . The generalized Schrödinger equation can be written as

$$\begin{aligned} \Phi'' + \left( \frac{2}{r} - (1 + \bar{\alpha} - \bar{\gamma}) \frac{m'}{m} \right) \Phi' + \left[ \frac{(\bar{\gamma} - \bar{\alpha} - 1) m'}{r m} + \bar{\gamma} \frac{m''}{m} - (\bar{\alpha}\bar{\gamma} + 2\bar{\gamma}) \frac{m'^2}{m^2} \right. \\ \left. - \frac{\sqrt{m} \hat{L}^2}{r^2} + \frac{2m}{\hbar^2} (E - V(r)) \right] \Phi = 0. \end{aligned} \quad (246)$$

Again we consider  $\Phi(r, \theta, \phi) = \frac{1}{r} \chi(r) Y(\theta, \phi)$ , with a spherically symmetric factor  $\frac{\chi(r)}{r}$  and  $m(r) = \frac{1}{(1 + kr^2)^2}$  and a generalized angular function  $Y(\theta, \phi)$  which separate the equation (246) into

$$\text{Angular part: } L^2 Y(\theta, \phi) = \hbar^2 l(l+1) Y(\theta, \phi), \quad (247)$$

$$\text{Radial part: } \chi'' + \frac{4(\bar{\alpha} - \bar{\gamma} + 1) kr}{(1 + kr^2)} \chi' + \left[ \frac{4\sigma_1 k - kl(l+1)}{(1 + kr^2)} + \frac{\frac{2E}{\hbar^2} + 4\sigma_2 k}{(1 + kr^2)^2} - \frac{\mu^2 kr^2}{(1 + kr^2)^4} - \frac{l(l+1)}{r^2} \right] \chi = 0, \quad (248)$$

where  $Y_{l,m}(\theta, \phi) = \sqrt{\frac{(l-m)!}{(l+m)!}} e^{im\phi} P_l^m(\cos \theta)$  are spherical harmonics, and  $l, m$  are quantum numbers.

Now we consider the quasi polynomial solution,

$$\chi(r) = \exp\left(-\frac{\mu k r^2}{2(1+k r^2)}\right) (1+k r^2)^d (k r^2)^s U(r), \quad (249)$$

so that Eq. (248) becomes

$$U''(r) + \left[ \frac{4s}{r} + \frac{4(\bar{\alpha} - \bar{\gamma} + 1)kr}{(1+kr^2)} - \frac{2\mu kr}{(1+kr^2)^2} \right] U'(r) + \left[ \frac{4\sigma_r k}{(1+kr^2)} + \frac{4\epsilon_r k}{(1+kr^2)^2} + \frac{4\mu k(d + \bar{\alpha} - \bar{\gamma})}{(1+kr^2)^3} \right] U(r) = 0, \quad (250)$$

where

$$s = \frac{l+1}{2} \text{ and } s = -\frac{l}{2}, \quad (251)$$

$$\sigma_r = \sigma_1 + \frac{l(l+1)}{4} + d \left( d + \frac{3}{2} + 2s + 2\bar{\alpha} - 2\bar{\gamma} \right) + 2s(1 + \bar{\alpha} - \bar{\gamma}), \quad (252)$$

$$\epsilon_r = \frac{E}{2\hbar^2 k} + \sigma_2 - \mu \left( d + s + \bar{\gamma} - \bar{\alpha} + \frac{1}{4} \right) - d(d + 1 + 2\bar{\alpha} - 2\bar{\gamma}). \quad (253)$$

On applying again the transformation (212), we can express Eq. (250) as

$$z(1-z)^2 U''(z) + \left[ 2s + \frac{1}{2} + \left( 2d + 2\bar{\alpha} - 2\bar{\gamma} - 2s - \mu - \frac{1}{2} \right) z + (2\mu - 2d - 2\bar{\alpha} - 2\bar{\gamma}) z^2 - \mu z^3 \right] U'(z) + [\epsilon_r + \sigma_r + \mu(d + \bar{\alpha} - \bar{\gamma}) - z(\epsilon_r + 2\mu(d + \bar{\alpha} - \bar{\gamma})) + z^2 \mu(d + \bar{\alpha} - \bar{\gamma})] U(z) = 0, \quad |z| < 1. \quad (254)$$

This is again of the form of confluent Heun equation which can be solved using Bethe-ansatz method. Using the results of Sec. IV to solve equation (254) as

$$U(z) = \Pi_{i=0}^n (z - z_i) \quad (255)$$

provided the following restrictions on the parameters hold:

$$d = n + \bar{\gamma} - \bar{\alpha}, \quad (256)$$

$$-\epsilon_r = n(n+1) + \mu \sum_{i=1}^n z_i, \quad (257)$$

$$-\epsilon_r - \sigma_r = -\mu \sum_{i=1}^n z_i^2 + 2(\mu - 1) \sum_{i=1}^n z_i + 2n^2 - 2ns - \frac{n}{2}. \quad (258)$$

The energy eigenvalues are

$$E_{n,l} = \left(2n + 2s + \frac{1}{2}\right) \hbar\omega_0 + \left(-\mu \sum_{i=1}^n z_i + \bar{\gamma} - (\bar{\gamma} - \bar{\alpha})(\bar{\gamma} - \bar{\alpha} - 1) + \sigma_1\right) 2\hbar^2 k, \quad (259)$$

with

$$\sigma_1 = \mu \sum_i^n z_i^2 + (2 - \mu) \sum_{i=1}^n z_i - 2n^2 - 2s - \frac{l(l+1)}{4} + (\bar{\gamma} - \bar{\alpha}) \left(\bar{\gamma} - \bar{\alpha} - \frac{3}{2}\right). \quad (260)$$

Since for  $s = -\frac{l}{2}$  we get unbounded solutions, we consider  $s = \frac{l+1}{2}$  and eigenfunctions for the system (243) are

$$\Phi_{n,l}(r, \theta, \phi) = \mathcal{N}_{n,l} e^{\frac{-\omega_0 r^2}{2 \hbar(1+kr^2)}} (kr^2)^{\frac{l}{2}} (1+kr^2)^{n+\bar{\gamma}-\bar{\alpha}} \prod_{i=0}^n \left(\frac{k r^2}{(1+kr^2)} - z_i\right), \quad (261)$$

where  $\mathcal{N}_{n,l}$  is normalization constant and the roots can be found out by using the relation,

$$\sum_{j \neq i}^n \frac{2}{z_i - z_j} = \frac{\mu z_i^2 + (2n - \mu)z_i + (l + \frac{3}{2})}{z_i(z_i - 1)}, \quad (262)$$

The energy eigenvalues (259) are also of quadratic nature similar to the Higgs oscillator and Mathews-Lakshmanan oscillator. While solving the system (182) quantum mechanically the ordering parameters play important role since  $\sigma_1$  and  $\sigma_2$  (vide (127) and (128)) are also being related with  $n$ . This is possible when the ordering parameters are treated as arbitrary. To obtain the eigenfunctions for different values of  $n$ , we have to find out the roots  $z_i$ ,  $i = 0, 1, 2, \dots, n$ , by solving the equation in  $z_i$  of degree  $n+1$ . Hence it is possible only to obtain a few states of the system (243).

**Bound state:**  $n = 0, l = 0$

For  $n = 0$ , Eq. (260) results in

$$\sigma_1 = (\bar{\gamma} - \bar{\alpha}) \left(\bar{\gamma} - \bar{\alpha} - \frac{3}{2}\right) - 1. \quad (263)$$

Then, the ground state solution is

$$\Phi_{0,0}(r, \theta, \phi) = \mathcal{N}_{0,0} \exp \left[ \frac{-\omega_0 r^2}{2 \hbar(1+kr^2)} \right] (1+kr^2)^{\bar{\gamma}-\bar{\alpha}}, \quad (264)$$

$$E_{0,0} = \frac{3}{2} \hbar\omega_0 + (\bar{\alpha} + \bar{\gamma} - 2) \hbar^2 k, \quad (265)$$

which is bounded when  $\bar{\alpha} - \bar{\gamma} > 0$  and the normalization constant,  $\mathcal{N}_{n,l}$ , is found to be

$$\mathcal{N}_{0,0} = \left[ \frac{\mu^{3/2} \sqrt{k}}{\pi (\operatorname{erf}(\sqrt{\mu}) - 2\sqrt{\mu} e^{-\mu})} \right]^{1/2}, \quad k > 0, \quad (266)$$

$$\mathcal{N}_{0,0} = \left[ -\frac{\mu^{3/2} \sqrt{|k|}}{\pi} \right]^{1/2}, \quad k < 0, \quad (267)$$

where  $\bar{\alpha} - \bar{\gamma} = \frac{5}{4}$ .

**Bound state:**  $n = 1, l = 0$  and  $l = 1$

The first excited state is

$$\Phi_{1,0}(r, \theta, \phi) = \mathcal{N}_{1,0} \exp \left[ \frac{-\omega_0 r^2}{2 \hbar (1 + kr^2)} \right] (1 + kr^2)^{1+\bar{\gamma}-\bar{\alpha}} \left( \frac{kr^2}{1 + kr^2} - z_1 \right), \quad (268)$$

$$E_{1,0} = \frac{7}{2} \hbar \omega_0 + (\bar{\alpha} + \bar{\gamma} - 9 - 2\mu z_1) \hbar^2 k, \quad (269)$$

which is obtained from

$$\sigma_1 = \mu z_1^2 + (2 - \mu) z_1 - 3 + \left( \bar{\gamma} - \bar{\alpha} - \frac{3}{2} \right) (\bar{\gamma} - \bar{\alpha}), \quad (270)$$

and  $z_1$  satisfies

$$\mu z_1^2 + (2 - \mu) z_1 + \frac{3}{2} = 0. \quad (271)$$

Eq. (271) results in

$$z_1 = \left( \frac{1}{2} - \frac{1}{\mu} \right) \pm \frac{1}{2\mu} \sqrt{\mu^2 - 10\mu + 4}. \quad (272)$$

The normalization constant can be obtained as

$$\mathcal{N}_{1,0} = \left[ \frac{2k\sqrt{k}}{-\sqrt{\pi} \left( \frac{\partial^3}{\partial \mu^3} + 2 z_1 \frac{\partial^2}{\partial \mu^2} + z_1^2 \frac{\partial}{\partial \mu} \right) \frac{erf(\sqrt{\mu})}{\sqrt{\mu}}} \right]^{1/2}, \quad k > 0, \quad (273)$$

$$\mathcal{N}_{1,0} = \left[ \frac{8\mu^{7/2} \sqrt{|k|}}{\sqrt{\pi} (15 + 3\mu z_1 + z_1^2)} \right]^{1/2}, \quad k < 0, \quad (274)$$

where  $\bar{\alpha} - \bar{\gamma} = \frac{9}{4}$ .

The bound state for  $n = 1$  and  $l = 1$  can be obtained as

$$\Phi_{1,1}(r, \theta, \phi) = \mathcal{N}_{1,1} \exp \left[ \frac{-\omega_0 r^2}{2 \hbar (1 + kr^2)} \right] \sqrt{kr} (1 + kr^2)^{1+\bar{\gamma}-\bar{\alpha}} \left( \frac{kr^2}{1 + kr^2} - z_1 \right), \quad (275)$$

$$E_{1,1} = \frac{9}{2} \hbar \omega_0 + (\bar{\alpha} + \bar{\gamma} - 14 - 2\mu z_1) \hbar^2 k, \quad (276)$$

where the root  $z_1$  satisfies the equation

$$\mu z_1^2 + (2 - \mu) z_1 + \frac{5}{2} = 0 \quad (277)$$

with the constraint on the ordering parameters  $\sigma_1$  as

$$\sigma_1 = \mu z_1^2 + (2 - \mu) z_1 - \frac{9}{2} + \left( \bar{\gamma} - \bar{\alpha} - \frac{3}{2} \right) (\bar{\gamma} - \bar{\alpha}). \quad (278)$$



The normalization constant  $\mathcal{N}_{1,1}$  is of the form as

$$\mathcal{N}_{1,1} = \left[ \frac{\sqrt{k}}{2\pi\sqrt{\pi} \left( \frac{\partial^4}{\partial \mu^4} - 2 z_1 \frac{\partial^3}{\partial \mu^3} + z_1^2 \frac{\partial^2}{\partial \mu^2} \right) \frac{\text{erf}(\sqrt{\mu})}{\sqrt{\mu}}} \right]^{1/2}, \quad k > 0, \quad (279)$$

$$\mathcal{N}_{1,1} = \left[ \frac{\mu^{9/2}\sqrt{k}}{\pi \left( \frac{7!!}{4} + 5!! \mu z_1 + \mu^2 3!! z_1^2 \right)} \right]^{1/2}, \quad k < 0, \quad (280)$$

where  $\bar{\alpha} - \bar{\gamma} = \frac{9}{4}$ . As in the case of one dimensional system (167), we again fixed the ordering parameter to normalize the solutions (264), (268) and (275).

For  $n = 2$ , we will get the coupled equations of  $z_1$  and  $z_2$  from (260) and (262) which can not be solved analytically. Hence, the system (242) is quasi exactly solvable.

## VI. CONCLUSION

We have studied the classical and quantum dynamics of two one dimensional nonlinear oscillators, namely the (i) Higgs oscillator and (ii) a  $k$ -dependent nonpolynomial rational potential. Both the systems share the same mass functions but differ in their potential forms and the dynamics of the two systems are different. We obtained explicitly the solutions of the Higgs oscillator and the non-polynomial  $k$ -dependent potential for their classical dynamics. The former solution is expressed in terms of trigonometric functions whereas for the second system the classical solution can be given in terms of Jacobian elliptic functions. The  $k$ -dependent nature of the potential gives rise to a complex solution. In the quantum study, the Higgs oscillator is exactly solvable for all possible choices of ordering parameters, whereas the second system is quasi exactly solvable and it possesses quasi-polynomial solutions with a requirement that the ordering parameters are arbitrary. We also studied the classical and quantum dynamics of the three dimensional generalization of the above two nonlinear systems.

## Acknowledgement

VC wishes to acknowledge DST for the financial support of the project (No. SR/WOS-A/PM-64/2018(G)) under Women Scientist Scheme A. ML acknowledges the financial sup-

port under a DST-SERB Distinguished Fellowship program (Grant No. SB/DF/04/2017).

---

- [1] P. M. Mathews and M. Lakshmanan, Q. Appl. Math. **32**, 215 (1974)
- [2] P. M. Mathews and M. Lakshmanan, Nuovo Cimento A **26**, 299 (1975)
- [3] M. Lakshmanan and K. Eswaran, J. Phys. A: Math. Gen. **8**, 1658 (1975)
- [4] M. F. Rañada and M. Santander, J. Math. Phys., **43**, 431 (2002)
- [5] M. Lakshmanan and V. K. Chandrasekar, The European Physical Journal Special Topics, **222**, 665688 (2013)
- [6] C. Quesne, J. Math. Phys. **59**, 042104 (2018).
- [7] P. W. Higgs, J. Phys. A: Math. Gen. **12**, 309 (1979); H. I. Leemon, J. Phys. A: Math. Gen. **12**, 489 (1979)
- [8] J. F. Cariñena, M. F. Rañada, M. Santander and M. Senthilvelan, Nonlinearity **17**, 1941 (2004)
- [9] J. F. Cariñena, M. F. Rañada and M. Santander, Annals of Physics, **322**, 434 (2007)
- [10] R. Delbourgo, A. Salam and J. Strathdee, Phys. Rev., **187**, 1999 (1969)
- [11] V. Chithiika Ruby, V. K. Chandrasekar, M. Senthilvelan and M. Lakshmanan, J. Math. Phys., **56**, 012103 (2015)
- [12] S. Karthiga, V. Chithiika Ruby and M. Senthilvelan, Phys. Lett. A., **382**, 1645-1650 (2018)
- [13] M. S. Marinov and V. S. Popov, J. Phys. A: Math. Gen., **8**, 1575 (1975)
- [14] B. Midya and B. Roy, J. Phys. A: Math. Theor., **42**, 285301 (2009).
- [15] A. Schulze-Halberg and B. Roy, J. Math. Phys., **54**, 122104 (2013).
- [16] A. Trabelsi, F. Madouri, A. Merdaci and A. Almatar, *Classification scheme for kinetic energy operators with position-dependent mass* arXiv.org:1302.3963v1 (2013).
- [17] I. S. Gradshteyn and I. M. Ryzhik. Table of Integrals, Series and Products. Academic Press, New York, (1980).
- [18] S. Karthiga, V. Chithiika Ruby, M. Senthilvelan and M. Lakshmanan, J. Math. Phys., **58**, 102110 (2017).
- [19] Y.-Z. Zhang, J. Phys. A: Math. Theor., **45**, 065206 (2012); D. Agboola and Y. -Z. Zhang. J. Math. Phys., **53**, 042101 (2012)
- [20] C. Quesne, Acta Polytech., **58**, 118-127 (2018).