

KITCHEN & KETTLE

First Aid Checklist

Practical first aid for the cuts,
burns, bites, and mishaps of
everyday life — plus a few
remedies that actually work.

WORKS BECAUSE IT WORKS.

How to use this checklist

This isn't a medical textbook. It's the stuff that happens in real kitchens and households — things you can handle yourself, and a clear line for when you shouldn't.

Each section covers one type of injury. Read the steps, check them off as you go, and keep this somewhere you can grab it with one hand while the other hand is under cold water.

If you're not sure, get help. This checklist covers minor injuries. If something seems serious — heavy bleeding that won't stop, deep puncture wounds, head injuries, anything involving eyes or joints — skip the checklist and call a doctor or 911.

STOCK YOUR FIRST AID KIT

- ☐ Adhesive bandages (assorted sizes)
- ☐ Sterile gauze pads & medical tape
- ☐ Antiseptic wipes or spray
- ☐ Mild soap (for cleaning wounds)
- ☐ Antibiotic ointment
- ☐ Tweezers (sharp point)
- ☐ Small scissors
- ☐ Hydrocortisone cream
- ☐ Calamine lotion
- ☐ Aloe vera gel or burn cream
- ☐ Ace bandage / elastic wrap
- ☐ Instant cold pack
- ☐ Pain reliever (ibuprofen or acetaminophen)
- ☐ Antihistamine (Benadryl or similar)
- ☐ Thermometer
- ☐ Non-stick sterile dressings
- ☐ Heating pad
- ☐ Small needle (sterilized, for splinters)
- ☐ Disposable gloves
- ☐ Duct tape (yes, duct tape — see splinters)
- ☐ Rubbing alcohol

Emergency Contacts

Fill these in and keep this page where anyone in the house can find it.

911 / EMERGENCY

POISON CONTROL

FAMILY DOCTOR

PEDIATRICIAN

NEAREST HOSPITAL

NEAREST URGENT CARE

PHARMACY

VETERINARIAN

NEIGHBOR / NEARBY

HOUSEHOLD INFO

HOME ADDRESS

LANDLINE (IF ANY)

KNOWN ALLERGIES

MEDICAL CONDITIONS

KITCHEN & KETTLE • WORKS BECAUSE IT WORKS.

Cuts & Scrapes

For minor cuts

1. Wash your hands first.
2. Rinse the wound under clean running water. No hydrogen peroxide — it damages tissue.
3. Clean around the wound with mild soap. Try not to get soap in the wound itself.
4. Apply antibiotic ointment.
5. Cover with a bandage. Change daily or when wet or dirty.

Pressure stops bleeding. If a cut won't stop bleeding, apply firm, steady pressure with a clean cloth for 10–15 minutes without peeking. Most bleeding stops in that window. If it soaks through, add more cloth on top — don't remove the first layer.

FOLK REMEDY THAT WORKS

Yarrow for cuts. Yarrow grows in fields and along roadsides — feathery leaves, flat clusters of small white flowers. Crush a few fresh leaves between your fingers and press them directly onto a bleeding cut. Yarrow is a natural styptic — it helps slow bleeding and has been used as a field dressing for centuries. Once the bleeding stops, clean the wound properly and bandage as usual. This is a stopgap for when you're outside and bleeding, not a replacement for cleaning and bandaging.

When to get medical help

1. Bleeding doesn't stop after 15 minutes of direct pressure
2. The wound is deep, gaping, or you can see fat or muscle
3. It's a puncture wound (especially from a nail or animal bite)
4. It's on the face, a joint, or anywhere scarring matters
5. Numbness, tingling, or you can't move the area
6. Signs of infection: redness spreading, warmth, pus, fever

KITCHEN & KETTLE • WORKS BECAUSE IT WORKS.

Burns

For minor burns (red, painful, no blistering)

1. **Cool it immediately.** Run under cool (not cold) water for at least 10 minutes. This is the single most important step — it stops the burn from deepening.
2. Remove any jewelry or tight clothing near the burn before swelling starts.
3. Apply aloe vera gel or a burn cream. No butter, no oil, no ice — these trap heat or damage tissue.
4. Cover loosely with sterile gauze if the area might get rubbed or dirty.
5. Ibuprofen or acetaminophen helps with pain.

For blistering burns

1. Cool under water as above. Do not pop blisters — the skin is protecting the wound underneath.
2. Cover loosely with a non-stick sterile dressing.
3. Change dressing daily. Watch for infection.
4. Burns larger than 3 inches across, or on the face, hands, feet, groin, or a major joint — get medical help.

Kitchen tip: For a minor contact burn from a pan handle or baking sheet, cool it, then a thin layer of antibiotic ointment and a bandage is usually enough. The second you feel pain, get it under water — the faster you cool it, the less damage.

KITCHEN & KETTLE • WORKS BECAUSE IT WORKS.

Splinters & Slivers

If the splinter is sticking out

1. Wash the area with soap and water.
2. Clean tweezers with rubbing alcohol.
3. Grasp the splinter as close to the skin as possible and pull in the direction it entered.
4. Wash again and apply antibiotic ointment.

If the splinter is under the surface

1. Soak the area in warm water for a few minutes to soften the skin.
2. If you can see the splinter but can't grab it, use a sterilized needle to gently lift the skin over the tip.
3. Once the tip is exposed, grab with tweezers and pull.

FOLK REMEDY THAT WORKS

Duct tape for splinters. For superficial splinters — especially fiberglass, wood slivers, or cactus spines — press a strip of duct tape firmly over the area and pull it off in the direction opposite to how the splinter entered. The adhesive grabs the tips and pulls them out. Works best on clusters of tiny slivers where tweezers are hopeless. Follow with soap and water.

When to get help: The splinter is deep, under a fingernail, or the area becomes red, warm, or pus-filled. Splinters carry bacteria — don't ignore one that's been in for more than a day.

KITCHEN & *KETTLE* • WORKS BECAUSE IT WORKS.

Sprains & Strains

R.I.C.E. method — first 48 hours

1. **Rest.** Stop using the joint. Don't try to "walk it off."
2. **Ice.** Cold pack or bag of frozen peas wrapped in a thin towel. 20 minutes on, 20 minutes off. No ice directly on skin.
3. **Compression.** Wrap with an elastic bandage — snug but not tight. If the area tingles, goes numb, or turns blue, loosen immediately.
4. **Elevate.** Keep the injured area raised above heart level when you can.

FOLK REMEDY THAT WORKS

Comfrey (knitbone) poultice for sprains. Comfrey — sometimes called knitbone — has been used for centuries to speed healing of sprains, strains, and bruises. Crush fresh leaves or rehydrate dried leaves with warm water to make a paste, apply it to the swollen area, and wrap loosely with a cloth. Leave on for several hours or overnight. The plant contains allantoin, which promotes cell repair. **Important:** Comfrey is for topical use only. Do not ingest it — the plant contains compounds that can damage the liver when taken internally. Do not apply to open wounds or broken skin.

After 48 hours: Switch to warmth (heating pad, warm compress) to increase blood flow and healing. Gentle movement helps — don't immobilize longer than necessary.

When to get medical help

1. You can't put any weight on it at all
2. The joint looks deformed or out of place
3. You heard a pop or crack at the time of injury
4. Numbness or tingling below the injury
5. Swelling that doesn't improve after 24 hours of R.I.C.E.

KITCHEN & KETTLE • WORKS BECAUSE IT WORKS.

Insect Bites & Stings

For most bites and stings

1. **Remove the stinger if there is one.** Scrape it out sideways with a fingernail or credit card edge — don't pinch it, which squeezes more venom in.
2. Wash with soap and water.
3. Apply ice (wrapped in cloth) for 10 minutes to reduce swelling.
4. Hydrocortisone cream or calamine lotion for itching.
5. Oral antihistamine (Benadryl) if swelling or itching is significant.

For ticks

1. Use fine-tipped tweezers. Grasp as close to the skin as possible (by the head, not the body).
2. Pull straight up with steady, even pressure. Don't twist or jerk.
3. Clean the bite with rubbing alcohol or soap and water.
4. Save the tick in a sealed bag if you want it identified later.
5. Watch for a bullseye rash, fever, or joint pain in the following weeks.

FOLK REMEDY THAT WORKS

Baking soda paste for itching. Mix a little baking soda with water to make a thick paste. Apply to mosquito bites or bee stings and let it dry. The alkalinity helps neutralize the itch. Not a substitute for antihistamines if you're having a strong reaction, but it'll calm a normal bite down.

FOLK REMEDY THAT WORKS

Narrow leaf plantain poultice. If you're outdoors and get stung, look for narrow leaf plantain — the common weed with long, ribbed leaves, not the banana-like fruit. Chew or crush a few leaves into a moist wad, press it onto the sting, and hold it there. The leaves draw out the sting and reduce swelling faster than waiting it out. Works on bee stings, wasp stings, and mosquito bites. Spit poultice isn't elegant, but it works.

Allergic reactions — get emergency help if: Difficulty breathing, swelling of the face or throat, hives spreading beyond the sting site, dizziness, or nausea. These are signs of anaphylaxis. Use an EpiPen if you have one and call 911.

KITCHEN & KETTLE • WORKS BECAUSE IT WORKS.

Ears

Water trapped in ear

1. Tilt your head so the affected ear faces down.
2. Gently tug on the earlobe and jiggle it while tilting.
3. Try yawning, chewing, or the Valsalva maneuver (pinch nose, close mouth, blow gently).

FOLK REMEDY THAT WORKS

Rubbing alcohol in the ear for trapped water. A few drops of rubbing alcohol in the affected ear, let it sit for 30 seconds, then tilt to drain. The alcohol breaks the surface tension of the water and evaporates quickly, taking the water with it. Works best right after swimming or showering. Don't use this if you have ear tubes, a perforated eardrum, or an active ear infection.

Ear pain / possible infection

1. A warm (not hot) compress held against the ear can relieve pain.
2. Over-the-counter pain reliever.
3. Do not put anything in the ear — no drops, no oil, no cotton swabs — unless a doctor tells you to.
4. See a doctor if: pain is severe, there's drainage or pus, hearing is affected, or fever develops.

KITCHEN & *KETTLE* • WORKS BECAUSE IT WORKS.

Minor Kitchen Injuries

Minor knife nick on fingertip

1. Rinse under cool water.
2. Apply firm pressure with a clean paper towel for 5–10 minutes. Don't lift to check too often — it breaks the clot.
3. Once bleeding stops, clean gently, apply antibiotic ointment, and wrap with a bandage.
4. If you can't bend the finger or feel it properly, get it checked — you may have hit a tendon or nerve.

Oil splatter burn

1. Cool water immediately, same as any burn. Ten full minutes.
2. Oil burns tend to be small but can be deep. Treat as a second-degree burn — don't pop blisters.
3. Aloe or burn gel, loose gauze covering.

Chili pepper in the eye

1. Flush with cool water — lots of it. Tilt your head and let water run across the eye from inner corner to outer.
2. Do not rub — it spreads the capsaicin.
3. Milk or yogurt on a damp cloth held gently over the closed eye can help (dairy fat binds capsaicin).
4. Wash your hands thoroughly with soap before touching anything else.

PREVENTION

When handling hot peppers, wear disposable gloves or coat your hands with a thin layer of oil first (capsaicin is oil-soluble and washes off more easily). After handling peppers, wash hands with dish soap — not hand soap — before touching your face.

KITCHEN & *KETTLE* • WORKS BECAUSE IT WORKS.

Nosebleeds

What to do

1. **Sit up and lean forward.** Tilting your head back sends blood down your throat, which can cause nausea or choking. Lean forward so it drains out.
2. **Pinch the soft part of your nose** — the fleshy sides below the bridge — with your thumb and forefinger. Hold steady pressure for 10–15 minutes without releasing to check.
3. Breathe through your mouth. Spit out any blood that reaches your throat.
4. After the bleeding stops, avoid blowing your nose or bending over for a few hours.

Most nosebleeds are from dry air or minor irritation — not a medical emergency. A humidifier at night helps prevent them. If you get them frequently, a dab of petroleum jelly inside each nostril at bedtime keeps the lining from drying out.

When to get medical help

1. Bleeding doesn't slow after 20 minutes of continuous pressure
2. Bleeding is heavy and you feel lightheaded or weak
3. It follows a head injury (possible skull fracture)
4. You're on blood thinners and can't get it to stop

KITCHEN & *KETTLE* • WORKS BECAUSE IT WORKS.

Choking

If the person can cough or make sounds

1. **Encourage them to keep coughing.** A strong cough is the body's way of clearing the airway. Don't slap their back — it can lodge the object deeper.
2. Stay with them. If coughing weakens or stops, move to the next section.

If the person cannot breathe, cough, or speak

1. **Call 911.** Have someone call while you start. If you're alone, do abdominal thrusts for 2 minutes first, then call.
2. **Stand behind them.** Wrap your arms around their waist. Make a fist with one hand, thumb side against their stomach, just above the navel and below the ribcage.
3. Grab your fist with your other hand and **thrust sharply inward and upward.** Repeat until the object comes out or they lose consciousness.
4. If they lose consciousness, lower them to the ground, start CPR if you know it, and continue until help arrives.

If you're alone and choking

1. Call 911 first if you can — even if you can't speak, an open line will bring help.
2. Make a fist and place it above your navel. Use your other hand to thrust inward and upward.
3. Or lean over the back of a chair or counter edge, driving it into your upper abdomen with your body weight.

For an infant (under 1 year)

1. Sit down and lay the baby face-down along your forearm, head lower than chest. Support the head and jaw.
2. Give **5 firm back blows** between the shoulder blades with the heel of your hand.
3. If the object doesn't come out, turn the baby face-up on your thigh, head still lower than chest. Give **5 chest thrusts** using two fingers on the breastbone.
4. Alternate 5 back blows and 5 chest thrusts. Call 911 after one minute if the object hasn't cleared.

KITCHEN & KETTLE • WORKS BECAUSE IT WORKS.

Eye Injuries

Debris in the eye (dust, sand, eyelash)

1. **Do not rub.** Rubbing can scratch the cornea.
2. Blink rapidly — your tears may flush it out.
3. Pull the upper eyelid over the lower eyelid and release. The lower lashes can sweep debris away.
4. If that doesn't work, flush with clean water or saline. Tilt your head so the affected eye is lower, and pour water from the inner corner outward.

Chemical splash

1. **Flush immediately.** Hold the eye open under a gentle stream of cool water for at least 15 minutes. Use a sink, shower, or hose.
2. Remove contact lenses if you wear them — don't delay flushing to do this.
3. Do not rub. Do not apply anything to the eye.
4. Call Poison Control (1-800-222-1222) or go to the ER after flushing. Bring the chemical container if you can.

Blow or impact to the eye

1. Apply a cold compress gently — no pressure. A bag of frozen peas wrapped in a cloth works.
2. Do not apply pressure or rub.
3. Get medical help if: vision changes, you see flashes or floaters, there's blood in the eye, the eye looks sunken or protruding, or you can't move it normally.

KITCHEN & *KETTLE* • WORKS BECAUSE IT WORKS.

Poisoning

If you suspect poisoning

1. **Call Poison Control immediately:** 1-800-222-1222. They're available 24/7 and their guidance is free. Put this number in your phone.
2. Stay calm. Have the container or bottle with you — the label tells them exactly what was ingested.
3. Do **not** induce vomiting unless Poison Control specifically tells you to. Some substances cause more damage coming back up.
4. Do not give the person anything to eat or drink unless instructed.

Call 911 instead of Poison Control if the person: collapses, has a seizure, stops breathing, or can't be woken up. Poison Control handles the substance questions; 911 handles life-threatening emergencies.

Inhaled poison (fumes, smoke, gas)

1. Get the person to fresh air immediately. Open windows and doors.
2. Do not enter an enclosed space yourself if fumes are strong — you could become a second victim.
3. Call 911. Check breathing and start CPR if needed and you know how.

Poison on the skin

1. Remove contaminated clothing carefully — don't pull it over the face.
2. Rinse skin with water for at least 15 minutes.
3. Call Poison Control for further instructions.

KITCHEN & *KETTLE* • WORKS BECAUSE IT WORKS.

When to Call a Doctor

Most household injuries are minor and heal on their own. These are the signals that something needs professional attention.

GO TO THE ER OR CALL 911

- ☐ Heavy bleeding that won't stop with direct pressure
- ☐ Difficulty breathing
- ☐ Chest pain or pressure
- ☐ Head injury with confusion, vomiting, or loss of consciousness
- ☐ Severe allergic reaction (swelling face/throat, trouble breathing)
- ☐ Severe burn — large, deep, or on face/hands/feet/joints
- ☐ Possible broken bone — deformity, can't move it, severe pain
- ☐ Poisoning — call Poison Control first (1-800-222-1222) and follow their instructions

SEE A DOCTOR WITHIN 24 HOURS

- ☐ Deep cut that may need stitches
- ☐ Animal bite that broke the skin
- ☐ Puncture wound (especially from a nail or dirty object)
- ☐ Signs of infection: spreading redness, warmth, swelling, pus, fever
- ☐ Burn that blisters over more than 3 inches
- ☐ Sprain that can't bear weight or has significant swelling
- ☐ Something stuck in the eye that won't flush out
- ☐ Tick bite followed by rash, fever, or joint pain (Lyme concern)

Trust your gut. If something feels wrong and you can't pin down why, that's a valid reason to call a doctor. This list covers common patterns — it doesn't cover everything. You know your body and your family better than a checklist does.

KITCHEN & *KETTLE* • WORKS BECAUSE IT WORKS.

KITCHEN & KETTLE

Take care of each other.

Most injuries are small. Most recoveries are quick. The skill isn't in having no accidents — it's in staying calm and knowing what to do when they happen.

Print extras. Tuck one in the kitchen drawer. Give one to a new parent.

KITCHEN & KETTLE • WORKS BECAUSE IT WORKS.