

QUANTUM MECHANICS

IVAN IORSH • AUTUMN 2015 • ITMO UNIVERSITY

Last Revision: October 14, 2015

Table of Contents

1	Introduction	2
1.1	Schrödinger formalism	2
1.2	Heisenberg formalism	2
	Building an operator's matrix	3
	$f_n m(t)$ time dependence	3
	Operator matrix properties	4
1.3	Switching to a different state basis	6
1.4	Pauli uncertainty principle	8
	Single-slit electron diffraction	8
	Black holes	9
	Quantum Pencil	13
1.5	Problems	14
2	Analytical Solutions	15
2.1	Rectangular quantum well	15
2.2	Harmonic oscillator	15
2.3	Spherically symmetric potential	15
2.4	Problems	15
	Rectangular quantum well	15
3	Quasi-classical approximation	16
3.1	Problems	16
4	Spin	17
4.1	Angular momentum	17
	Classical	17
	Quantum	17
4.2	Problems	17
5	Perturbation theory	18
5.1	Time-independent	18
5.2	Time-dependent	18
5.3	Problems	18

6 Problem Solutions	19
6.1 Introduction	19
6.2 Analytical solutions	19
6.3 Quasiclassical approximation	19
6.4 Spin	19
6.5 Perturbation theory	19
References	20

Abstract

Quantum Mechanics Lecture Notes.

TODO:

- Language - Русский or English?
- Определиться с стилистикой доказательств - всякие новомодные ЧТД \square и прочее

Dual language Example

Двуязычность можно например реализовать
вот так... blah blah blahy

And here some english text

1 Introduction

1.1 Schrödinger formalism

$$\hat{f}\Phi = E\Phi \quad (1.1)$$

$$\Phi \rightarrow dP = |\Phi|^2 dq \quad (1.2)$$

$$x \leftrightarrow \hat{x} \quad (1.3)$$

$$p_x \leftrightarrow -i\hbar \frac{\partial}{\partial x} \quad (1.4)$$

$$f \leftrightarrow \hat{f} \quad (1.5)$$

$$\bar{f} = \int \hat{f} dp = \int \Phi^* \hat{f} \Phi dq \quad (1.6)$$

$$[\hat{f}, \hat{g}] = \hat{f}\hat{g} - \hat{g}\hat{f} \quad (1.7)$$

$$\{\hat{f}, \hat{g}\} = \hat{f}\hat{g} + \hat{g}\hat{f} \quad (1.8)$$

[1]

1.2 Heisenberg formalism

Schrödinger was good at math, which is why his quantum mechanics formalism is full of complex mathematical constructs. Heisenberg, on the other hand, had a lot of difficulty with math, which is why his matrix quantum mechanics formalism is limited almost exclusively to linear algebra constructs

Roman ...

Name	Schrödinger	Heisenberg
State Basis	Wave function of basis states $\{\Phi_n\}$	Column vector of basis states $\begin{pmatrix} \phi_1 \\ \dots \\ \phi_n \end{pmatrix}$
Observables	Operator $\bar{f} = \int \Phi_n^* \hat{f} \Phi_m$	Operator matrix $\begin{pmatrix} \phi_{11} & \dots & \phi_{n1} \\ & \dots & \\ \phi_{1n} & \dots & \phi_{nn} \end{pmatrix}$
Schrödinger Equation	$\hat{f}\Phi = E\Phi$	$\begin{pmatrix} \phi_{11} & \dots & \phi_{n1} \\ & \dots & \\ \phi_{1n} & \dots & \phi_{nn} \end{pmatrix} \begin{pmatrix} \psi_1 \\ \dots \\ \psi_n \end{pmatrix} = \lambda \begin{pmatrix} \psi_1 \\ \dots \\ \psi_n \end{pmatrix}$

Building an operator's matrix

For a system with a discrete state basis, $\{\Psi_n\}$, any state of the system can be described as a linear combination of the basis' wave functions:

$$\Psi = \sum_n a_n \Psi_n \quad (1.9)$$

Observable \bar{f} for such a wave function can be decomposed into a sum over the basis state wave functions:

$$\begin{aligned}
 \bar{f} &= \int \Psi^* \hat{f} \Psi dq = \int \sum_n a_n^* \Psi_n^* \hat{f} \sum_m a_m \Psi_m dq \\
 &= \sum_n \sum_m a_n^* a_m \int \Psi_n^* \hat{f} \Psi_m dq = \sum_n \sum_m a_n^* a_m f_{nm}(t) \\
 &= \sum_n \sum_m a_n^* f_{nm}(t) a_m
 \end{aligned} \quad (1.10)$$

Where f_{nm} is the operator matrix.

$f_{nm}(t)$ time dependence

Move to appendix?

Solutions to the time-independent Schrödinger equation:

$$\hat{H}\Psi_n = E_n \Psi_n \quad (1.11)$$

$$\Psi_n(t) = e^{-\frac{i}{\hbar} E_n t} \Phi_n \quad (1.12)$$

Which, in operator matrix terms translates into

$$\begin{aligned}
f_{nm}(t) &= \int \Psi_n^* \hat{f} \Psi_m dq = \int \Phi_n^* (e^{-\frac{i}{\hbar} E_n t})^* \hat{f} \Phi_m e^{-\frac{i}{\hbar} E_m t} dq \\
&= e^{+\frac{i}{\hbar} E_n t} e^{-\frac{i}{\hbar} E_m t} \int \Phi_n^* \hat{f} \Phi_m dq = e^{i \frac{E_n - E_m}{\hbar} t} \int \Phi_n^* \hat{f} \Phi_m dq \\
&= f_{nm} e^{i \omega_{nm} t}
\end{aligned} \tag{1.13}$$

Operator matrix properties

1. The operator matrix is hermitian ¹ Transposed operator:

$$\left(\int \Phi \hat{f} \Psi dq \right)^T = \int \Psi (\hat{f})^T \Phi dq \tag{1.14}$$

Complex conjugate:

$$(\hat{f})^* = \hat{f}^* \tag{1.15}$$

Hermitian conjugate:

$$\bar{f}^* = \int \Psi^* \hat{f}^\dagger \Psi dq \tag{1.16}$$

In operator matrix terms:

$$\begin{aligned}
(f_{nm}^*) &= \int \varphi_n^* \hat{f}^\dagger \varphi_m dq = \int \varphi_n^* (\hat{f}^*)^T \varphi_m dq \\
&= \int \varphi_m (\hat{f}^* \varphi_n^*) dq = \left(\int \varphi_m^* \hat{f}^\dagger \varphi_n dq \right)^* = (f_{mn})^*
\end{aligned} \tag{1.17}$$

Which means, if f_{nm} is real, meaning $f_{nm}^* = f_{nm}$ that

$$f_{nm} = f_{mn}^* = f_{nm}^\dagger \tag{1.18}$$

2. The matrix' diagonal elements are time-independent and real

$$f_{nn} = \int \Psi_n \hat{f} \Psi_n dq \equiv \bar{f}_n \tag{1.19}$$

Where \bar{f}_n is the value of observable f in basis state n .

3. The matrix of the product of two operators is the product of their matrices

For operators \hat{f} and \hat{g} , what is the operator matrix for operator $\hat{f} \times \hat{g} \rightarrow (\hat{f} \times \hat{g})_{nm}$?

¹ $H^\dagger = H = (H^*)^T$

Move to appendix?

$$\hat{f}\varphi_n = \sum_m f_{mn}\varphi_m \quad (1.20)$$

$$\begin{aligned} \int \varphi_k^* dq \times \hat{f}\varphi_n &= \int \varphi_k^* dq \times \sum_m f_{mn}\varphi_m \\ \int \varphi_k^* \hat{f}\varphi_n dq &= \sum_m f_{mn} \int \varphi_k^* \varphi_m dq f_{kn} = \sum_m f_{mn} \delta_{km} = f_{kn} \end{aligned} \quad (1.21)$$

Because for state basis φ_n , φ_n and φ_m are orthogonal for all $m \neq n$.

Using 1.18, we can write:

$$\begin{aligned} \hat{f}\hat{g}\varphi_n &= \hat{f}(\hat{g}\varphi_n) = \hat{f} \sum_k g_{kn}\varphi_k = \sum_k g_{kn}\hat{f}\varphi_k \\ &= \sum_k g_{kn} \sum_m f_{mk}\varphi_m = \sum_{k,m} g_{kn}f_{mk}\varphi_m \\ &= \sum_{k,m} f_{mk}g_{kn}\varphi_m \end{aligned} \quad (1.22)$$

And knowing that:

$$(\hat{f}\hat{g})\varphi_n = \sum_m (\hat{f}\hat{g})_{nm}\varphi_m \quad (1.23)$$

We end up with:

$$(\hat{f}\hat{g})_{nn} = \sum_k f_{mk}g_{kn} \quad (1.24)$$

4. The operator's matrix is equivalent to the operator

$$\Psi = \sum_m c_m \varphi_m \quad (1.25)$$

$$\hat{f}\Psi = f\Psi \quad (1.26)$$

$$\begin{aligned} \hat{f} \sum_m c_m \varphi_m &= f \sum_m c_m \varphi_m \\ \int \varphi_n^* \hat{f} \sum_m c_m \varphi_m dq &= \int \varphi_n^* f \sum_m c_m \varphi_m dq \\ \sum_m c_m \int \varphi_n^* \hat{f} \varphi_m dq &= f \sum_m c_m \int \varphi_n^* \varphi_m dq \\ \sum_m c_m f_{nm} &= f \sum_m c_m \delta_{nm} \\ \sum_m c_m f_{nm} &= f c_n \end{aligned} \quad (1.27)$$

$$(1.28)$$

$$\begin{aligned}\sum_m c_m f_{nm} &= f c_n \\ \sum_m f_{nm} - f \delta_{nm} c_m &= 0 \Rightarrow \\ || f_{nm} - f \delta_{nm} || &= 0\end{aligned}\tag{1.29}$$

$$(1.30)$$

1.3 Switching to a different state basis

$\{\varphi_n(q)\}$ and $\{\varphi'_n(q)\}$ are two different basis's.

$$\varphi'_n(q) = \sum_m S_{mn} \varphi_n(q) \tag{1.31}$$

$$\varphi'_n = \hat{S} \varphi_n \tag{1.32}$$

Where \hat{S} is the transition operator. If the new basis $\{\varphi'_n(q)\}$ is orthogonal, meaning:

$$\int \varphi_m'^* \varphi'_n dq = \delta_{mn} \tag{1.33}$$

then $\hat{S}^\dagger = \hat{S}^{-1}$.

$$\begin{aligned}\int \varphi_m'^* \varphi'_n dq &= \delta_{mn} \\ \int \hat{S}^* \varphi_m^* \hat{S} \varphi_n dq &= \delta_{mn} \\ \int \varphi_m^* (\hat{S}^*)^T \hat{S} \varphi_n dq &= \delta_{mn} \\ \int \varphi_m^* \sum_l S_{ml}^* S_{ln} \varphi_n dq &= \delta_{mn} \\ \sum_l S_{ml}^* S_{ln} \int \varphi_m^* \varphi_n dq &= \delta_{mn} \\ \delta_{mn} (\sum_l S_{ml}^* S_{ln} - 1) &= 0 \Rightarrow \\ \hat{S}^\dagger &= \hat{S}^{-1} \quad (S_{mn}^\dagger = S_{nm}^*)\end{aligned}\tag{1.34}$$

Operators in the new basis can be written as:

$$\begin{aligned}\int \varphi_m'^* \hat{f} \varphi'_n dq &= \int (\hat{S}^* \varphi_m^*) (\hat{f} \hat{S} \varphi_n) dq = \\ &= \int \varphi_m^* \hat{S}^{*T} \hat{f} \hat{S} \varphi_n dq = \int \varphi_m^* \hat{f}' \varphi_n dq \Rightarrow \\ \hat{f}' &= \hat{S}^{*T} \hat{f} \hat{S} = \hat{S}^\dagger \hat{f} \hat{S} = \hat{S}^{-1} \hat{f} \hat{S}\end{aligned}\tag{1.35}$$

The operator's matrix's trace is the sum of the operator matrix's diagonal elements:

$$Sp\hat{f} = \sum_n f_{nn} \quad (1.36)$$

1. The trace of the product of two operators is invariant to the order of the operators

$$Sp(\hat{f}\hat{g}) = Sp(\hat{g}\hat{f}) \quad (1.37)$$

According to 1.36, $Sp\hat{f} = \sum_n f_{nn}$, therefore (using 1.24):

$$\begin{aligned} Sp(\hat{f}\hat{g}) &= \sum_n (\hat{f} \times \hat{g})_{nn} \\ &= \sum_n \sum_k f_{nk} g_{kn} \quad \text{and} \end{aligned} \quad (1.38)$$

$$\begin{aligned} Sp(\hat{g}\hat{f}) &= \sum_n (\hat{g} \times \hat{f})_{nn} \\ &= \sum_n \sum_k g_{nk} f_{kn}, \quad n \rightarrow k; k \rightarrow n \\ &= \sum_k \sum_n g_{kn} f_{nk} = \sum_n \sum_k f_{nk} g_{kn} \end{aligned} \quad (1.39)$$

2. The trace of the product of three or more operators is invariant to the cyclic permutation of the operators

$$Sp(\hat{f}\hat{g}\hat{h}) = Sp(\hat{h}\hat{f}\hat{g}) = Sp(\hat{g}\hat{h}\hat{f}) \quad (1.40)$$

According to 1.37

$$Sp(\hat{f}(\hat{g}\hat{h})) = Sp((\hat{g}\hat{h})\hat{f}) \quad \text{and} \quad (1.41)$$

$$Sp((\hat{f}\hat{g})\hat{h}) = Sp(\hat{h}(\hat{f}\hat{g})) \quad (1.42)$$

Which is equivalent to 1.40 because matrix multiplication is associative.

The operator's matrix's trace is invariant to the basis.

$$Sp\hat{f}' = Sp\hat{S}^{-1}\hat{f}\hat{S} = Sp\hat{S}\hat{S}^{-1}\hat{f} = Sp\hat{f} \quad (1.43)$$

Commutators Two operators commute if and only if they share a set of basis states. Or, in other words, there exist as basis in which they are both diagonal.

If \hat{f} and \hat{g} commute, then:

$$[\hat{f}, \hat{g}] = \hat{f}\hat{g} - \hat{g}\hat{f} = 0 \quad (1.44)$$

Which means that

$$\sum_k f_{mk} g_{kn} = \sum_k g_{mk} f_{kn} \quad (1.45)$$

If $\{\varphi_n\}$ are eigenfunctions of \hat{f} , then $f_{nm} \neq 0$ only if $n = m \rightarrow$

$$\begin{aligned} f_{mm} g_{mn} &= g_{mn} f_{nn} \\ g_{mn} (f_{mm} - f_{nn}) &= 0 \end{aligned} \quad (1.46)$$

Meaning that $g_{mn} = 0$ if $m \neq n$

1.4 Pauli uncertainty principle

The Pauli uncertainty principle states that if the operators of two observables do not commute, than we cannot measure both observables with arbitrary precision at the same time. In other words, that to more certain are we about one observable, the more uncertain we are about the other. For example:

$$[\hat{x}, \hat{p}_x] = i\hbar \quad (1.47)$$

$$\Delta x \Delta p_x \geq \hbar \quad (1.48)$$

or, for an exact relation,

$$\sigma_x \sigma_{p_x} \geq \frac{\hbar}{2} \quad (1.49)$$

$$\sigma_x = \sqrt{\langle \Delta x^2 \rangle} \quad \sigma_{p_x} = \sqrt{\langle \Delta p_x^2 \rangle} \quad (1.50)$$

$$(1.51)$$

Single-slit electron diffraction




Figure 1.1: Single-slit electron diffraction

$$b \sin(\theta) = \lambda \quad (1.52)$$

$$\Delta p_x = p \sin(\theta) \quad (1.53)$$

$$\Delta x = d \quad (1.54)$$

Using the relation for the de Broglie wavelength

$$\lambda = \frac{h}{p} = \frac{2 * \pi * \hbar}{p} \quad (1.55)$$

We get, considering

$$\Delta x \Delta p_x = d \sin(\theta) \frac{2\pi\hbar}{\lambda} = 2\pi\hbar \quad (1.56)$$

$$\Delta x \Delta p_x \approx \hbar \quad (1.57)$$

Black holes

And now let's talk about black holes

Ivan Iorsh

For an white dwarf star we can say that it's electrically neutral, has about the same number of protons as neutrons and is approximately spherical:

$$\bar{e} \rightarrow N \quad (1.58)$$

$$\bar{p} \rightarrow N \quad (1.59)$$

$$\bar{n}_0 \rightarrow N \quad (1.60)$$

$$V = \frac{4}{3}\pi R^3 \quad (1.61)$$

$$M_{\odot} = 2m_p N \quad (1.62)$$

Knowing the total volume we can estimate the average volume occupied by each electron, d_e :




Figure 1.2: Spherical horse in a vacuum

$$d_e = \left(\frac{\frac{4}{3}\pi R^3}{N} \right)^{\frac{1}{3}} = \frac{R}{N^{\frac{1}{3}}} \left(\frac{4}{3}\pi \right)^{\frac{1}{3}} \quad (1.63)$$

TODO: MOVE TO END: Average electron position, taking into account the fact that the electrons are confined:

$$\langle \Delta d^2 \rangle = \langle d^2 \rangle + \langle \Delta d \rangle^2 = \langle d^2 \rangle \quad (1.64)$$

And considering that electron positions d are of the same order as their **dispersion** $\Delta d \approx d$; and the same for their momentum, $\Delta p \approx p$ the Pauli uncertainty principle can be written as (for one electron):

$$\Delta p \Delta d \gtrsim \frac{\hbar}{2}; \quad v = \frac{p}{m_e} \quad (1.65)$$

$$\Delta p \approx \frac{\hbar}{2\Delta d} = \frac{\hbar}{2d} \quad (1.66)$$

$$E_{kinetic} = \frac{m_e v^2}{2} = \frac{p^2}{2m_e} = \frac{\hbar^2}{8d^2 m_e} \quad (1.67)$$

And for N electrons:

$$E_{kN} = \frac{N\hbar^2}{8d^2 m_e} \quad (1.68)$$

For an average star,

$$M_{\odot} \approx 2 * 10^{33} \text{g} \quad (1.69)$$

$$R_{\odot} \approx 6 * 10^5 \text{m} \quad (1.70)$$

$$m_p \approx 10^{-27} \text{kg} \quad (1.71)$$

$$m_e \approx 10^{-30} \text{kg} \quad (1.72)$$

$$\hbar = 1.05 * 10^{-34} \text{J} * \text{s} \quad (1.73)$$

The average speed of an electron in the star is **Explicit calc**

$$V = \frac{p}{m_e} = \frac{\hbar}{m_e} \frac{N^{\frac{1}{3}}}{R} \left(\frac{4}{3}\pi \right)^{-\frac{1}{3}} = \frac{\hbar}{m_e} \left(\frac{M}{2m_p} \right)^{\frac{1}{3}} \frac{1}{R} \left(\frac{4}{3}\pi \right)^{-\frac{1}{3}} = \quad (1.74)$$

$$= \dots \quad (1.75)$$

$$\approx 10^8 \frac{\text{m}}{\text{s}} \approx \frac{1}{3} c \quad (1.76)$$

Which is highly relativistic. /hlEnd Part to move

Energy of a star:

$$E_{full} = E_{kinetic} + U_{gravitational} \quad (1.77)$$

$$F = \frac{Gm_1 m_2}{r^2} - \frac{\partial d}{\partial r} \quad (1.78)$$

$$U = - \frac{Gm_1 m_2}{r} \quad (1.79)$$




Figure 1.3: Forces integration schematic

The potenetial energy of the star:

$$dU = \frac{-G(\frac{4}{3}\pi r^3 \rho)(4\pi r^2 dr \rho)}{r} = -\frac{16}{3}G\pi^2 \rho^2 r^4 \quad (1.80)$$

$$U = \int_0^R dU = - \int_0^R \frac{16}{3}G\pi^2 \rho^2 r^4 = -\frac{16}{15}G\pi^2 \rho^2 r^5 \Big|_0^R = -\frac{16}{15}G\pi^2 \rho^2 R^5 \quad (1.81)$$

$$M = \frac{4}{3}\pi \rho R^3 \quad (1.82)$$

$$\frac{M^2}{R} = \frac{16}{9}\pi^2 \rho^2 R^5 \quad (1.83)$$

$$U = -G \frac{9}{15} \frac{M^2}{R} \quad (1.84)$$

The kinetic energy of the star's electrons: For a single electron:

$$\Delta p \Delta d \gtrsim \frac{\hbar}{2}; \quad v \frac{p}{m_e} \quad (1.85)$$

$$\Delta p \approx \frac{\hbar}{2\Delta d} = \frac{\hbar}{2d} \quad (1.86)$$

$$E_{kinetic} = \frac{m_e v^2}{2} = \frac{p^2}{2m_e} = \frac{\hbar^2}{8d^2 m_e} \quad (1.87)$$

And for N electrons:

$$E_{kN} = \frac{N\hbar^2}{8d^2 m_e} \quad (1.88)$$

Now the total energy of the star is:

$$E = \frac{N\hbar^2}{8d^2m_e} - G \frac{9}{15} \frac{M^2}{R} = \quad (1.89)$$

$$= \frac{N^{\frac{5}{3}}\hbar^2}{8R^2m_e(\frac{4}{3}\pi)^{\frac{2}{3}}} - G \frac{9}{15} \frac{M^2}{R} = \quad (1.90)$$

$$= \frac{M^{\frac{5}{3}}\hbar^2}{8R^2m_e(\frac{4}{3}\pi)^{\frac{2}{3}}(2m_p)^{\frac{5}{3}}} - G \frac{9}{15} \frac{M^2}{R} \quad (1.91)$$

For a stable star, its stable radius should be in a minimum of energy,

$$\frac{\partial E}{\partial R} = 0 \quad (1.92)$$

For our star,

$$\frac{\partial E}{\partial R} = - \frac{M^{\frac{5}{3}}\hbar^2}{4R^3m_e(\frac{4}{3}\pi)^{\frac{2}{3}}(2m_p)^{\frac{5}{3}}} + G \frac{9}{15} \frac{M^2}{R^2} \quad (1.93)$$

$$\frac{\partial E}{\partial R} = 0 \Rightarrow \quad (1.94)$$

$$\frac{M^{\frac{5}{3}}\hbar^2}{4R^3m_e(\frac{4}{3}\pi)^{\frac{2}{3}}(2m_p)^{\frac{5}{3}}} = G \frac{9}{15} \frac{M^2}{R^2} \quad (1.95)$$

$$\frac{15}{36} \frac{GM^{-\frac{1}{3}}}{m_e(\frac{4}{3}\pi)^{\frac{2}{3}}(2m_p)^{\frac{5}{3}}} = R \quad (1.96)$$

Or simply,

$$M^{\frac{1}{3}}R = \text{const} \quad (1.97)$$

Which means that for every stellar mass there exists a certain stable radius. The problem with this equation is that it does not take into account that electrons in such a star are highly relativistic **TODO: MOVE RELATIVISTIC THING HERE**, meaning that the kinetic energy of the star cannot be accurately represented as in 1.88.

For a relativistic electron,

$$E_k = \sqrt{m^2c^4 + p^2c^2} \approx \quad (1.98)$$

$$pc \ll mc^2, \quad \approx mc^2 + \frac{p^2}{2m} \quad (1.99)$$

$$pc \gg mc^2, \quad \approx cp \quad (1.100)$$

Taking into account that $v_e \approx \frac{1}{3}c$, and that $p = \frac{\hbar}{2d}$,

$$E_{kinetic} = \frac{c\hbar}{2d} = \frac{c\hbar N^{\frac{4}{3}}}{2R(\frac{4}{3}\pi)^{\frac{1}{3}}} = \frac{c\hbar M^{\frac{4}{3}}}{2R(\frac{4}{3}\pi)^{\frac{1}{3}}(2m_p)^{\frac{4}{3}}} \quad (1.101)$$

$$E = \frac{c\hbar M^{\frac{4}{3}}}{2R(\frac{4}{3}\pi)^{\frac{1}{3}}(2m_p)^{\frac{4}{3}}} - G \frac{9}{15} \frac{M^2}{R} = \quad (1.102)$$

$$= \frac{1}{R} \left(\frac{c\hbar M^{\frac{4}{3}}}{2(\frac{4}{3}\pi)^{\frac{1}{3}}(2m_p)^{\frac{4}{3}}} - GM^2 \frac{9}{15} \right) \quad (1.103)$$

Which means that a stable R doesn't exist for such stars - if the expression in parenthesis in 1.103 is greater than zero, than the star expands until its electrons are no longer relativistic, and it settles to a radius defined by 1.97, or if the expression in parenthesis in 1.103 is less than zero, the star's kinetic energy is insufficient to withstand its gravitational pull and it collapses into a black hole. The mass at which this happens, M_{cr} , is

$$M_{cr} = \left(\frac{15c\hbar}{9G2(2m_p)^{\frac{4}{3}}(\frac{4}{3}\pi)^{\frac{1}{3}}} \right)^{\frac{3}{2}} = \quad (1.104)$$

...explicit calc...

$$\approx 10^{30} \text{ kg} \approx 1.4M_{\odot} \quad (1.105)$$

Where M_{\odot} is the mass of our sun. Meaning that no white dwarf star with a mass over $1.4M_{\odot}$ can stably exist. Nobel Prize lecture, Subrahmanyan Chandrasekhar[2].

Quantum Pencil




Figure 1.4: Vertical pencil diagram

If we place a pencil with mass $m = 10^{-6}\text{kg}$ on its tip $d_0 = 10^{-10}\text{m}$, because of the uncertainty principle, its center of mass will start to move,

$$\Delta x \Delta p \approx \frac{\hbar}{2} \quad (1.106)$$

$$\Delta x < d_0 \quad (1.107)$$

$$\Delta p > \frac{\hbar}{2d_0} \quad (1.108)$$

$$p \sim \Delta p \quad (1.109)$$

$$v = \frac{p}{m} = \frac{\hbar}{2d_0 m} \quad (1.110)$$

We can say that the pencil has fallen when its center of mass is no longer over the "tip" of the pencil (Fig., 1.4).

$$t \sim \frac{d_0}{v} = \frac{2d_0^2 m}{\hbar} \approx \frac{2 * (10^{-10})^2 * 10^{-6}}{10^{-34}} \approx 10^8 \text{s} \quad (1.111)$$

Which means that the pencil can stably exist in a vertical position for over 3 years (compared to other solutions, which give unrealistic estimates of about 3 seconds)[3].

1.5 Problems

2 Analytical Solutions

2.1 Rectangular quantum well

2.2 Harmonic oscillator

2.3 Spherically symmetric potential

2.4 Problems

Rectangular quantum well

Double quantum well

Double quantum barrier

Minimal transistor size

3 Quasi-classical approximation

3.1 Problems

4 Spin

4.1 Angular momentum

Classical

Figure 4.1: Forces integration schematic

Quantum

4.2 Problems

5 Perturbation theory

5.1 Time-independent

5.2 Time-dependent

5.3 Problems

6 Problem Solutions

6.1 Introduction

6.2 Analytical solutions

6.3 Quasiclassical approximation

6.4 Spin

6.5 Perturbation theory

References

- [1] J. J. Sakurai and J. J. Napolitano, Modern quantum mechanics. Pearson Higher Ed, 2014.
- [2] S. Chandrasekhar, “On stars, their evolution and their stability,” Reviews of modern physics, vol. 56, no. 2, p. 137, 1984.
- [3] D. Easton, “The quantum mechanical tipping pencil – a caution for physics teachers,” European Journal of Physics, vol. 28, no. 6, p. 1097, 2007.