


Chapter 2. Getting to Know Your Data

- ❑ Data Objects and Attribute Types
- ❑ Basic Statistical Descriptions of Data
- ❑ Data Visualization
- ❑ Measuring Data Similarity and Dissimilarity
- ❑ Summary




Similarity, Dissimilarity, and Proximity

- ❑ **Similarity measure or similarity function** គេស្រាវជ្រាវដើម្បីវាស់ កម្រិតនៃការស្របគ្នា
 - ❑ A real-valued function that quantifies the similarity between two objects
 - ❑ Measure how two data objects are alike: The higher value, the more alike
 - ❑ Often falls in the range $[0,1]$: 0: no similarity; 1: completely similar
0: គ្មានការស្របគ្នា; 1: ស្របគ្នា
- ❑ **Dissimilarity (or distance) measure** វាស់ពីការខុសគ្នា
 - ❑ Numerical measure of how different two data objects are វាស់ពីការខុសគ្នា រវាង 2 វត្ថុទិន្នន័យ
 - ❑ In some sense, the inverse of similarity: The lower, the more alike
 - ❑ Minimum dissimilarity is often 0 (i.e., completely similar) $[0,1]$; ស្របគ្នា
 - ❑ Range $[0, 1]$ or $[0, \infty)$, depending on the definition $[0, \infty]$; ចម្ងាយ
- ❑ **Proximity** usually refers to either similarity or dissimilarity

Data Matrix and Dissimilarity Matrix

- Data matrix

- A data matrix of n data points with l dimensions


$$D = \begin{pmatrix} x_{11} & x_{12} & \dots & x_{1l} \\ x_{21} & x_{22} & \dots & x_{2l} \\ \vdots & \vdots & \ddots & \vdots \\ x_{n1} & x_{n2} & \dots & x_{nl} \end{pmatrix}$$


- Dissimilarity (distance) matrix

- n data points, but registers only the distance $d(i, j)$ (typically metric)

- Usually symmetric, thus a triangular matrix

- **Distance functions** are usually different for real, boolean, categorical, ordinal, ratio, and vector variables

- Weights can be associated with different variables based on applications and data semantics


$$\begin{pmatrix} 0 & & & \\ d(2,1) & 0 & & \\ \vdots & \vdots & \ddots & \\ d(n,1) & d(n,2) & \dots & 0 \end{pmatrix}$$

for analyzing unstructured data

Standardizing Numeric Data

- Z-score:
$$z = \frac{x - \mu}{\sigma}$$
 - X: raw score to be standardized, μ : mean of the population, σ : standard deviation
 - the distance between the raw score and the population mean in units of the standard deviation
 - negative when the raw score is below the mean, “+” when above
- An alternative way: Calculate the mean absolute deviation

$$s_f = \frac{1}{n} (|x_{1f} - m_f| + |x_{2f} - m_f| + \dots + |x_{nf} - m_f|)$$


where

$$m_f = \frac{1}{n} (x_{1f} + x_{2f} + \dots + x_{nf}).$$

- standardized measure (z-score):
$$z_{if} = \frac{x_{if} - m_f}{s_f}$$

standardizing column 1 into standard scale
kinnu 222 + 10msl 10000 column 10000
- Using mean absolute deviation is more robust than using standard deviation

Example: Data Matrix and Dissimilarity Matrix



Data Matrix

point	attribute1	attribute2
$x1$	1	2
$x2$	3	5
$x3$	2	0
$x4$	4	5

Dissimilarity Matrix (by **Euclidean Distance**)

	$x1$	$x2$	$x3$	$x4$
$x1$	0			
$x2$	3.61	0		
$x3$	2.24	5.1	0	
$x4$	4.24	1	5.39	0

Distance on Numeric Data: Minkowski Distance

- **Minkowski distance**: A popular distance measure

$$d(i, j) = \sqrt[p]{|x_{i1} - x_{j1}|^p + |x_{i2} - x_{j2}|^p + \dots + |x_{il} - x_{jl}|^p}$$

where $i = (x_{i1}, x_{i2}, \dots, x_{il})$ and $j = (x_{j1}, x_{j2}, \dots, x_{jl})$ are two l -dimensional data objects, and p is the order (the distance so defined is also called L- p norm)

- Properties

- $d(i, j) > 0$ if $i \neq j$, and $d(i, i) = 0$ (Positivity)

- $d(i, j) = d(j, i)$ (Symmetry)

- $d(i, j) \leq d(i, k) + d(k, j)$ (Triangle Inequality)

- A distance that satisfies these properties is a **metric**

- Note: There are nonmetric dissimilarities, e.g., set differences

Special Cases of Minkowski Distance

□ $p = 1$: (L_1 norm) **Manhattan (or city block) distance**

□ E.g., the Hamming distance: the number of bits that are different between two binary vectors

$$d(i, j) = |x_{i1} - x_{j1}| + |x_{i2} - x_{j2}| + \cdots + |x_{il} - x_{jl}|$$

□ $p = 2$: (L_2 norm) **Euclidean distance**

$$d(i, j) = \sqrt{|x_{i1} - x_{j1}|^2 + |x_{i2} - x_{j2}|^2 + \cdots + |x_{il} - x_{jl}|^2}$$

□ $p \rightarrow \infty$: (L_{\max} norm, L_{∞} norm) **“supremum” distance**


□ The maximum difference between any component (attribute) of the vectors

$$d(i, j) = \lim_{p \rightarrow \infty} \sqrt[p]{|x_{i1} - x_{j1}|^p + |x_{i2} - x_{j2}|^p + \cdots + |x_{il} - x_{jl}|^p} = \max_{f=1}^l |x_{if} - x_{jf}|$$

අනන්ත වෙනසක් ඇති දත්ත

Example: Minkowski Distance at Special Cases

point	attribute 1	attribute 2
x1	1	2
x2	3	5
x3	2	0
x4	4	5



Manhattan (L_1)

L	x1	x2	x3	x4
x1	0			
x2	5	0		
x3	3	6	0	
x4	6	1	7	0


Euclidean (L_2)

L2	x1	x2	x3	x4
x1	0			
x2	3.61	0		
x3	2.24	5.1	0	
x4	4.24	1	5.39	0

Supremum (L_∞)

L_∞	x1	x2	x3	x4
x1	0			
x2	3	0		
x3	2	5	0	
x4	3	1	5	0

point	attribute 1	attribute 2
x1	1	2
x2	3	5
x3	2	0
x4	4	5




$$L_1 \text{ norm} = 6 = 3 + 3$$

$$L_2 \text{ norm} = \sqrt{18} = \sqrt{3^2 + 3^2}$$

$$L_\infty = 3 = \text{ระยะห่างมากที่สุด}$$

point	attribute 1	attribute 2
x1	1	2
x2	3	5
x3	2	0
x4	4	5




$$L_1 \text{ norm} = 7 = 5 + 2$$

$$L_2 \text{ norm} = \sqrt{29} = \sqrt{5^2 + 2^2}$$

$$L_\infty = 5 = \max$$

point	attribute 1	attribute 2
x1	1	2
x2	3	5
x3	2	0
x4	4	5



ទំហំ: ប្រភេទ: ឆ្នាំ: ឆ្នាំ: A + ឆ្នាំ B

ទំហំ: ប្រភេទ: ឆ្នាំ: ឆ្នាំ: C ឆ្នាំ