

5 POSTOPERATIVE RESOURCE UTILIZATION AND SURVIVAL AMONG LIVER TRANSPLANT RECIPIENTS WITH A MELD SCORE GREATER THAN OR EQUAL TO 40: A RETROSPECTIVE COHORT STUDY

Filipe S. Cardoso, Constantine J. Karvellas, Norman Kneteman, Glenda Meeberg, Pedro Fidalgo, Sean M. Bagshaw

Introduction:

Cirrhotic patients with Model for End-stage Liver Disease (MELD) score ≥ 40 have high risk of death without liver transplant (LT). This study aimed to evaluate these patients' outcomes after transplant.

Methods:

The retrospective cohort included 519 adult cirrhotic patients who underwent LT at one Canadian center between 2002 and 2012. Primary exposure was severity of end-stage liver disease measured by MELD score at transplant (≥ 40 vs. < 40). Primary outcome was duration of first intensive care unit (ICU) stay after LT. Secondary outcomes were duration of first hospital stay after LT, rate of ICU readmission, re-transplant rate, and survival rates.

Results:

On the day of LT, 5% (28/519) of patients had a MELD score ≥ 40 . These patients had longer first ICU stay after LT (14 vs. 2 days; $p < 0.001$). MELD score ≥ 40 at transplant was independently associated with first ICU stay after transplant ≥ 10 days (OR, 3.21). These patients had longer first hospital stay after LT (45 vs. 18 days; $p < 0.001$); however, there was no significant difference in the rate of ICU readmission (18% vs. 22%; $p = 0.58$) or re-transplant rate (4% vs. 4%; $p = 1.00$). Cumulative survival at 1 month, 3 months, 1 year, 3 years, and 5 years was 98%, 96%, 90%, 79%, and 72%, respectively. There was no significant difference in cumulative survival stratified by MELD score ≥ 40 vs. < 40 at transplant ($p = 0.59$).

Conclusions:

Cirrhotic patients with MELD score ≥ 40 at transplant utilize greater postoperative health resources; however, derive similar long-term survival benefit with LT.

References:

Shawcross DL, Austin MJ, Abeles RD et al. The impact of organ dysfunction in cirrhosis: survival at a cost? *J Hepatol* 2012; 56(5):1054-62.
Alexopoulos S, Matsuoka L, Cho Y et al. Outcomes after liver transplantation in patients achieving a model for

end-stage liver disease score of 40 or higher. Transplantation 2013; 95(3):507-12.

Oberkofler CE, Dutkowski P, Stocker R et al. Model of end stage liver disease (MELD) score greater than 23

predicts length of stay in the ICU but not mortality in liver transplant recipients. Crit Care 2010; 14(3):R117.

Univeristy of Alberta Hospital, University of Alberta, Edmonton, Alberta, Canada