

International Journal of Computer Mathematics

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gcom20>

A new fifth order weighted runge kutta formula

D.J. Evans^a & A.R. Yaakub^a

^a Parallel Algorithms Research Centre, Loughborough University of Technology, Loughborough, Leicestershire, LE11 3TU, UK

Version of record first published: 19 Mar 2007.

To cite this article: D.J. Evans & A.R. Yaakub (1996): A new fifth order weighted runge kutta formula, International Journal of Computer Mathematics, 59:3-4, 227-243

To link to this article: <http://dx.doi.org/10.1080/00207169608804467>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

A NEW FIFTH ORDER WEIGHTED RUNGE KUTTA FORMULA

D. J. EVANS and A. R. YAAKUB

*Parallel Algorithms Research Centre,
 Loughborough University of Technology,
 Loughborough, Leicestershire, LE11 3TU, UK*

(Received August 9 1995)

In this paper a new weighted Runge Kutta fifth order method with 5 stages is introduced. By comparison with the currently recommended RK4(5) Merson and RK5(6) Nystrom methods, the new method gives improved results.

KEY WORDS: Runge Kutta, Merson, Nystrom, weighted mean.

C. R. CATEGORIES: G.1.7.

1 INTRODUCTION

From recent publications (see Evans & Yaakub [1993] and Evans & Sanugi [1987], [1993]), fourth order linear and non-linear methods using a variety of means for solving initial value problems of the form $y' = f(x, y)$ have been shown to have the form

$$y_{n+1} = y_n + \frac{h}{3} \left[\sum_{i=1}^3 \text{Means} \right], \quad (1.1)$$

where *Means* = principal means include arithmetic mean (AM), geometric mean (GM) [5], contraharmonic mean (C_0M) [2], centroidal mean (C_eM) [3], root mean square (RMS) [11], harmonic mean (H_aM) [10] and heronian mean (H_eM) [4] which involve k_i , $1 \leq i \leq 4$

where

$$\begin{aligned} k_1 &= f(x_n, y_n) \\ k_2 &= f(x_n + a_1 h, y_n + a_1 h k_1) \\ k_3 &= f(x_n + (a_2 + a_3) h, y_n + a_2 h k_1 + a_3 h k_2) \\ k_4 &= f(x_n + (a_4 + a_5 + a_6) h, y_n + a_4 h k_1 + a_5 h k_2 + a_6 h k_3) \end{aligned} \quad (1.2)$$

From the above discussion, a comparison of the parameters a_i , $1 \leq i \leq 6$ in equation (1.2), show that a_1 is fixed, a_2, a_4, a_6 are decreasing and a_3, a_5 as increasing as shown in Table 1.

From Table 1, we can see that $a_2 + a_3 = 1/2$ and $a_4 + a_5 + a_6 = 1$. In this paper, our concern is to establish a new fifth order formula based on the last two k values i.e., k_3 , and k_4 , which involve the parameters $a_2 + a_3 = 1/2$ and $a_4 + a_5 + a_6 = 1$ and weighted so we establish a new weighted Runge-Kutta formula (WRK) similar to [12].

2 THE FOURTH ORDER ARITHMETIC MEAN WEIGHTED RUNGE-KUTTA FORMULA

The standard fourth order arithmetic mean (AM) Runge-Kutta formula for solving IVPs may be written in the form

$$y_{n+1} = y_n + \frac{h}{3} \left[\sum_{i=1}^3 \left(\frac{k_i + k_{i+1}}{2} \right) \right] \quad (2.1)$$

or

$$y_{n+1} = y_n + \frac{h}{3} \left[\frac{k_1 + k_2}{2} + \frac{k_2 + k_3}{2} + \frac{k_3 + k_4}{2} \right]$$

where k_i , $1 \leq i \leq 4$ as we mentioned above.

Now, based on the parameters involved in the last two k values i.e., k_3 and k_4 , we attempt to develop a new fourth order method called the weighted Runge-Kutta (WRK) formula in the form

$$y_{n+1} = y_n + h \left[\sum_{i=1}^3 w_i \left(\frac{k_i + k_{i+1}}{2} \right) \right] \quad (2.2)$$

Table 1 The values of parameters a_i , $1 \leq i \leq 6$ based on the various integration formulas.

| | C_0M | C_eM | RMS | AM | GM | H_eM | H_aM |
|-------|----------------|-------------------|------------------|---------------|-----------------|-------------------|----------------|
| a_1 | $\frac{1}{2}$ | $\frac{1}{2}$ | $\frac{1}{2}$ | $\frac{1}{2}$ | $\frac{1}{2}$ | $\frac{1}{2}$ | $\frac{1}{2}$ |
| a_2 | $\frac{1}{8}$ | $\frac{1}{24}$ | $\frac{1}{16}$ | 0 | $-\frac{1}{16}$ | $-\frac{1}{48}$ | $-\frac{1}{8}$ |
| a_3 | $\frac{3}{8}$ | $\frac{11}{24}$ | $\frac{7}{16}$ | $\frac{1}{2}$ | $\frac{9}{16}$ | $\frac{25}{48}$ | $\frac{5}{8}$ |
| a_4 | $\frac{1}{4}$ | $\frac{1}{12}$ | $\frac{1}{8}$ | 0 | $-\frac{1}{8}$ | $-\frac{1}{24}$ | $-\frac{1}{4}$ |
| a_5 | $-\frac{3}{4}$ | $-\frac{132}{25}$ | $-\frac{56}{17}$ | 0 | $\frac{24}{5}$ | $\frac{600}{47}$ | $\frac{20}{7}$ |
| a_6 | $\frac{3}{2}$ | $\frac{73}{66}$ | $\frac{33}{28}$ | 1 | $\frac{11}{12}$ | $\frac{289}{300}$ | $\frac{9}{10}$ |

where $\sum_{i=1}^3 w_i = 1$ and $k_1 = f(y_n)$

$$\begin{aligned}k_2 &= f(y_n + a_1 h k_1) \\k_3 &= f(y_n + a_2 h k_1 + (\tfrac{1}{2} - a_2) h k_2) \\k_4 &= f(y_n + a_3 h k_1 + a_4 h k_2 + (1 - a_3 - a_4) h k_3).\end{aligned}\quad (2.3)$$

By use of the standard procedure of adjustment of the parameters, a fourth order accuracy formula is obtained from solving the 7 non-linear equations involving 7 variables obtained by comparison with terms of specified order in the Taylor series of expansion, i.e.,

$$hf: \quad 1. \quad -x(1) - x(2) - x(3) = 0, \quad (2.4-i)$$

$$h^2 ff_y: \quad 2. \quad -2.*x(4)*x(1) - x(2) - 2.*x(4)*x(2) - 3.*x(3) = 0, \quad (2.4-ii)$$

$$\begin{aligned}h^3 f f_y^2: \quad 2. \quad &-3.*x(4)*x(2) + 6.*x(4)*x(5)*x(2) - 3.*x(3) \\&- 3.*x(4)*x(3) + 6.*x(4)*x(5)*x(3) + 3.*x(6)*x(3) \\&+ 3.*x(7)*x(3) - 6.*x(4)*x(7)*x(3) = 0,\end{aligned}\quad (2.4-iii)$$

$$\begin{aligned}h^3 f^2 f_{yy}: \quad 8. \quad &-12.*x(4)**2*x(1) - 3.*x(2) - 12.*x(4)**2*x(2) \\&- 15.*x(3) = 0,\end{aligned}\quad (2.4-iv)$$

$$\begin{aligned}h^4 f f_y^3: \quad 1. \quad &-6.*x(4)*x(3) + 12.*x(4)*x(5)*x(3) \\&- 12.*x(4)*x(5)*x(6)*x(3) + 6.*x(4)*x(7)*x(3) \\&- 12.*x(4)*x(5)*x(7)*x(3) + 6.*x(4)*x(6)*x(3) = 0,\end{aligned}\quad (2.4-v)$$

$$\begin{aligned}h^4 f^2 f_y f_{yy}: \quad 8. \quad &-6.*x(4)*x(2) - 6.*x(4)**2*x(2) \\&+ 12.*x(4)*x(5)*x(2) + 12.*x(4)**2*x(5)*x(2) - 15.*x(3) \\&- 6.*x(4)*x(3) - 6.*x(4)**2*x(3) + 12.*x(4)*x(5)*x(3) \\&+ 12.*x(4)**2*x(5)*x(3) + 15.*x(6)*x(3) + 15.*x(7)*x(3) \\&- 24.*x(4)*x(7)*x(3) - 12.*x(4)**2*x(7)*x(3) = 0,\end{aligned}\quad (2.4-vi)$$

$$h^4 f^3 f_{yyy}: \quad 4. \quad -8.*x(4)**3*x(1) - x(2) - 9.*x(3) - 8.*x(4)**3*x(2) = 0 \quad (2.4-vii)$$

where $x(1) = w_1$, $x(2) = w_2$, $x(3) = w_3$, $x(4) = a_1$, $x(5) = a_2$, $x(6) = a_3$ and $x(7) = a_4$.

Equations (2.4-i)–(2.4-vii) are then solved simultaneously using the NAG routine (Subroutine C05NBF) for solving a system of non-linear equations to give the

required parameters, i.e.,

$$x(1) = w_1 = 0.3333333333, \quad x(2) = w_2 = 0.3333333333, \quad x(3) = w_3 = 0.3333333333$$

$$x(4) = a_1 = 0.5000000000, \quad x(5) = a_2 = 0, \quad x(6) = a_3 = 0, \quad x(7) = a_4 = 0$$

Thus, the WRK method gives the same result as the standard fourth order arithmetic mean (AM) Runge-Kutta method i.e.,

$$y_{n+1} = y_n + \frac{h}{3} \left[\frac{k_1 + k_2}{2} + \frac{k_2 + k_3}{2} + \frac{k_3 + k_4}{2} \right]$$

where

$$k_1 = f(y_n)$$

$$k_2 = f\left(y_n + \frac{1}{2}hk_1\right)$$

$$k_3 = f\left(y_n + \frac{1}{2}hk_2\right)$$

$$k_4 = f(y_n + hk_3).$$

3 A NEW FIFTH ORDER ARITHMETIC MEAN WEIGHTED RUNGE- KUTTA FORMULA

We now extend the same procedure to obtain the fifth order formula in the form

$$y_{n+1} = y_n + h \left[\sum_{i=1}^4 w_i \left(\frac{k_i + k_{i+1}}{2} \right) \right] \quad (3.1)$$

where

$$\sum_{i=1}^4 w_i = 1$$

and

$$k_1 = f(y_n)$$

$$k_2 = f(y_n + a_1hk_1)$$

$$k_3 = f(y_n + a_2hk_1 + a_3hk_2)$$

$$k_4 = f(y_n + a_4hk_1 + a_5hk_2 + (\frac{1}{2} - a_4 - a_5)hk_3)$$

$$k_5 = f(y_n + a_6hk_1 + a_7hk_2 + a_8hk_3 + (1 - a_6 - a_7 - a_8)hk_4). \quad (3.2)$$

The Taylor series expansion of $y(x_{n+h})$ up to sixth order is given by

$$\begin{aligned}
 y(x_n + h) = & y_n + hf + \frac{1}{2} h^2 ff_y + \frac{1}{6} h^3 (ff_y^2 + f^2 f_{yy}) \\
 & + \frac{1}{24} h^4 (f^3 f_{yyy} + 4f^2 f_y f_{yy} + ff_y^3) \\
 & + \frac{1}{120} h^5 (ff_y^4 + 11f^2 f_y^2 f_{yy} + 4f^3 f_{yy}^2 + 7f^3 f_y f_{yyy} + f^4 f_{yyyy}) \\
 & + \frac{1}{720} h^6 (f^5 f_{yyyy} + 11f^4 f_y f_{yyy} + 15f^4 f_{yy} f_{yyy} + 32f^3 f_y^2 f_{yyy} \\
 & + 34f^3 f_y f_{yy}^2 + 26f^2 f_y^3 f_{yy}^2 + ff_y^5) + O(h^7) \dots
 \end{aligned} \quad (3.3)$$

By substituting equation (3.2) into (3.1) and subtract from equation (3.3), we obtain 12 equations with 12 parameters, i.e.,

$$hf: \quad 1 - x(1) - x(2) - x(3) - x(4) = 0, \quad (3.4-i)$$

$$\begin{aligned}
 h^2 ff_y: \quad & 2 - 2*x(5)*x(1) - 2*x(5)*x(2) - 2*x(6)*x(2) \\
 & - 2*x(7)*x(2) - x(3) - 2*x(6)*x(3) - 2*x(7)*x(3) - 3*x(4) = 0, \quad (3.4-ii)
 \end{aligned}$$

$$\begin{aligned}
 h^3 ff_y^2: \quad & 2 - 6*x(5)*x(7)*x(2) - 3*x(6)*x(3) - 3*x(7)*x(3) - 6*x(5)*x(7)*x(3) \\
 & + 6*x(6)*x(8)*x(3) + 6*x(7)*x(8)*x(3) - 6*x(5)*x(9)*x(3) \\
 & + 6*x(6)*x(9)*x(3) \\
 & + 6*x(6)*x((7)*x(3) - 3*x(4) - 3*x(6)*x(4) - 3*x(7)*x(4) \\
 & + 6*x(6)*x(8)*x(4) + 6*x(7)*x(8)*x(4) - 6*x(5)*x(9)*x(4) \\
 & + 6*x(6)*x(9)*x(4) + 6*x(7)*x(9)*x(4) \\
 & + 3*x(10)*x(4) + 3*x(11)*x(4) - 6*x(5)*x(11)*x(4) \\
 & + 3*x(12)*x(4) - 6*x(6)*x(12)*x(4) - 6*x(7)*x(12)*x(4) = 0, \quad (3.4-iii)
 \end{aligned}$$

$$\begin{aligned}
 h^3 f^2 f_{yy}: \quad & 8 - 12*x(5)**2*x(1) - 12*x(5)**2*x(2) - 12*x(6)**2*x(2) \\
 & - 24*x(6)*x(7)*x(2) - 12*x(7)**2*x(2) - 3*x(3) \\
 & - 12*x(6)**2*x(3) - 24*x(6)*x(7)*x(3) - 12*x(7)**2*x(3) - 15*x(4) = 0, \\
 & \dots \quad (3.4-iv)
 \end{aligned}$$

$$\begin{aligned}
 h^4 ff_y^3: \quad & 1 - 6*x(5)*x(7)*x(3) + 12*x(5)*x(7)*x(8)*x(3) \\
 & + 12*x(5)*x(7)*x(9)*x(3) - 6*x(6)*x(4) - 6*x(7)*x(4) \\
 & - 6*x(5)*x(7)*x(4) + 12*x(6)*x(8)*x(4) + 12*x(7)*x(8)*x(4)
 \end{aligned}$$

$$\begin{aligned}
& + 12*x(5)*x(7)*x(8)*x(4) - 12*x(5)*x(9)*x(4) \\
& + 12*x(6)*x(9)*x(4) + 12*x(7)*x(9)*x(4) + 12*x(5)*x(7)*x(9)*x(4) \\
& + 6*x(6)*x(10)*x(4) + 6*x(7)*x(10)*x(4) \\
& - 12*x(6)*x(8)*x(10)*x(4) - 12*x(7)*x(8)*x(10)*x(4) \\
& + 12*x(5)*x(9)*x(10)*x(4) - 12*x(6)*x(9)*x(10)*x(4) \\
& - 12*x(7)*x(9)*x(10)*x(4) + 6*x(6)*x(11)*x(4) \\
& + 6*x(7)*x(11)*x(4) - 12*x(6)*x(8)*x(11)*x(4) \\
& - 12*x(7)*x(8)*x(11)*x(4) + 12*x(5)*x(9)*x(11)*x(4) \\
& - 12*x(6)*x(9)*x(11)*x(4) - 12*x(7)*x(9)*x(11)*x(4) \\
& + 6*x(6)*x(12)*x(4) + 6*x(7)*x(12)*x(4) \\
& - 12*x(5)*x(7)*x(12)*x(4) - 12*x(6)*x(8)*x(12)*x(4) \\
& - 12*x(7)*x(8)*x(12)*x(4) + 12*x(5)*x(9)*x(12)*x(4) \\
& - 12*x(6)*x(9)*x(12)*x(4) - 12*x(7)*x(9)*x(12)*x(4) = 0, \quad (3.4-v)
\end{aligned}$$

$$\begin{aligned}
h^4 f^2 f_y f_{yy}: \quad & 8 - 12*x(5)**2*x(7)*x(2) - 24*x(5)*x(6)*x(7)*x(2) \\
& - 24*x(5)*x(7)**2*x(2) - 6*x(6)*x(3) - 6*x(6)**2*x(3) \\
& - 6*x(7)*x(3) - 12*x(5)**2*x(7)*x(3) - 12*x(6)*x(7)*x(3) \\
& - 24*x(5)*x(6)*x(7)*x(3) - 6*x(7)**2*x(3) \\
& - 24*x(5)*x(7)**2*x(3) + 12*x(6)*x(8)*x(3) \\
& + 12*x(6)**2*x(8)*x(3) + 12*x(7)*x(8)*x(3) \\
& + 24*x(6)*x(7)*x(8)*x(3) + 12*x(7)**2*x(8)*x(3) \\
& - 12*x(5)*x(9)*x(3) - 12*x(5)**2*x(9)*x(3) \\
& + 12*x(6)*x(9)*x(3) + 12*x(6)**2*x(9)*x(3) + 12*x(7)*x(9)*x(3) \\
& + 24*x(6)*x(7)*x(9)*x(3) \\
& + 12*x(7)**2*x(9)*x(3) - 15*x(4) - 6*x(6)*x(4) \\
& - 6*x(6)**2*x(4) - 6*x(7)*x(4) - 12*x(6)*x(7)*x(4) \\
& - 6*x(7)**2*x(4) + 12*x(6)*x(8)*x(4) + 12*x(6)**2*x(8)*x(4) \\
& + 12*x(7)*x(8)*x(4) + 24*x(6)*x(7)*x(8)*x(4) \\
& + 12*x(7)**2*x(8)*x(4) - 12*x(5)*x(9)*x(4) \\
& - 12*x(5)**2*x(9)*x(4) + 12*x(6)*x(9)*x(4) \\
& + 12*x(6)**2*x(9)*x(4) + 12*x(7)*x(9)*x(4)
\end{aligned}$$

$$\begin{aligned}
& + 24*x(6)*x(7)*x(9)*x(4) + 12*x(7)**2*x(9)*x(4) + 15*x(10)*x(4) \\
& + 15*x(11)*x(4) - 24*x(5)*x(11)*x(4) - 12*x(5)**2*x(11)*x(4) \\
& + 15*x(12)*x(4) - 24*x(6)*x(12)*x(4) \\
& - 12*x(6)**2*x(12)*x(4) - 24*x(7)*x(12)*x(4) \\
& - 24*x(6)*x(7)*x(12)*x(4) - 12*x(7)**2*x(12)*x(4) = 0, \quad (3.4-vi)
\end{aligned}$$

$$\begin{aligned}
h^4 f^3 f_{yyy}: & 4 - 8*x(5)**3*x(1) - 8*x(5)**3*x(2) - 8*x(6)**3*x(2) \\
& - 24*x(6)**2*x(7)*x(2) - 24*x(6)*x(7)**2*x(2) \\
& - 8*x(7)**3*x(2) - x(3) - 8*x(6)**3*x(3) - 24*x(6)**2*x(7)*x(3) \\
& - 24*x(6)*x(7)**2*x(3) - 8*x(7)**3*x(3) - 9*x(4) = 0, \quad (3.4-vii)
\end{aligned}$$

$$\begin{aligned}
h^5 ff^4: & 1 - 30*x(5)*x(7)*x(4) + 60*x(5)*x(7)*x(8)*x(4) \\
& + 60*x(5)*x(7)*x(9)*x(4) + 30*x(5)*x(7)*x(10)*x(4) \\
& - 60*x(5)*x(7)*x(8)*x(10)*x(4) - 60*x(5)*x(7)*x(9)*x(10)*x(4) \\
& + 30*x(5)*x(7)*x(11)*x(4) + 30*x(5)*x(7)*x(12)*x(4) \\
& - 60*x(5)*x(7)*x(8)*x(11)*x(4) - 60*x(5)*x(7)*x(9)*x(11)*x(4) \\
& - 60*x(5)*x(7)*x(8)*x(12)*x(4) - 60*x(5)*x(7)*x(9)*x(12)*x(4) = 0, \\
& \dots \quad (3.4-viii)
\end{aligned}$$

$$\begin{aligned}
h^5 f^2 f^2 f_{yy}: & 22 - 60*x(5)**2*x(7)**2*x(2) - 15*x(6)**2*x(3) \\
& - 30*x(5)*x(7)*x(3) - 30*x(5)**2*x(7)*x(3) \\
& - 30*x(6)*x(7)*x(3) - 60*x(5)*x(6)*x(7)*x(3) \\
& - 15*x(7)**2*x(3) - 60*x(5)*x(7)**2*x(3) \\
& - 60*x(5)**2*x(7)**2*x(3) + 60*x(6)**2*x(8)*x(3) \\
& + 60*x(5)*x(7)*x(8)*x(3) + 60*x(5)**2*x(7)*x(8)*x(3) \\
& + 120*x(6)*x(7)*x(8)*x(3) + 120*x(5)*x(6)*x(7)*x(8)*x(3) \\
& + 60*x(7)**2*x(8)*x(3) \\
& + 120*x(5)*x(7)**2*x(8)*x(3) - 60*x(6)**2*x(8)**2*x(3) \\
& - 120*x(6)*x(7)*x(8)**2*x(3) \\
& - 60*x(7)**2*x(8)**2*x(3) - 60*x(5)*x(6)*x(9)*x(3) \\
& + 60*x(6)**2*x(9)*x(3) + 60*x(5)**2*x(7)*x(9)*x(3) \\
& + 120*x(6)*x(7)*x(9)*x(3) + 120*x(5)*x(6)*x(7)*x(9)*x(3) \\
& + 60*x(7)**2*x(9)*x(3)
\end{aligned}$$

$$\begin{aligned}
& + 120*x(5)*x(7)**2*x(9)*x(3) + 120*x(5)*x(6)*x(8)*x(9)*x(3) \\
& - 120*x(6)**2*x(8)*x(9)*x(3) \\
& + 120*x(5)*x(7)*x(8)*x(9)*x(3) - 240*x(6)*x(7)*x(8)*x(9)*x(3) \\
& - 120*x(7)**2*x(8)*x(9)*x(3) \\
& - 60*x(5)**2*x(9)**2*x(3) + 120*x(5)*x(6)*x(9)**2*x(3) \\
& - 60*x(6)**2*x(9)**2*x(3) \\
& + 120*x(5)*x(7)*x(9)**2*x(3) - 120*x(6)*x(7)*x(9)**2*x(3) \\
& - 60*x(7)**2*x(9)**2*x(3) - 15*x(4) \\
& - 90*x(6)*x(4) - 45*x(6)**2*x(4) - 90*x(7)*x(4) \\
& - 30*x(5)*x(7)*x(4) - 30*x(5)**2*x(7)*x(4) \\
& - 90*x(6)*x(7)*x(4) - 60*x(5)*x(6)*x(7)*x(4) - 45*x(7)**2*x(4) \\
& - 60*x(5)*x(7)**2*x(4) \\
& + 180*x(6)*x(8)*x(4) + 120*x(6)**2*x(8)*x(4) + 180*x(7)*x(8)*x(4) \\
& + 60*x(5)*x(7)*x(8)*x(4) \\
& + 60*x(5)**2*x(7)*x(8)*x(4) + 240*x(6)*x(7)*x(8)*x(4) \\
& + 120*x(5)*x(6)*x(7)*x(8)*x(4) \\
& + 120*x(7)**2*x(8)*x(4) + 120*x(5)*x(7)**2*x(8)*x(4) \\
& - 60*x(6)**2*x(8)**2*x(4) \\
& - 120*x(6)*x(7)*x(8)**2*x(4) - 60*x(7)**2*x(8)**2*x(4) \\
& - 180*x(5)*x(9)*x(4) - 60*x(5)**2*x(9)*x(4) \\
& + 180*x(6)*x(9)*x(4) - 60*x(5)*x(6)*x(9)*x(4) \\
& + 120*x(6)**2*x(9)*x(4) + 180*x(7)*x(9)*x(4) \\
& + 60*x(5)**2*x(7)*x(9)*x(4) + 240*x(6)*x(7)*x(9)*x(4) \\
& + 120*x(5)*x(6)*x(7)*x(9)*x(4) \\
& + 120*x(7)**2*x(9)*x(4) + 120*x(5)*x(7)**2*x(9)*x(4) \\
& + 120*x(5)*x(6)*x(8)*x(9)*x(4) \\
& - 120*x(6)**2*x(8)*x(9)*x(4) + 120*x(5)*x(7)*x(8)*x(9)*x(4) \\
& - 240*x(6)*x(7)*x(8)*x(9)*x(4) \\
& - 120*x(7)**2*x(8)*x(9)*x(4) - 60*x(5)**2*x(9)**2*x(4) \\
& + 120*x(5)*x(6)*x(9)**2*x(4) \\
& - 60*x(6)**2*x(9)**2*x(4) + 120*x(5)*x(7)*x(9)**2*x(4)
\end{aligned}$$

$$\begin{aligned}
& -120*x(6)*x(7)*x(9)**2*x(4) \\
& -60*x(7)**2*x(9)**2*x(4) + 30*x(10)*x(4) + 90*x(6)*x(10)*x(4) \\
& + 30*x(6)**2*x(10)*x(4) + 90*x(7)*x(10)*x(4) \\
& + 60*x(6)*x(7)*x(10)*x(4) + 30*x(7)**2*x(10)*x(4) \\
& -180*x(6)*x(8)*x(10)*x(4) - 60*x(6)**2*x(8)*x(10)*x(4) \\
& -180*x(7)*x(8)*x(10)*x(4) - 120*x(6)*x(7)*x(8)*x(10)*x(4) \\
& -60*x(7)**2*x(8)*x(10)*x(4) \\
& + 180*x(5)*x(9)*x(10)*x(4) + 60*x(5)**2*x(9)*x(10)*x(4) \\
& -180*x(6)*x(9)*x(10)*x(4) - 60*x(6)**2*x(9)*x(10)*x(4) \\
& -180*x(7)*x(9)*x(10)*x(4) - 120*x(6)*x(7)*x(9)*x(10)*x(4) \\
& -60*x(7)**2*x(9)*x(10)*x(4) - 15*x(10)**2*x(4) \\
& + 30*x(11)*x(4) - 60*x(5)*x(11)*x(4) + 90*x(6)*x(11)*x(4) \\
& + 30*x(6)**2*x(11)*x(4) + 90*x(7)*x(11)*x(4) \\
& + 60*x(6)*x(7)*x(11)*x(4) + 30*x(7)**2*x(11)*x(4) \\
& -180*x(6)*x(8)*x(11)*x(4) - 60*x(6)**2*x(8)*x(11)*x(4) \\
& -180*x(7)*x(8)*x(11)*x(4) - 120*x(6)*x(7)*x(8)*x(11)*x(4) \\
& -60*x(7)**2*x(8)*x(11)*x(4) \\
& + 180*x(5)*x(9)*x(11)*x(4) + 60*x(5)**2*x(9)*x(11)*x(4) \\
& -180*x(6)*x(9)*x(11)*x(4) - 60*x(6)**2*x(9)*x(11)*x(4) \\
& -180*x(7)*x(9)*x(11)*x(4) - 120*x(6)*x(7)*x(9)*x(11)*x(4) \\
& -60*x(7)**2*x(9)*x(11)*x(4) - 30*x(10)*x(11)*x(4) \\
& + 60*x(5)*x(10)*x(11)*x(4) - 15*x(11)**2*x(4) \\
& + 60*x(5)*x(11)**2*x(4) - 60*x(5)**2*x(11)**2*x(4) \\
& + 30*x(12)*x(4) + 30*x(6)*x(12)*x(4) + 30*x(6)**2*x(12)*x(4) \\
& + 30*x(7)*x(12)*x(4) - 120*x(5)*x(7)*x(12)*x(4) \\
& -60*x(5)**2*x(7)*x(12)*x(4) + 60*x(6)*x(7)*x(12)*x(4) \\
& -120*x(5)*x(6)*x(7)*x(12)*x(4) + 30*x(7)**2*x(12)*x(4) \\
& -120*x(5)*x(7)**2*x(12)*x(4) - 180*x(6)*x(8)*x(12)*x(4) \\
& -60*x(6)**2*x(8)*x(12)*x(4) \\
& -180*x(7)*x(8)*x(12)*x(4) - 120*x(6)*x(7)*x(8)*x(12)*x(4) \\
& -60*x(7)**2*x(8)*x(12)*x(4)
\end{aligned}$$

$$\begin{aligned}
& + 180*x(5)*x(9)*x(12)*x(4) + 60*x(5)**2*x(9)*x(12)*x(4) \\
& - 180*x(6)*x(9)*x(12)*x(4) - 60*x(6)**2*x(9)*x(12)*x(4) \\
& - 180*x(7)*x(9)*x(12)*x(4) - 120*x(6)*x(7)*x(9)*x(12)*x(4) \\
& - 60*x(7)**2*x(9)*x(12)*x(4) - 30*x(10)*x(12)*x(4) \\
& + 60*x(6)*x(10)*x(12)*x(4) + 60*x(7)*x(10)*x(12)*x(4) \\
& - 30*x(11)*x(12)*x(4) + 60*x(5)*x(11)*x(12)*x(4) \\
& + 60*x(6)*x(11)*x(12)*x(4) - 120*x(5)*x(6)*x(11)*x(12)*x(4) \\
& + 60*x(7)*x(11)*x(12)*x(4) \\
& - 120*x(5)*x(7)*x(11)*x(12)*x(4) - 15*x(12)**2*x(4) \\
& + 60*x(6)*x(12)**2*x(4) - 60*x(6)**2*x(12)**2*x(4) \\
& + 60*x(7)*x(12)**2*x(4) - 120*x(6)*x(7)*x(12)**2*x(4) \\
& - 60*x(7)**2*x(12)**2*x(4) = 0, \tag{3.4-ix}
\end{aligned}$$

$$\begin{aligned}
h^5 f^3 f_{yy}^2: & 8 - 60*x(5)**2*x(6)*x(7)*x(2) - 60*x(5)**2*x(7)**2*x(2) \\
& - 15*x(6)**2*x(3) - 30*x(6)*x(7)*x(3) \\
& - 60*x(5)**2*x(6)*x(7)*x(3) - 15*x(7)**2*x(3) \\
& - 60*x(5)**2*x(7)**2*x(3) + 30*x(6)**2*x(8)*x(3) \\
& + 60*x(6)*x(7)*x(8)*x(3) + 30*x(7)**2*x(8)*x(3) \\
& - 30*x(5)**2*x(9)*x(3) + 30*x(6)**2*x(9)*x(3) \\
& + 60*x(6)*x(7)*x(9)*x(3) + 30*x(7)**2*x(9)*x(3) - 15*x(4) \\
& - 15*x(6)**2*x(4) - 30*x(6)*x(7)*x(4) \\
& - 15*x(7)**2*x(4) + 30*x(6)**2*x(8)*x(4) \\
& + 60*x(6)*x(7)*x(8)*x(4) + 30*x(7)**2*x(8)*x(4) \\
& - 30*x(5)**2*x(9)*x(4) + 30*x(6)**2*x(9)*x(4) \\
& + 60*x(6)*x(7)*x(9)*x(4) + 30*x(7)**2*x(9)*x(4) \\
& + 15*x(10)*x(4) + 15*x(11)*x(4) - 60*x(5)**2*x(11)*x(4) \\
& + 15*x(12)*x(4) - 60*x(6)**2*x(12)*x(4) \\
& - 120*x(6)*x(7)*x(12)*x(4) - 60*x(7)**2*x(12)*x(4) = 0, \tag{3.4-x}
\end{aligned}$$

$$\begin{aligned}
h^5 f^3 f_y f_{yyy}: & 28 - 40*x(5)**3*x(7)*x(2) - 120*x(5)*x(6)**2*x(7)*x(2) \\
& - 240*x(5)*x(6)*x(7)**2*x(2) \\
& - 120*x(5)*x(7)**3*x(2) - 15*x(6)*x(3) - 20*x(6)**3*x(3)
\end{aligned}$$

$$\begin{aligned}
& -15*x(7)*x(3) - 40*x(5)**3*x(7)*x(3) \\
& - 60*x(6)**2*x(7)*x(3) - 120*x(5)*x(6)**2*x(7)*x(3) \\
& - 60*x(6)*x(7)**2*x(3) - 240*x(5)*x(6)*x(7)**2*x(3) \\
& - 20*x(7)**3*x(3) - 120*x(5)*x(7)**3*x(3) + 30*x(6)*x(8)*x(3) \\
& + 40*x(6)**3*x(8)*x(3) + 30*x(7)*x(8)*x(3) \\
& + 120*x(6)**2*x(7)*x(8)*x(3) + 120*x(6)*x(7)**2*x(8)*x(3) \\
& + 40*x(7)**3*x(8)*x(3) - 30*x(5)*x(9)*x(3) \\
& - 40*x(5)**3*x(9)*x(3) + 30*x(6)*x(9)*x(3) \\
& + 40*x(6)**3*x(9)*x(3) + 30*x(7)*x(9)*x(3) \\
& + 120*x(6)**2*x(7)*x(9)*x(3) + 120*x(6)*x(7)**2*x(9)*x(3) \\
& + 40*x(7)**3*x(9)*x(3) - 65*x(4) \\
& - 15*x(6)*x(4) - 20*x(6)**3*x(4) - 15*x(7)*x(4) \\
& - 60*x(6)**2*x(7)*x(4) - 60*x(6)*x(7)**2*x(4) \\
& - 20*x(7)**3*x(4) + 30*x(6)*x(8)*x(4) + 40*x(6)**3*x(8)*x(4) \\
& + 30*x(7)*x(8)*x(4) \\
& + 120*x(6)**2*x(7)*x(8)*x(4) + 120*x(6)*x(7)**2*x(8)*x(4) \\
& + 40*x(7)**3*x(8)*x(4) - 30*x(5)*x(9)*x(4) \\
& - 40*x(5)**3*x(9)*x(4) + 30*x(6)*x(9)*x(4) \\
& + 40*x(6)**3*x(9)*x(4) + 30*x(7)*x(9)*x(4) \\
& + 120*x(6)**2*x(7)*x(9)*x(4) + 120*x(6)*x(7)**2*x(9)*x(4) \\
& + 40*x(7)**3*x(9)*x(4) + 65*x(10)*x(4) \\
& + 65*x(11)*x(4) - 120*x(5)*x(11)*x(4) - 40*x(5)**3*x(11)*x(4) \\
& + 65*x(12)*x(4) - 120*x(6)*x(12)*x(4) \\
& - 40*x(6)**3*x(12)*x(4) - 120*x(7)*x(12)*x(4) \\
& - 120*x(6)**2*x(7)*x(12)*x(4) - 120*x(6)*x(7)**2*x(12)*x(4) \\
& - 40*x(7)**3*x(12)*x(4) = 0, \tag{3.4-xi}
\end{aligned}$$

$$\begin{aligned}
h^5 f_{yyyy}^{(4)}: & 32 - 80*x(5)**4*x(1) - 80*x(5)**4*x(2) \\
& - 80*x(6)**4*x(2) - 320*x(6)**3*x(7)*x(2) \\
& - 480*x(6)**2*x(7)**2*x(2) - 320*x(6)*x(7)**3*x(2) \\
& - 80*x(7)**4*x(2) - 5*x(3) - 80*x(6)**4*x(3)
\end{aligned}$$

$$\begin{aligned}
& -320*x(6)**3*x(7)*x(3) - 480*x(6)**2*x(7)**2*x(3) \\
& -320*x(6)*x(7)**3*x(3) - 80*x(7)**4*x(3) - 85*x(4) = 0 \quad (3.4\text{-xii})
\end{aligned}$$

where

$$\begin{aligned}
x(1) &= w_1, x(2) = w_2, x(3) = w_3, x(4) = w_4, x(5) = a_1, x(6) = a_2. \\
x(7) &= a_3, x(8) = a_4, x(9) = a_5, x(10) = a_6, x(11) = a_7 \text{ and } x(12) = a_8.
\end{aligned}$$

Similarly, equations (3.4-i)–(3.4-xii) are solved simultaneously by using the NAG routine (Subroutine C05NBF) for solving a system of non-linear equations to give the required parameters, i.e.,

$$\begin{aligned}
w_1 = x(1) &= 0.2615038147, w_2 = x(2) = -0.2765809214, w_3 = x(3) = 0.5947141647 \\
w_4 = x(4) &= 0.4203629420, a_1 = x(5) = 1.5471214403, a_2 = x(6) = 0.1756458393 \\
a_3 = x(7) &= 0.1243059001, a_4 = x(8) = 0.1009316694, a_5 = x(9) = 0.1100539630 \\
a_6 = x(10) &= 0.99974318, x_7 = x(11) = -0.0928890403, x_8 = x(12) = -0.6201812828 \\
&\dots \quad (3.5)
\end{aligned}$$

Thus, the new fifth order WRK method can be written as follows

$$\begin{aligned}
y_{n+1} = y_n + h \left[0.2615038147 \left(\frac{k_1 + k_2}{2} \right) - 0.2765809214 \left(\frac{k_2 + k_3}{2} \right) \right. \\
\left. + 0.5947141647 \left(\frac{k_3 + k_4}{2} \right) + 0.4203629420 \left(\frac{k_4 + k_5}{2} \right) \right] \quad (3.6)
\end{aligned}$$

where

$$\begin{aligned}
k_1 &= f(y_n) \\
k_2 &= f(y_n + 1.5471214403hk_1) \\
k_3 &= f(y_n + 0.1756458393hk_1 + 0.1243059001hk_2) \\
k_4 &= f(y_n + 0.1009316694hk_1 + 0.1100539630hk_2 + 0.2890143692hk_3) \\
k_5 &= f(y_n + 0.9997431862hk_1 - 0.0928890403hk_2 - 0.6201812828hk_3 \\
&\quad + 0.7133271396hk_4). \quad \dots \quad (3.7)
\end{aligned}$$

By rationalizing the coefficients in equation (3.5) the fifth order WRK method can also be written in rational form as given in Appendix 1.

3.1 Error Analysis

By substituting the values of $a_i, 1 \leq i \leq 8$ and $w_i, 1 \leq i \leq 4$ in (3.5) into (3.1) and (3.2) using Mathematica and evaluating all the terms up to (h^6) to represent the local

truncation error (LTE) for this method we have

$$\begin{aligned}
 LTE = h^6 & \left[\frac{1}{720} f f_y^5 + 0.0018022816 f^2 f_y^3 f_{yy} - 0.0166861138 f^3 f_y f_{yy}^2 \right. \\
 & + 0.0082646021 f^3 f_y^2 f_{yyy} + 0.0041171137 f^4 f_{yy} f_{yyy} \\
 & \left. - 0.0023096163 f^4 f_y f_{yyy} + 0.0000588245 f^5 f_{yyyy} \right] \\
 & \dots
 \end{aligned} \tag{3.8}$$

4 STABILITY ANALYSIS OF THE FIFTH ORDER WRK METHOD

We examine the stability region for the fifth order arithmetic mean WRK method with the test equation $y' = \lambda y$ and we obtain

$$\begin{aligned}
 k_1 &= \lambda(y_n) \\
 k_2 &= \lambda(y_n + 1.5471214403 h k_1) \\
 k_3 &= \lambda(y_n + 0.1756458393 h k_1 + 0.1243059001 h k_2) \\
 k_4 &= \lambda(y_n + 0.1009316694 h k_1 + 0.1100539630 h k_2 + 0.2890143692 h k_3) \\
 k_5 &= \lambda(y_n + 0.9997431862 h k_1 - 0.0928890403 h k_2 - 0.6201812828 h k_3 \\
 & \quad + 0.7133271396 h k_4). \dots
 \end{aligned} \tag{4.1}$$

By substituting k_i , $1 \leq i \leq 5$ in equation (4.1) and w_i , $1 \leq i \leq 4$ in equation (3.5) into the fifth order WRK formula, i.e.,

$$y_{n+1} = y_n + h \left[\sum_{i=1}^4 w_i \left(\frac{k_i + k_{i+1}}{2} \right) \right] \tag{4.2}$$

we obtain

$$\begin{aligned}
 y_{n+1} &= y_n + (h\lambda)y_n + 0.5(h\lambda)^2 y_n + 0.166667(h\lambda)^3 y_n + 0.0416667(h\lambda)^4 y_n \\
 & \quad + 0.00833333(h\lambda)^5 y_n + 0(h^6). \dots
 \end{aligned} \tag{4.3}$$

By rationalizing the coefficients in equation (4.3) we have

$$\begin{aligned}
 y_{n+1} &= y_n + (h\lambda)y_n + \frac{1}{2}(h\lambda)^2 y_n + \frac{1}{6}(h\lambda)^3 y_n + \frac{1}{24}(h\lambda)^4 y_n + \frac{1}{120}(h\lambda)^5 + 0(h^6) \\
 & \dots
 \end{aligned} \tag{4.4}$$

By substituting $h\lambda = z$ in (4.4), we can show that

$$y_{n+1} = y_n + y_n \left[z + \frac{z^2}{2} + \frac{z^3}{6} + \frac{z^4}{24} + \frac{z^5}{120} \right] + 0(z^6). \quad (4.5)$$

Following Evans and Sanugi [1991], we write

$$\frac{y_{n+1}}{y_n} = Q$$

in the equation (4.5), to obtain

$$Q = 1 + z + \frac{1}{2}z^2 + \frac{1}{6}z^3 + \frac{1}{24}z^4 + \frac{1}{120}z^5 + 0(z^6). \quad (4.6)$$

To determine the stability region of the fifth order WRK formula in the complex plane that satisfy the condition

$$\left| \frac{y_{n+1}}{y_n} \right| = |Q| < 1$$

i.e.,

$$\left| 1 + z + \frac{z^2}{2} + \frac{z^3}{6} + \frac{z^4}{24} + \frac{z^5}{120} \right| < 1 \quad (4.7)$$

by the use of Mathematica, we can plot the graphic surface defined by equation (4.7) as shown in Figure 1 can also plot the stability region defined by the formula in equation (4.7) as shown in Figure 2.

5 NUMERICAL EXAMPLE

We consider the IVP

$$y' = x - y + 1, \quad y(0) = 1, \quad 0 \leq x \leq 1 \quad (5.1)$$




Figure 1 Graphic surface defined by fifth order WRK formula.




Figure 2 Stability region for fifth order WRK formula.

Table 2 Absolute error by various formula for solving equation (5.1).

| x | <i>Classical</i> <i>AM-RK4</i> | <i>RK4(5)</i> <i>Merson</i> | <i>RK5(6)</i> <i>Nystrom</i> | <i>WRK-RK5(5)</i> |
|-----|-----------------------------------|--------------------------------|---------------------------------|-------------------|
| 0.1 | 0.8196E-07 | 0.1252E-07 | 0.1707E-08 | 0.1380E-08 |
| 0.2 | 0.1483E-06 | 0.2266E-07 | 0.3089E-08 | 0.2498E-08 |
| 0.3 | 0.2013E-06 | 0.3075E-07 | 0.4192E-08 | 0.3391E-08 |
| 0.4 | 0.2429E-06 | 0.3710E-07 | 0.5058E-08 | 0.4091E-08 |
| 0.5 | 0.2747E-06 | 0.4196E-07 | 0.5721E-08 | 0.4627E-08 |
| 0.6 | 0.2983E-06 | 0.4556E-07 | 0.6211E-08 | 0.5024E-08 |
| 0.7 | 0.3149E-06 | 0.4810E-07 | 0.6557E-08 | 0.5303E-08 |
| 0.8 | 0.3256E-06 | 0.4974E-07 | 0.6781E-08 | 0.5484E-08 |
| 0.9 | 0.3315E-06 | 0.5063E-07 | 0.6902E-08 | 0.5583E-08 |
| 1.0 | 0.3332E-06 | 0.5090E-07 | 0.6939E-08 | 0.5613E-08 |

and the exact solution is $y(x) = x + \exp(-x)$. The absolute errors in the numerical solution using formula (3.1)–(3.2) compared with the fourth order RK formula (classical formula), RK5(6)-Nystrom [7] and RK4(5)-Merson [9] are shown in Table 2.

From Table 2, we can see that the accuracy obtained from using WRK-RK5(5) is better than the RK4, RK4(5)-Merson and RK5(6)-Nystrom methods. When we make a work comparison with the fifth order RK5(6)-Nystrom method, then this new fifth order method saves one function evaluation. From the above discussion we can conclude that contrary to Butcher [1] a fifth order arithmetic mean weighted Runge-Kutta method with five stages does exist. The study of fifth order weighted Runge-Kutta methods for a variety of means are under further investigation.

References

- [1] J. C. Butcher, *The numerical analysis of ordinary differential equations: Runge-Kutta and general linear methods*, John Wiley & Sons, Chichester, (1987).
- [2] D. J. Evans and A. R. Yaakub, A New Fourth Order Runge-Kutta Method Based On The Contraharmonic Mean Formula, Loughborough University of Technology, Department of Computer Studies, *Report No. 849* (1993).
[Contributed Talks at the VI-th SERC Numerical Analysis Summer School, Leicester University, 25–29 July 1994]
- [3] D. J. Evans and A. R. Yaakub, A New 4th Order Runge-Kutta Method Based On The Centroidal Mean Formula, Loughborough University of Technology, Department of Computer Studies, *Report No. 851*. (1993)
- [4] D. J. Evans and N. Yaacob, A Fourth Order Runge-Kutta Method Based On The Heronian Mean Formula, Loughborough University of Technology, Department of Computer Studies, *Report No. 879* (1994).
- [5] D. J. Evans and B. B. Sanugi, A New 4th Order Runge-Kutta Method for Initial Value Problems, in S.O. Fatunla (Ed.), *Computational Mathematics II*, Boole Press, Dublin (1987).
- [6] D. J. Evans and B. B. Sanugi, A Comparison of Nonlinear Trapezoidal Formulae for Solving Initial Value Problems, *Intern. J. Computer Math.*, **41**, (1991) 65–79.
- [7] J. D. Lambert, *Computational Methods In Ordinary Differential Equations*, John Wiley & Sons, London (1973).
- [8] J. D. Lambert, *Numerical Methods For Ordinary Differential Systems*, John Wiley & Sons, London (1991).
- [9] R. H. Merson, An Operational Method For The Study of Integrating processes, *Proc. Symp. Data Processing*, Weapon Research Establishment, Salisbury, S. Australia (1957).
- [10] B. B. Sanugi and D. J. Evans, A New Fourth Order Runge-Kutta Formula Based on Harmonic Mean, Loughborough University of Technology, Department of Computer Studies, *Report No. 799* (1993).
- [11] A. R. Yaakub and D. J. Evans, A New Fourth Order Runge-Kutta Methods Based On The Root Mean Square Formula, Loughborough University of Technology, Department of Computer Studies, *Report No. 862* (1993).
- [12] Bahrom Sanugi and Nazeeruddin Yaacob, A New 5th Order 5 stage Runge-Kutta Method for Initial Value Type Problems in ODE's. *Int. Jour. Comp. Maths* (1995) (in press).

APPENDIX 1

By use of Mathematica to rationalize the coefficients in equation (3.5), we obtain

$$\begin{aligned}
 w_1 &= 0.2615038147 = \frac{28449}{108790}, & w_2 &= -0.2765809214 = \frac{40890}{147841} \\
 w_3 &= 0.5947141647 = \frac{211318}{355327}, & w_4 &= 0.4203629420 = \frac{65601}{156058} \\
 a_1 &= 1.5471214403 = \frac{479767}{310103}, & a_2 &= 0.1756458393 = \frac{14081}{80167} \\
 a_3 &= 0.1243059001 = \frac{7768}{62491}, & a_4 &= 0.1009316694 = \frac{20551}{203613} \\
 a_5 &= 0.1100539630 = \frac{15153}{137687}, & a_6 &= 0.9997431862 = \frac{151822}{151861} \\
 a_7 &= -0.0928890403 = \frac{4909}{52848}, & a_8 &= -0.6201812828 = \frac{91411}{147394}
 \end{aligned}$$

$$a_{11} = \frac{1}{2} - a_4 - a_5 = 0.2890143692 = \frac{32604}{112811}$$

$$a_{22} = 1 - a_6 - a_7 - a_8 = 0.7133271396 = \frac{548973}{769595}$$

and this new fifth order WRK method can be written in rational form as

$$y_{n+1} = y_n + h \left[\frac{28449}{108790} \left(\frac{k_1 + k_2}{2} \right) - \frac{40890}{147841} \left(\frac{k_2 + k_3}{2} \right) \right. \\ \left. + \frac{211318}{355327} \left(\frac{k_3 + k_4}{2} \right) + \frac{65601}{156058} \left(\frac{k_4 + k_5}{2} \right) \right]$$

where

$$k_1 = f(y_n)$$

$$k_2 = f\left(y_n + \frac{479767}{310103} h k_1\right)$$

$$k_3 = f\left(y_n + \frac{14081}{80167} h k_1 + \frac{7768}{62491} h k_2\right)$$

$$k_4 = f\left(y_n + \frac{20551}{203613} h k_1 + \frac{15153}{137687} h k_2 + \frac{32604}{112811} h k_3\right)$$

$$k_5 = f\left(y_n + \frac{151822}{151861} h k_1 - \frac{4909}{52848} h k_2 - \frac{91411}{147394} h k_3 + \frac{548973}{769595} h k_4\right).$$