The Integral is the Area Under the Curve

Kieron Eisma

November 2026

Introduction

Anybody who took calculus in high school was most certainly taught the following. To find the area under the curve $y = f(x)$ over some interval $[a, b]$, one uses the formula

$$A = \int_a^b f(x)\,dx$$

by rote, without question. Under the definition of the Riemann (Darboux) integral, where we are using rectangles to approximate the area under a sufficiently well-behaved function, one can see why one can interpret the integral as such.

In this paper, we shall use measure theory to give an alternative proof.

1 Geometric Preliminaries




Figure 1: The function f and set A_f. Obviously, the space X is an abstract set; this diagram is only meant to serve as a visual aid. Image generated by ChatGPT 6.

We let $(X, \mathcal{M}, \mu)$ be a σ-finite measure space. It will suffice to prove the desired result for $f \in L^+(\mu)$, that is, for non-negative functions f, for if $f \in L^1(\mu)$, we can simply apply the result to the positive and negative parts of f. To keep things clean, we also assume that $f(x) < \infty$ for all $x \in X$.

The output space of our function is $[0, \infty]$ (we use the extended real number system). Thus, the area bounded by our curve will be a subset of $X \times [0, \infty]$. Denote this set by A_f. Geometrically, for a point $(x, y) \in X \times [0, \infty]$ to be contained in A_f, we must have that $y \leq f(x)$, as illustrated in Figure 1.

Hence, we can define

$$A_f = \{(x, y) \in X \times [0, \infty] : y \leq f(x)\},$$

which is equivalent to writing

$$A_f = \{(x, y) \in X \times [0, \infty] : f(x) - y \geq 0\}.$$

2 Proving the Result

Any hope of proving this result rests on our ability to measure the size of this set.

Lemma 2.1 *The set A_f is $\mathcal{M} \otimes \mathcal{B}_{\mathbb{R}}$-measurable.*

Proof: Define the map $W : X \times [0, \infty] \to [0, \infty]$ by $W(x, y) = f(x) - y$. Then,

$$A_f = W^{-1}([0, \infty]).$$

Thus, if we can show that W is measurable, then we have our result. Indeed, consider the maps $(z, y) \mapsto z - y$ and $(x, y) \mapsto (f(x), y)$. The first map is measurable by continuity, and the second is measurable by Proposition 2.4 in Folland [Fol99]. The composition of measurable functions is measurable, so it follows that W is measurable, and hence that A_f is $\mathcal{M} \otimes \mathcal{B}_{\mathbb{R}}$-measurable. $\square$

Now that we know A_f is measurable, what is its measure? In the result below, m denotes the standard Lebesgue measure.

Theorem 2.1 (The Integral is the Area Under the Curve.)

$$(\mu \times m)(A_f) = \int f \, d\mu.$$

Proof: σ-finiteness allows us to conclude[1]

$$(\mu \times m)(A_f) = \int m((A_f)_x) \, d\mu(x). \tag{1}$$

Recal that the x-section of A_f is defined as

[1] Theorem 2.36 in [Fol99]

$$(A_f)_x = \{y \in [0, \infty] : (x, y) \in A_f\} = [0, f(x)].$$

Substituting the above equation into Equation 1, and using the properties of the Lebesgue measure, we see that

$$\begin{aligned}(\mu \times m)(A_f) &= \int (f(x) - 0)\, d\mu(x) \\ &= \int f(x)\, d\mu(x),\end{aligned}$$

which is the desired claim. ■

This is for the entire "X-line", so if we want to integrate $f \in L^+(\mu)$ over some set $E \in \mathcal{M}$, we can apply Theorem 2.1 to the function $g := f\chi_E \in L^+(\mu)$, where χ_E denotes the indicator function of the set E, to obtain

$$A_g = \int f\chi_E \, d\mu := \int_E f \, d\mu;$$

The indicator kills the function outside the target interval E. As stated before, for a more general function $f \in L^1(\mu)$, we can apply this result to the positive and negative parts of f, respectively (which are measurable), giving us what we want!

References

[Fol99] Gerald B. Folland. *Real Analysis: Modern Techniques and Their Applications*. 2nd ed. Vol. 234. Pure and Applied Mathematics. New York: John Wiley & Sons, 1999. ISBN: 978-0-471-31716-6.