

# INTRODUCTION TO SOCIAL NETWORK ANALYSIS AND NETWORK SCIENCE METHODS

Workshop, ICHPS 2023, Scottsdale, AZ

---

James O'Malley, Ph.D.

Department of Biomedical Data Science

The Dartmouth Institute for Health Policy and  
Clinical Practice

Geisel School of Medicine at Dartmouth

Email: [James.OMalley@Dartmouth.edu](mailto:James.OMalley@Dartmouth.edu)

# Link to Computing Materials Used in Workshop

- **Github** site with R scripts and data used in the illustrative examples presented in workshop:

<https://github.com/kiwijomalley/ICHPS-Social-Network-Analysis-Workshop-2023>

- **GRANDPA algorithm**
  - Use to generate random networks for analysis that emulate a base network.
  - Very useful if network data is confidential
  - Bobak CA, Zhao Y, Levy JJ, and O'Malley AJ. (2022). GRANDPA: GeneRAtive Network sampling using Degree and Property Augmentation applied to the analysis of partially confidential healthcare networks. doi.org/10.48550/arXiv.2211.15000
  - Presented in a conference poster by Carly Bobak on January 9, 2023

# Example network and data: Small physician clinic in the Boston area

- **Relational data (adjacency matrix form):**  
ICHPS\_PhysPractBin.txt
- **Relational data (edgelist form):** ICHPS\_ClinEdgeList.txt
- **Node attribute data:** ICHPS\_nodecov.dat
- **References:**
  - Keating, Ayanian, Cleary, and Marsden (2007)
  - O'Malley and Marsden (2008)

# Segments of Code for Three Key Problems

- I. Descriptive analyses involving networks
  - Visualizing a network
  - Descriptive features of networks
- II. Statistical analyses of social influence or peer effects
  - Formation of weight matrix
  - Cross-sectional cases
- III. Statistical analysis of relational data
  - Exponential random graph models
  - Latent-space models

# I. Descriptive Analyses

- **R script:** WorkshopICHPS2023.R
  - Loads network
  - Forms directed and undirected binary-valued network objects
  - Makes plots of network
  - Computes summary measures of the network and of physicians' positions within the network

# Loading Physician Practice Network

Adjacency matrix



- `reldata <- scan("ICHPS_PhysPractBin.txt")`
- `nr <- sqrt(length(reldir))` #Number of physicians
- `reldir <- matrix(reldir,ncol=nr,nrow=nr,byrow=T)`

## #Directed

- `reldir <- ifelse(reldir>0,1,0)` #Directed
  - Can use more stringent threshold (greater # conversations) to obtain sparser network by using threshold of 1

## #Undirected

- `relnet <- ifelse(reldir+t(reldir)>0,1,0)` #Undirected
  - Use “And Rule” (threshold of >1) to obtain sparser network
- `pnet <-`  
`network(reldir,directed=TRUE,matrixtype="adjacency")`

## Aside: Forming an edgelist from an adjacency matrix with IDs from 1 to nr

- `idto <- seq(1,nr)`
- `edgelist=c()`
- `for (i in 1:nr) {`
  - `edge <- idto[(reldir[i,]==1)]`
  - `nc=length(edge)`
  - `edgelist=rbind(edgelist,cbind(rep(i,nc),edge))`
- `}`
- `edgelist <- data.frame(edgelist)`
- `names(edgelist) <- c("source","target")`
- `pnet <-`  
`network(reldir,directed=TRUE,matrixtype="edgelist")`

# More general loading of network data

- `pnet <-  
network(yourdatafile,directed=TRUE,matrixtype="edgelist  
")`



Use one of: adjacency, edgelist, incidence



# Loading attribute data

# Load attribute data


- `covdata <- scan("ICHPS_nodecov.dat")`
- `covdata <- matrix(covdata,nrow=nr,byrow=T)`
- `colnames(covdata) <-  
c("id","male","whexpert","pctwom","numsess","practice",  
"sumhrt")`

# Software for descriptive analysis of network

- R is convenient
  - sna package contains many built-in features
  - igraph package
    - R: `install.packages("sna","igraph")`
    - R: `library(sna)`
    - R: `library(igraph)`
- Python
  - networkx package

# Plotting network using SNA

- **#Plot from adjacency matrix**
- `plot(pnet,mode="fruchtermanreingold",displaylabels=T)`  
`gplot(pnet,gmode="digraph",mode="fruchtermanreingold")`




# Using igraph

- Let **edgelist** be a N by 2 matrix in R with the network represented as an edgelist
- `nodes <- unique(c(edgelist[,1],edgelist[,2]))`
- **# Make graph object from edgelist:**
- `gnet <- graph_from_data_frame(d=edgelist, vertices=nodes, directed=TRUE)`
- `print(gnet, e=TRUE, v=TRUE)`

# Plotting network in igraph

- `V(gnet)$color <- covdata$practice`
- `V(gnet)$size <- 10*(covdata$whexpert+1)`
- `par(mar=c(0,0,0,0))`
- `plot(gnet,`
- `vertex.color = V(gnet)$color, # Color of nodes`
- `vertex.size = V(gnet)$size, # Size of nodes`
- `vertex.label.color = "black", # change color of labels`
- `vertex.label.cex = .75, # change size of labels to 75% of original size`
- `edge.curved=.25, # add a 25% curve to the edges`
- `edge.color="grey20", # change edge color to grey`
- `edge.arrow.size=0.3)`
- `dev.copy2pdf(file=paste(outdir,"PhysNetNicePlot.pdf",sep=""), width=6, height=6) #to file`

# Visualization of network of physicians' professional relationships within a medical practice



# Computing summary measures of networks in R

- Use SNA or igraph
  - You might detach igraph if you want to use SNA as igraph masks functions (has other functions by the same name) or precede function name with package name (e.g., igraph::grecip)
  - detach("package:igraph", unload = TRUE)
- Dyad and Triad Census'
  - dyadc=dyad.census(reldir)
  - recip=grecip(reldir)
  - triadc\_mut=triad.census(reldir,g="graph")
  - triadc\_dir=triad.census(reldir,g="digraph")
  - trans=gtrans(reldir,mode="digraph")

# Summary measures cont.

- Degree distributions
  - `idegree=degree(reldir,cmode="indegree")`
  - `odegree=degree(reldir,cmode="outdegree")`
  - `central=centralization(reldir,degree)`
- Centrality measures
  - `closecent=closeness(reldir,gmode="digraph")`
  - `bcent=betweenness(reldir,gmode="digraph")`
  - `eigcent=evcent(reldir,gmode="digraph",use.eigen=FALSE)`
  - `powcent=bonpow(reldir,gmode="digraph")`



# Aside: Manual computation of eigenvector centrality

# Column standardize as per Katz (1953): undirected network

- `on=as.vector(rep(1,nrow(relmut)))`
- `rscale=as.vector(relmut %*% on)`
- `scnet=diag(rscale-1,nrow=length(cscale))`
- `srelmut=relmut %*% scnet` #Row-stochastic adjacency mat
- `seigcent=evcent(srelmut,gmode="digraph",use.eigen=FALSE)`

Can't  
be 0!

#manual

- `sesys=eigen(srelmut,symmetric=TRUE)` #First eigenvalue is 1
- `seigcent2=abs(sesys$vector[,1])` #Same as seigcent

# Differs from:

- `eigcent = evcent(relmut,gmode="digraph",use.eigen=FALSE)`

# R Output

```
sesys$values
```

```
[1] 1.00000000 0.58674579 -0.50090590 0.47303368 -0.43397276 0.42666878 -0.42491411 -  
0.39345582  
[9] 0.37479529 -0.37272600 0.34943660 -0.34833131 0.29465182 -0.29277386 -0.26530439  
0.26150139  
[17] -0.25725084 -0.25079539 -0.23908776 0.23292529 0.17182520 -0.15703102 -0.14330681 -  
0.12994348  
[25] 0.12570645 -0.10902035 -0.10191533 0.09780081 0.06373494 -0.06033018 0.04989629 -  
0.04039014  
[33] 0.01273312
```

```
> -sesys$vector[,1]
```


```
[1] 0.14294834 0.16081688 0.08934271 0.17868542 0.07147417 0.12507980 0.23229105  
0.14294834  
[9] 0.28589668 0.08934271 0.10721125 0.12507980 0.12507980 0.03573708 0.17868542  
0.23229105  
[17] 0.12507980 0.07147417 0.30376522 0.03573708 0.30376522 0.05360563 0.08934271  
0.26802813  
[25] 0.08934271 0.03573708 0.42884501 0.10721125 0.10721125 0.21442251 0.14294834  
0.03573708  
[33] 0.16081688
```

## II. Statistical models involving comparative analysis of multiple networks

- Do social network characteristics correlate with other variables of interest?
- Often involves use of regression and hierarchical regression models
- **Original data:** Keating, Ayanian, Cleary, and Marsden (2007)
- **R script:** ICHPS2023models.R

# Example plot of network and statistical analysis (O'Malley and Marsden, 2008)

- `pnet <- network(physnetwork, directed=TRUE, matrixtype="adjacency",  
vertex.attr=nodecov,  
vertex.attrnames = c("male", "whexpert",  
"pctwom", "numsess", "practice", "bcma", "bima",  
"bpp", "wnhlth", "numcat", "pctcat"))`
- `plot(pnet, mode = "fruchtermanreingold", displaylabels=T)`



# Simplest model

- `model1a <- ergm(pnet~edges)`
- Evaluating log-likelihood at the estimate.
- Formula: `pnet ~ edges`
- Iterations: 5 out of 20
- Monte Carlo MLE Results:
  - Estimate Std. Error MCMC % p-value
  - edges -1.70084 0.08517 0 <1e-04 \*\*\*
  - ---
  - Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1
- Null Deviance: 1463.9 on 1056 degrees of freedom
- Residual Deviance: 908.6 on 1055 degrees of freedom
- AIC: 910.6 BIC: 915.5 (Smaller is better.)

What is this the log-odds of?

# Larger model cont.

- Formula: `pnet ~ edges + mutual + nodecov("whexpert") + nodecov("pctwom") + nodecov("numsess") + nodematch("male", diff = F) + nodematch("bcma", diff = F) + nodematch("bima", diff = F) + nodematch("bpp", diff = F) + nodematch("wnhlth", diff = F)`

- Monte Carlo MLE Results:

	Estimate	Std. Error	MCMC %	p-value
edges	-4.560879	0.510615	0	< 1e-04 ***
mutual	0.851292	0.293985	0	0.003862 **
nodecov.whexpert	-0.391256	0.279232	0	0.161455
nodecov.pctwom	-0.001583	0.004530	0	0.726830
nodecov.numsess	0.159686	0.039360	0	< 1e-04 ***
nodematch.male	0.646502	0.181881	0	0.000396 ***
nodematch.bcma	0.739658	0.278889	0	0.008119 **
nodematch.bima	0.277871	0.199195	0	0.163321
nodematch.bpp	1.396742	0.255284	0	< 1e-04 ***
nodematch.wnhlth	-0.046937	0.221694	0	0.832366

---

- Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

- AIC: 838.3 BIC: 887.9 (Smaller is better.)

Ego (originator) covariate

Homophily covariate

# Separate node-match (homophily) coefficients under alternative specification

- `model1f <- ergm(pnet~edges + mutual + nodecov("whexpert") + nodecov("pctwom") + nodecov("numsess") + nodematch("male",diff=T) + nodematch("practice",diff=T))`


- Estimate Std. Error MCMC % p-value

• edges	-1.134295	0.417598	0 0.00671 **
• mutual	0.615252	0.316615	0 0.05226 .
• nodecov.whexpert	-0.509118	0.293552	0 0.08315 .
• nodecov.pctwom	-0.022326	0.005604	0 < 1e-04 ***
• nodecov.numsess	-0.007532	0.045438	0 0.86837
• <b>nodematch.male.0</b>	<b>1.534152</b>	<b>0.247805</b>	<b>0 &lt; 1e-04 ***</b>
• <b>nodematch.male.1</b>	<b>-1.112795</b>	<b>0.420071</b>	<b>0 0.00819 **</b>
• nodematch.practice.1	1.320874	0.849377	0 0.12022
• nodematch.practice.2	0.466084	0.215630	0 0.03088 *
• nodematch.practice.3	2.131637	0.423777	0 < 1e-04 ***
• nodematch.practice.4	1.981935	0.489350	0 < 1e-04 ***

Allows  
nodematch.male  
coefficient to vary  
by sex

# Assessing Goodness of Fit with Respect to In-Degree Distribution

- `model1e.gof <-  
gof(model1e~idegree,control=control.gof.ergm(nsim=100)  
,verbose=T)`
- `plot(model1e.gof)`





# Estimating Latent-space models in R

- `ergmm(formula, response = NULL, family = "Bernoulli", fam.par = NULL, control = control.ergmm(), user.start = list(), prior = ergmm.prior(), tofit = c("mcmc", "mkl", "mkl.mbc", "procrustes", "klswitch"), Z.ref = NULL, Z.K.ref = NULL, seed = NULL, verbose = FALSE)`
  - “family” command allows different distributions in the exponential family (as for generalized linear models)
  - “prior” command gives some control over prior distributions for Bayesian analysis
  - “tofit” controls which elements of estimation are performed
- `?ergmm` to get help and then `terms.ergmm` to get list of terms that are supported
  - Note that there are no mutual, triadic or higher-order network statistics are allowed in this model!

# Equating Latentnet's ERGMM and Statnet's ERGM

- ERGMM Formula: `pnet ~ nodematch("male", diff = F)`
- Attribute: **edges (included by default in ERGMM)**
- Model: Bernoulli
- MCMC sample of size 4000, draws are 10 iterations apart, after burnin of 10000 iterations.
- Covariate coefficients posterior means:

	Estimate	2.5%	97.5%	2*min(Pr(>0),Pr(<0))	
(Intercept)	-2.15650	-2.44573	-1.8870		< 2.2e-16 ***
nodematch.male	0.78345	0.43672	1.1398		< 2.2e-16 ***

- ERGM Formula: `pnet ~ edges + nodematch("male", diff = F)`

- Iterations: 5 out of 20

- Monte Carlo MLE Results:


	Estimate	Std. Error	MCMC %	p-value
edges	-2.1552	0.1437	0	<1e-04 ***
nodematch.male	0.7898	0.1794	0	<1e-04 ***

**WARNING:** Currently, the ERGMM procedure in R can only estimate models in which edges are conditionally independent. So cannot include network statistics involving mutual or triadic terms!

# Euclidean distance with $d = 3$ dimensions and 3 clusters (or groups)

```
lmodel7 <- ergmm(pnet ~  
  receiver+rsender+euclidean(d=3,G=3)+nodematch("male",diff=T))
```

Obvious  
application is to  
clusters actors  
into groups!




# Assessment of Goodness of fit


- `lmodel7i.gof <-  
gof(lmodel7, GOF=~idegree, control=ergmm.control(nsim=  
100), verbose=T)`
- `plot(lmodel7i.gof)`
- `lmodel7o.gof <-  
gof(lmodel7, GOF=~odegree, control=ergmm.control(nsim=  
=100), verbose=T)`
- `plot(lmodel7o.gof)`

# Goodness of fit of degree distributions with Euclidean latent-space

Goodness-of-fit diagnostics




Goodness-of-fit diagnostics




# Goodness of fit of degree distributions with Bilinear latent-space

Goodness-of-fit diagnostics



Goodness-of-fit diagnostics



# III. Statistical analyses of social influence or peer effects

- Use Inam function
  - Uses numerical approximation to second derivatives in Newton-family optimization routines
- Develop your own estimation routines
  - Use optim function for maximum likelihood inference
- Amenable to Bayesian inference

# Data wrangling required to form peer actor outcome or attribute weighted average outcome

## Data manipulation

- `on <- as.vector(rep(1,nr))`
- `x <- as.matrix(cbind(on,regdata[,c("male",  
"pctwom","numalters")]))`

## Peer outcome predictor

- `hrtalt <- wtreldir %*% as.vector(covdata$sumhrt)`
- `regdata <-  
data.frame(covdata,hrtalt=hrtalt,noalters=noalters,numalters=numalters)`



# Estimation of linear regression and autoregressive outcome models for cross-sectional network influence data in R using Inam function

## Autoregressive outcome model estimation in R

- `reg.adj <- lm(sumhrt~x+hrtalt-1, data=regdata)`
- `Inam1.adj <- Inam(regdata$sumhrt,x,wtreldir)`

## Network autocorrelation model estimation in R

- `Inam2.adj <- Inam(regdata$sumhrt,x,NULL,wtreldir)`

# Results and Peer-effect Estimate Interpretation (isoequal=0; assume isolates not influenced)

	Linear regression			Autoregressive outcome model		
Term	Estimate	SE	P-value	Estimate	SE	P-value
on	16.049	4.236	0.001	16.078	3.902	0.000
male	-0.327	2.492	0.897	-0.206	2.277	0.928
pctwom	-0.054	0.052	0.305	-0.052	0.047	0.275
numalters	-0.363	0.218	0.107	-0.353	0.199	0.077
hrtalt or rho1.1	0.299	0.195	0.136	0.277	0.171	0.106

Peer effect parameter  
under the autoregressive  
outcome model estimated  
by lnam

# Estimation of network autocorrelation model in R and Interpretation of Peer-Effect Estimates

```
Inam2.adj <- Inam(regdata$sumhrt,x,NULL,wtreldir)
```

Term	Autoregressive outcome model			Network autocorrelation model		
	Estimate	SE	P-value	Estimate	SE	P-value
on	16.078	3.902	0.000	16.471	4.089	0.000
male	-0.206	2.277	0.928	1.324	2.159	0.540
pctwom	-0.052	0.047	0.275	-0.023	0.050	0.645
numalters	-0.353	0.199	0.077	-0.209	0.227	0.356
rho1.1 or rho2.1	0.277	0.171	0.106	0.026	0.445	0.953

Inam also accommodates models with both autoregressive outcome and network autocorrelation terms

Much  
larger SE

# References: Descriptive network measures

- Wasserman S. and Faust K. (1994), Social Network Analysis. Cambridge: Cambridge University Press
- Bonacich, P. (1987), Power and Centrality: A Family of Measures," American Journal of Sociology, 92, 1170-1182
- Freeman, L. (1979), Centrality in Social Networks, Conceptual Clarification, Social Networks, 1, 215-239
- Faust, K. (1997), Centrality in Affiliation Networks, Social Networks, 19, 157-191
- O'Malley, A. J. and Marsden, P. V. (2008), The Analysis of Social Networks, Health Services & Outcomes Research Methodology, 8, 222-269

# References: Bipartite Networks and Comparison of Multiple Networks

- Borgatti, S. and Everett, M. (1997), Network Analysis of 2-Mode Data," Social Networks, 19, 243-269
- Barnett, M. L., Christakis, N. A., O'Malley, A. J., Onnela, J.-P., Keating, N. L., and Landon, B. E. (2012), Physician Patient-Sharing Networks and the Cost and Intensity of Care in US Hospitals," Medical Care, 50, 152-160
- Landon, B. E., Keating, N. L., Barnett, M. L., Onnela, J. P., Paul, S., O'Malley, A. J., Keegan, T., and Christakis, N. A. (2012), Variation in Patient-Sharing Networks of Physicians Across the United States, Journal of the American Medical Association, 308, 265-273.

# References: Bipartite Networks and Comparison of Multiple Networks cont.

- An C, O'Malley AJ, Rockmore DN, Stock CD. Analysis of the U.S. Patient Referral Network. *Statistics in Medicine*, 2018, 37, (5), 847-866. doi: 10.1002/sim.7565. PMID: 29205445
- Moen EL, Bynum JPW, Austin AM, Chakraborti G, Skinner JS, O'Malley AJ. Assessing variation in implantable cardioverter defibrillator therapy guideline adherence with physician and hospital patient-sharing networks. *Medical Care*, 2018, 56, (4), 350-357
- Moen EL, Austin AM, Bynum JP, Skinner JS, O'Malley AJ. An analysis of patient-sharing physician networks and implantable cardioverter defibrillator therapy. *Health Services and Outcomes Research Methodology*, 2016, 16, 132-153
- Landon BE, Keating NL, Onnela J-P, Zaslavsky AM, Christakis NAC, O'Malley AJ. Patient-Sharing Networks of Physicians and Healthcare Utilization and Spending Among Medicare Beneficiaries. *JAMA Internal Medicine*, 2018, 178 (1), 66-73
- Keating NL, O'Malley AJ, Onnela J-P, Landon BE. Assessing the impact of colonoscopy complications on use of colonoscopy among primary care physicians and other connected physicians: an observational study of older Americans. *BMJ Open*, 7, (6), e014239. doi:10.1136/bmjopen-2016-014239
- Moen EL, Bynum JPW, Skinner JS, O'Malley AJ. Physician network position and patient outcomes following implantable cardioverter defibrillator therapy. *Health Services Research*, 2019, 54 (4), 880-889. PMID: 30937894

# References: General Network Analyses and Applications

- Barabasi, A.-L. and Albert, R. (1999), Emergence of Scaling in Random Networks," *Science*, 286, 509-512
- Keating, N. L., Ayanian, J. Z., Cleary, P. D., and Marsden, P. V. (2007), Factors affecting influential discussions among physicians: a social network analysis of a primary care practice," *Journal of general internal medicine*, 22, 794-798
- Coleman, J., Katz, E., and Menzel, H. (1957), The diffusion of innovations among physicians," *Sociometry*, 20, 253-270
- Coleman, J., Katz, E., and et al. (1966), *Medical Innovation: A Diffusion Study*, Bobbs-Merrill
- Pollack, C. E., Soulos, P. R. & Gross, C. P. Physician's peer exposure and the adoption of a new cancer treatment modality. *Cancer* **121**, 2799–2807 (2015).
- Hidalgo, C. A., Blumm, N., Barabasi, A.-L., and Christakis, N. A. (2009), A Dynamic Network Approach for the Study of Human Phenotypes, *PLoS Computational Biology*, 5, e1000353. doi:10.1371/journal.pcbi.1000353

# References: cross-sectional models of relationships (social selection)

- Erdos, P. and Renyi, A. (1959), Random Graphs, Publicationes Mathematicae, 6, 290:297.
- Wang, W. and Wong, G. (1987), Stochastic Blockmodels for Directed Graphs, Journal of the American Statistical Association, 82, 8-19.
- Choi D, Wolfe P. and Airolidi E. (2010) Stochastic blockmodels with growing number of classes. arXiv:1011.4644.
- Fienberg, S. and S. Wasserman. (1981). Categorical data analysis of single sociometric relations. In Sociological Methodology, edited by S. Leinhardt, pp. 156-92. San Francisco: Jossey-Bass.
- Frank, O. and D. Strauss. (1986). Markov graphs. Journal of the American Statistical Association 81: 832-42.
- Goldenberg, A., Zheng, A. X., Fineberg, S. E. and Airolidi, E. M. (2009). A survey of statistical network models. In press: Foundations and Trends in Machine Learning.
- Handcock, M. S., Robins, G. L., Snijders, T. A. B., Moody, J. and Besag, J. (2003). Assessing degeneracy in statistical models of social networks. Journal of the American Statistical Association, 76: 33-50.
- Holland, P. and S. Leinhardt. (1981). An exponential family of probability-distributions for directed-graphs. Journal of the American Statistical Association 76(373): 33-50.



# References: cross-sectional models of relationships cont.

- Handcock, M. S., Hunter, D. R., Butts, C. T., Goodreau, S. M., Krivitsky, P. N., and Morris, M. (2010), *ergm: A Package to Fit, Simulate and Diagnose Exponential-Family Models for Networks*, <http://CRAN.R-project.org/package=ergm>. Version 2.2-6. Project home page at <http://statnetproject.org>
- Handcock, M. S., Robins, G. L., Snijders, T. A. B., Moody, J., and Besag, J. (2003), Assessing degeneracy in statistical models of social networks, *Journal of American Statistical Association*, 76, 33-50
- Robins, G. L., Snijders, T. A. B., Wang, P., Handcock, M. S., and Pattison, P. E. (2007), Recent developments in exponential random graph ( $p^*$ ) models for social networks, *Social Networks*, 29, 192-215
- Hoff, P. (2005). Bilinear mixed-effects models for dyadic data. *Journal of the American Statistical Association* 100: 286-95.
- Hoff, P., A. Raftery, and M. Handcock. (2002). Latent space approaches to social network analysis. *Journal of the American Statistical Association* 97: 1090-98.
- Hoff P. (2008), Modeling Homophily and Stochastic Equivalence in Symmetric Relational Data, in: *Advances in Neural Information Processing Systems*, volume 20, MIT Press, 657-664
- Paul S, Keating NL, Landon BE, O'Malley AJ. Results from using a new dyadic-dependence model to analyze sociocentric physician networks. *Social Science & Medicine*, 125, 2015, 51-59

# References: dynamic network models of relationships

- Krivitsky P.N. and Handcock M.S. (2010). A Separable Model for Dynamic Networks. arXiv:1011.1937v1[stat.ME].
- Nowicki K and Snijders T.A.B. (2001). Estimation and prediction for stochastic blockstructures. *Journal of the American Statistical Association*, 96, 1077-1087.
- O'Malley AJ, Christakis NA. (2011). Longitudinal Analysis of Large Social Networks: estimating the Effect of Health Traits on changes in Friendship Ties. *Statistics in Medicine*, 30, 9, 950-964
- Paul S. and O'Malley A.J. (2013). Hierarchical longitudinal models of relationships in social networks. *Journal of the Royal Statistical Society, Series C*.
- Pattison, P. and S. Wasserman. (1999). Logit models and logistic regressions for social networks: II. Multivariate relations. *British Journal of Mathematical and Statistical Psychology* 52 (Pt 2): 169-93.
- Van Duijn, M., T. Snijders, and B. Zijlstra. (2004). A Random Effects Model with Covariates for Directed Graphs. *Statistica Neerlandica* 58(2): 234-54.

# References: dynamic network models of relationships cont.

- Snijders, T. A. B. (2006). "Statistical methods for network dynamics." In S. R. Luchini et al., editors, Proceedings of the XLIII Scientific Meeting, Italian Statistical Society, pages 281-296, Padova: CLEUP.
- Snijders, T.A.B. (2005), Models for longitudinal social network data," in Models and Methods in Social Network Analysis, Cambridge University Press, 215-247.
- Steglich, C.E.G., Snijders, T.A.B. and Pearson, M. (2010). Dynamic Networks and Behavior: Separating Selection from Influence. Sociological Methodology, 40, 329-393.
- Westveld, A. H. and Hoff, P. D. (2011), A Mixed Effect Model for Longitudinal Relational and Network Data, With Applications To International Trade and Conflict," The Annals of Applied Statistics, 5, 843-872.

# References: social influence analyses

- Christakis NA, Fowler JH. (2007). The Spread of Obesity in a Large Social Network over 32 Years. *New England Journal of Medicine*, 357, 370--379
- Christakis NA, Fowler, JH. (2008). Dynamics of Smoking Behavior in a Large Social Network. *New England Journal of Medicine*, 358, 2249--2258
- Fowler JH, Christakis NA. (2008). Dynamic spread of happiness in a large social network: longitudinal analysis over 20 years in the Framingham Heart Study. *British Medical Journal*, 337, doi:10.1136/bmj.a2338.
- Cohen-Cole E, Fletcher JM. (2008). Detecting Implausible Social Network Effects in Acne, Height, and Headaches: Longitudinal Analyses. *British Medical Journal*, 337, a2533.
- Lyons, R. (2011), The spread of evidence-poor medicine via flawed social network analyses, *Statistics, Politics and Policy*, 2, 1-26, doi:10.2202/2151-7509.1024
- Shalizi CR, Thomas AC. (2011). Homophily and Contagion Are Generically Confounded in Observational Social Network Studies, *Sociological Methods and Research*, 40, 211--239.
- O'Malley AJ, Elwert F, Rosenquist JN, Zaslavsky AM, Christakis NA. Estimating peer effects in longitudinal dyadic data using instrumental variables. *Biometrics*, 2014, 70, 3, 506--515. (Published online: 29 APR 2014, DOI:10.1111/biom.12172).

# References: social influence analyses cont.

- Christakis, N. A. and Fowler, J. H. (2013), Social Contagion Theory: Examining Dynamic Social Networks and Human Behavior," *Statistics in Medicine*, 32, 556-577
- Manski, C. A. (1993), Identification of endogenous social effects: The Reflection Problem, *Review of Economic Studies*, 60, 531-542
- Marsden, P. V. and Friedkin, N. E. (1993), Network Studies of Social Influence, *Sociological Methods and Research*, 22, 127-151
- McPherson, M. L., Smith-Lovin, Cook, and et al (2001), Birds of a Feather: Homophily in Social Networks, *Annual Review of Sociology*, 27, 415-444
- VanderWeele, T. J. (2011), Sensitivity Analysis for Contagion Effects in Social Networks, *Sociological Methods & Research*, 40, 240-255
- VanderWeele, T. J., Ogburn, E. L., and Tchetgen Tchetgen, E. J. (2012), Why and When "Flawed" Social Network Analyses still yield Valid Tests of no Contagion, *Statistics, Politics, and Policy*, Manuscript 1050
- Keating NL, O'Malley AJ, Onnela J-P, Gray SQ, Landon BE. Influence of Peer Physicians on Intensity of End-of-Life Care for Cancer Decedents. *Medical Care*, 57, 6, 468-474. PMID: 31033059

# References: network-influence related models

- Land KC, Deane G. (1992). On the Large-Sample Estimation of Regression Models with Spatial or Network Effect Terms: A Two-Stage Least-Squares Approach. *Sociological Methodology*, (ed: Peter V. Marsden), Oxford, UK: Basil Blackwell, Ltd, 221-248
- Anselin L. (1988). *Spatial Econometrics: Methods and Models*, Kluwer Academic Publishers: Dordrecht, The Netherlands
- Sargan JD. (1958). The Estimation of Econometric Relationships Using Instrumental Variables, *Econometrica*, 26, 393-415
- Haining, R.P. (1978). The moving average model for spatial interaction. *Transactions of the Institute of British Geographers*, 3, 202-225
- Kelejian, H.H., Robinson, D.P., 1993. A suggested method of estimation for spatial interdependent models with autocorrelated errors, and an application to a county expenditure model. *Papers in Regional Science* 72, 297-312
- O'Malley AJ, Moen EL, Bynum JPW, Austin AM, Skinner JS. Modeling Peer Effect Modification by Network Position: The Diffusion of Implantable Cardioverter Defibrillators in the US Hospital Network. In Press: *Statistics in Medicine*